



Effect of NPK and Zinc Inorganic Fertilizers on Soil Health, Growth and Yield of Chickpea (*Cicer arietinum* L.)

Lokendra Meena^a, Ram Bharose^a, Arun Alfred David^a,
Tarence Thomas^a, Neha Toppo^{a*} and Harsh Kumar^a

^a Department of Soil Science and Agricultural Chemistry, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj - 211 007, U.P, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i173194

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/101405>

Original Research Article

Received: 15/04/2023

Accepted: 17/06/2023

Published: 04/07/2023

ABSTRACT

An objective of study effect on inorganic sources of NPK and Zn fertilizers on soil health with the experiment was laid out in Randomized Block Design with three levels of NPK (00%, 50% and 100%), and Zinc (00%, 50% and 100%) Recommended Dose of fertilizer applied in all treatments. The result showed that the soil properties, growth and yield of chickpea were significantly affected by application of NPK and Zinc. The highest growth and yield were observed in T₉ (100%NPK +100% Zinc), followed by T₈ (100% NPK and +50%Zinc), whereas the lowest growth and yield were observed in T₁ (00% NPK +00% Zinc). The treatment T₉ (NPK @ 100 % +Zn @ 100%) was recorded as best treatment for major soil parameters. The treatment T₉ (NPK @ 100% + Zn @ 100 %) also shows the significantly highest vegetative growth as well as yield attributes and net profit ₹45,186.00 ha⁻¹ with cost benefit ratio is 1:2.17.

Keywords: Chickpea; NPK; zinc; growth and yield; physico-chemical properties of soil.

*Corresponding author: E-mail: neha.shrishty@gmail.com;

1. INTRODUCTION

Soil plays a critical role in chickpea production by providing nutrients, water and physical support for growth. Physico-chemical properties of soil, including texture, pH, organic matter, fertility, moisture, drainage and temperature, play critical role in growth and development of chickpea. Proper management of these properties can lead to a healthy and productive chickpea crop. "Soil is the loose surface material that covers most land. It consists of inorganic particles and organic matter. Soil provides the structural support to plants used in agriculture and is also their source of water and nutrients. Soils vary greatly in their chemical and physical properties. Processes such as leaching, weathering and microbial activity combine to make a whole range of different soil types. Each type has strengths and weaknesses for agricultural production" [1].

"The three most significant nutrients, without any one of which plants could not endure, are known as the primary macronutrients: Nitrogen (N), Phosphorus (P), and Potassium (K). In particular, nitrogen is crucial to chlorophyll, which allows plants to carry out photosynthesis (the process by which they uptake sunlight to produce sugars from carbon dioxide and water). Nitrogen is also a major component in amino acids, the base of proteins" [2].

Nitrogen also aids in the compounds that let for storage and use of energy. The plants that are nutritious and healthy while developing have good nitrogen content in them. Phosphorus also encourages the growth of roots, promotes flowering, and is necessary for DNA. Phosphorus fertilizer extract from the phosphate rock [3].

Phosphorus is a vital component of ATP, the "energy unit" of plants. "ATP forms during photosynthesis, has phosphorus in its structure, and processes from the beginning of seedling growth through to the formation of grain and maturity. Thus, phosphorus is essential for the general health and vigour of all plants" [2].

Potassium is often referred to as the "value element," because of its contribution to many of the features we connect with quality, such as size, figure, color, and even taste, among others. Potassium provides strength to the plants and can resist diseases [2].

"Zinc is an important component of various enzymes that are responsible for driving many metabolic reactions in all crops. Growth and

development would stop if specific enzymes were not present in plant tissue. Carbohydrate, protein, and chlorophyll formation is significantly reduced in zinc-deficient plants" [4].

2. MATERIALS AND METHODS

A field experiment conducted at the Soil Science Research Farm, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, during the *Rabi* season 2021-22 growing chickpea *Var. Sadabahar* applied 3 levels of NPK and Zinc respectively 0 %, 50 % and 100 % including RDF for chickpea = 20:40:20 kg ha⁻¹ experiment is lead to observe the physical and chemical parameters. In physical parameters like that bulk density, particle density, pore space and water holding capacity through method by 100 ml graduated measuring cylinder and process by Muthuvel et al. [5]. "In chemical parameters through method by-Soil pH – method given by Jackson ML. [6] through using digital pH meter, Soil EC (dSm⁻¹) - method given by Wilcox, [7] through using digital EC meter, Organic Carbon (%)- Wet oxidation method given by Walkley and Black, [8], Available Nitrogen (kg ha⁻¹) - Kjeldhal Method [9], Available Phosphorus (kg ha⁻¹) - Colorimetric method by using Jasper single beam U.V. Spectrophotometer at 660 nm wavelength given by Olsen et al., [10], Available Potassium (kg ha⁻¹) - Flame photometric method by using Metzer Flame Photometer given by Toth and Prince, [11], Available Zinc (mg kg⁻¹) – DTPA extractant method by using AAS given by" Lindsay and Norvell, [12].

3. RESULTS AND DISCUSSION

3.1 Bulk Density (Mg m⁻³)

The data presented in Table and Fig. 1 shows the effect of NPK and Zinc on bulk density (Mg m⁻³) of soil after crop harvested. The response of NPK and Zinc on bulk density of soil was found non-significant. The maximum bulk density of soil 1.33 and 1.36 Mg m⁻³ was found at 0-15 and 15-30 cm in treatment T₁ (NPK @ 0 % + Zn @ 0 %) and minimum bulk density of soil 1.15 and 1.17 Mg m⁻³ was found at 0-15 and 15-30 cm in treatment T₉ (NPK @ 100 % + Zn @ 100 %) respectively. Similar result has been recorded by Hussain et al., [13]; Chinthra et al., [14].

3.2 Particle Density (Mg m⁻³)

The data presented in Table and Fig. 1 shows the effect of NPK and Zinc on particle density

(Mg m^{-3}) of soil after crop harvested. The response of NPK and Zinc on particle density of soil was found non-significant. The maximum particle density of soil 2.52 and 2.55 Mg m^{-3} was recorded at 0-15 and 15-30 cm in treatment T_1 (NPK @ 0 % + Zn @ 0 %) and minimum particle density of soil 2.38 and 2.45 Mg m^{-3} was recorded at 0-15 and 15-30 cm in treatment T_9 (NPK @ 100 % + Zn @ 100 %) respectively. Similar result has been recorded by Hussain et al., [13]; Chinthia et al., [14] and Dangi et al., [15].

3.3 Percent Pore Space (%)

The data presented in Table and Fig. 1 shows the effect of NPK and Zinc on percent pore space of soil after crop harvested. The response of NPK and Zinc on percent pore space of soil was found significant. The maximum percent pore space of soil 48.7 and 47.60 % was found at 0-15 and 15-30 cm in treatment T_9 (NPK @ 100 % + Zn @ 100 %) and minimum percent pore space of soil 45.11 and 44.77 % was found at 0-15 and 15-30 cm in treatment T_1 (NPK @ 0 % + Zn @ 0 %) respectively. Similar result has been recorded by Hussain et al., [13]; Chinthia et al., [14] and Dangi et al., [15].

3.4 Water Holding Capacity (%)

The data presented in Table and Fig. 1 shows the effect of NPK and Zinc on water holding capacity (%) of soil after crop harvested. The response of NPK and Zinc on water holding capacity of soil was found significant. The maximum water holding capacity of soil 44.58 and 42.09 % was found at 0-15 and 15-30 cm in treatment T_9 (NPK @ 100 % + Zn @ 100 %) and minimum water holding capacity of soil 39.23 and 38.05 % was found at 0-15 and 15-30 cm in treatment T_1 (NPK @ 0 % + Zn @ 0 %) respectively. Similar result has been recorded by Hussain et al., [13]; Chinthia et al., [14] and Dangi et al., [15].

3.5 Soil pH (1:2.5) w/v

The data presented in Table and Fig. 2 shows the effect of NPK and Zinc on pH of soil after crop harvested. The response NPK and Zinc on pH of soil was found non-significant. The maximum pH of soil 7.36 and 7.41 was found at 0-15 and 15-30 cm in treatment T_1 (NPK @ 0 % + Zn @ 0 %) and minimum pH of soil 7.27 and 7.27 was found at 15-30 cm in treatment T_9 (NPK @ 100 % + Zn @ 100 %) respectively. Similar result has been recorded by Parmar and Poonia [16]; Sahu et al., [17] and Deshlahare et al., [18].

3.6 Soil EC (dS m^{-1})

The data presented in Table and Fig. 2 shows the Effect of NPK and Zinc on EC of soil after crop harvested. The response of NPK and Zinc on EC of soil was found non-significant. The maximum EC of soil 0.476 and 0.482 dSm^{-1} was recorded at 0-15 and 15-30 cm in treatment T_9 (NPK @ 100% + Zn @ 100 %) and minimum EC of soil 0.445 and 0.448 dSm^{-1} was recorded at 15-30 cm in treatment T_1 [control (NPK @ 0% + Zn @ 0 %)] respectively. Similar result has been recorded by Parmar and Poonia [16]; Sahu et al., [17] and Deshlahare et al., [18].

3.7 Organic Carbon (%)

The data presented in Table and Fig. 2 shows the effect of NPK and Zinc on organic carbon of soil after crop harvested. The response of NPK and Zinc on organic carbon of soil was found non-significant. The maximum organic carbon of soil 0.43 and 0.38 % was found at 0-15 and 15-30 cm in treatment T_9 (NPK @ 100 % + Zn @ 100 %) and minimum organic carbon of soil 0.36 and 0.32 % was found at 0-15 and 15-30 cm in treatment T_1 (NPK @ 0 % + Zn @ 0 %) respectively. Similar result has been recorded by Parmar and Poonia [16]; Sahu et al., [17] and Deshlahare et al., [18].

3.8 Available Nitrogen (kg ha^{-1})

The data presented in Table and Fig. 2 shows the effect of NPK and Zinc on available nitrogen of soil after crop harvested. The response of NPK and Zinc on available nitrogen of soil was found significant. The maximum available nitrogen of soil 322.75 and 316.59 kg ha^{-1} was recorded at 0-15 and 15-30 cm in treatment T_9 (NPK @ 100% + Zn @ 100 %) and minimum available nitrogen of soil 290.65 and 286.32 kg ha^{-1} was recorded at 0-15 and 15-30 cm in treatment T_1 [control (NPK @ 0% + Zn @ 0 %)] respectively. Similar result has been recorded by Pingoliya et al., [19]; Yadav et al., [20] and Banjara and Majgahe [21].

3.9 Available Phosphorus (kg ha^{-1})

The data presented in Table and Fig. 2 shows the effect of NPK and Zinc on available phosphorus of soil after crop harvested. The response of NPK and Zinc on available phosphorus of soil was found significant. The maximum available phosphorus of 35.07 and

33.82 kg ha⁻¹ was recorded at 0-15 and 15-30 cm in treatment T₉ (NPK @ 100% + Zn @ 100 %) and minimum available phosphorus of soil 19.40 and 16.43 kg ha⁻¹ was recorded at 0-15 and 15-30

cm in treatment T₁ (NPK @ 0% + Zn @ 0 %) respectively. Similar result has been recorded by Pingoliya et al., [19] Yadav et al., [20]; Banjara and Majgahe [21].




Fig. 1. Effect of NPK and Zn on bulk density (Mg m⁻³), particle density (Mg m⁻³), pore space (%) and water holding capacity (%) of soil after crop harvest




Fig. 2. Effect of NPK and Zn on pH, electrical conductivity (dSm⁻¹), organic carbon (%), available nitrogen (kg ha⁻¹), available phosphorus (kg ha⁻¹), available potassium (kg ha⁻¹) and available zinc (mg kg⁻¹) of soil after crop harvest

Table 1. Effect of NPK and Zn on bulk density (Mg m^{-3}), particle density (Mg m^{-3}), pore space (%) and water holding capacity (%) of soil after crop harvest

Treatment	Bulk density (Mg m^{-3})		Particle density (Mg m^{-3})		Pore space (%)		Water holding capacity (%)	
	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm
T ₁ NPK @ 0 % + Zn @ 0 %	1.33	1.36	2.52	2.55	45.11	44.77	39.23	38.05
T ₂ NPK @ 0 % + Zn @ 50 %	1.30	1.32	2.50	2.53	45.97	45.80	39.03	38.64
T ₃ NPK @ 0 % + Zn @ 100 %	1.28	1.31	2.48	2.51	46.24	45.88	41.96	39.26
T ₄ NPK @ 50 % + Zn @ 0 %	1.27	1.29	2.51	2.54	45.55	45.11	39.76	38.29
T ₅ NPK @ 50 % + Zn @ 50 %	1.24	1.26	2.47	2.52	47.05	46.87	41.51	39.83
T ₆ NPK @ 50 % + Zn @ 100 %	1.21	1.25	2.44	2.49	47.38	47.07	42.27	41.87
T ₇ NPK @ 100 % + Zn @ 0 %	1.20	1.23	2.46	2.51	46.48	45.52	41.89	40.34
T ₈ NPK @ 100 % + Zn @ 50 %	1.18	1.21	2.42	2.48	47.60	47.52	43.05	41.25
T ₉ NPK @ 100 % + Zn @ 100 %	1.15	1.17	2.38	2.45	48.71	47.60	44.58	42.09
F-Test	NS	NS	NS	NS	S	S	S	S
S.Ed. (\pm)	-	-	-	-	0.61	0.59	0.68	0.79
C.D. at 0.5%	-	-	-	-	1.84	1.76	2.06	2.39

Table 2. Effect of NPK and Zn on pH, electrical conductivity (dSm^{-1}), organic carbon (%), available nitrogen (kg ha^{-1}), available phosphorus (kg ha^{-1}), available potassium (kg ha^{-1}) and available zinc (mg kg^{-1}) of soil after crop harvest

Treatment	Soil pH (1:2.5) w/v		Electrical Conductivity (dSm^{-1})		Organic Carbon (%)		Available Nitrogen (kg ha^{-1})		Available Phosphorus (kg ha^{-1})		Available Potassium (kg ha^{-1})		Available Zinc (mg kg^{-1})	
	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm
T ₁ NPK @ 0 % + Zn @ 0 %	7.36	7.41	0.445	0.448	0.36	0.32	290.65	286.32	19.40	16.43	182.23	178.55	0.284	0.292
T ₂ NPK @ 0 % + Zn @ 50 %	7.36	7.40	0.449	0.451	0.36	0.33	291.45	287.05	20.26	17.87	184.41	181.82	0.328	0.334
T ₃ NPK @ 0 % + Zn @ 100 %	7.35	7.40	0.453	0.456	0.38	0.34	293.37	289.91	22.87	19.09	187.58	185.56	0.288	0.296
T ₄ NPK @ 50 % + Zn @ 0 %	7.32	7.38	0.456	0.459	0.40	0.36	297.72	291.65	23.50	20.60	188.08	186.72	0.321	0.329
T ₅ NPK @ 50 % + Zn @ 50 %	7.31	7.38	0.460	0.465	0.41	0.38	300.68	294.27	25.24	22.28	191.56	189.80	0.331	0.338
T ₆ NPK @ 50 % + Zn @ 100 %	7.31	7.37	0.464	0.469	0.41	0.39	304.80	298.53	28.16	26.54	194.78	192.45	0.337	0.342
T ₇ NPK @ 100 % + Zn @ 0 %	7.28	7.31	0.467	0.473	0.42	0.39	310.06	304.26	30.45	28.27	199.81	195.72	0.340	0.344
T ₈ NPK @ 100 % + Zn @ 50 %	7.28	7.30	0.472	0.477	0.42	0.39	317.35	309.38	33.71	31.20	202.95	198.65	0.342	0.348
T ₉ NPK @ 100 % + Zn @ 100 %	7.27	7.29	0.476	0.482	0.43	0.38	322.75	316.59	35.07	33.82	208.42	204.67	0.345	0.353
F-Test	NS	NS	NS	NS	NS	NS	S	S	S	S	S	S	S	S
S.Ed. (\pm)	-	-	-	-	-	-	2.27	1.84	1.08	0.78	1.57	1.35	0.07	0.08
C.D. at 0.5%	-	-	-	-	-	-	4.62	3.71	2.19	1.61	2.25	1.78	0.16	0.19

3.10 Available Potassium (kg ha⁻¹)

The data presented in Table and Fig. 2 shows the effect of NPK and Zinc on available potassium of soil after crop harvested. The response of NPK and Zinc on available potassium of soil was found significant. The maximum available potassium of soil was recorded 208.42 and 204.67 kg ha⁻¹ was recorded at 0-15 and 15-30 cm in treatment T₉ (NPK @ 100% + Zn @ 100 %) and minimum available potassium of soil 182.23 kg and 178.55 kg ha⁻¹ was recorded at 0-15 and 15-30 cm in treatment T₁ [control (NPK @ 0% + Zn @ 0 %)] respectively. Similar result has been recorded by Pingoliya et al., [19], Yadav et al., [20]; Banjara and Majgahe [21].

3.11 Available Zn (mg kg⁻¹)

The data presented in Table and Fig. 2 shows the effect of NPK and Zinc on available Zn of soil after crop harvested. The response of NPK and Zinc on available Zn of soil was found significant. The maximum available Zn of soil 0.345 and 0.353 mg kg⁻¹ was recorded at 0-15 and 15-30 cm in treatment T₉ (NPK @ 100% + Zn @ 100 %) and minimum available Zn of soil 0.284 and 0.292 mg kg⁻¹ was recorded at 0-15 and 15-30 cm in treatment T₁ [control (NPK @ 0% + Zn @ 0 %)] respectively. Similar result has been recorded by Pingoliya et al., [19] Yadav et al., [20]; Banjara and Majgahe [21], [22-25].

4. CONCLUSION

The results of the experiment are concluded as the effect of NPK and Zinc on nitrogen (kg ha⁻¹), phosphorus (kg ha⁻¹), potassium (kg ha⁻¹), % pore space and water holding capacity (%) of soil after crop harvest was found significant except on bulk density (Mg m⁻³), particle density (Mg m⁻³), pH, EC (dsm⁻¹) and organic carbon (%) of soil after harvest. The treatment T₉ (NPK @ 100 % +Zn @ 100%) was recorded as best treatment for major soil parameters. Therefore, it can be recommended for farmers to obtain best combination Treatment (T₉) for higher farm income and sustainable agriculture.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Roy RN, Finck A, Blair GJ, Tandon HLS. Plant nutrition for food security. A guide for integrated nutrient management. FAO fertilizer and Plant Nutrition Bulletin 16. Rome, Italy, food and Agriculture Organization of the United Nations; 2006.
- Mishra PK, Bisht SC, Pooja R, Joshi GK, Singh G, Bisht JK, Bhatt JC. Bio associative effect of cold tolerant pseudomonas spp. and rhizobium leguminosarum-PR 1 on Iron acquisition, nutrient uptake and growth of lentil (*Lens culinaris* L.). European Journal of Soil Biology. 2011;47:35-43.
- Dixit GP. Project coordinator's report, all india coordinated research project on Chickpea. Indian Institute of Pulses Research, Kanpur; 2016.
- Singh A, Singh VK, Chandra R, Srivastava PC. Effect of the integrated nutrient management on chickpea based intercropping system and soil properties in Mollisols of the Tarai region. Journal of the Indian Society of Soil Science. 2009;60:38-44.
- Muthuvel P, Udayasoorian C, Natesan R, Ramaswamy PP. Introduction to Soil Analysis, Tamil Nadu Agricultural University Coimbatore-641002; 1992.
- Jackson ML. Soil chemical analysis Prentice Hall of India Ltd. New Delhi. 1958:219-221.
- Wilcox LV. Electrical conductivity Am. water work, Association. 1950;42:775-776.
- Walkley A, Black IA. Estimation of soil organic carbon by the chromic acid titration method. Soil Science. 1947;47:29-38.
- Subbiah BV, Asijja EC. A rapid procedure for estimation of available nitrogen in soil. Current Science. 1956;25(8):259-260.
- Olsen SR, Cole CV, Watanabe FS, Dean LA. Estimation of available phosphorus in soils by extraction with sodium bicarbonate (NaHCO₃). U.S.D.A. Circular. 1954;939:1-19.
- Toth SJ, Prince AL. Estimation of cation exchange capacity and exchangeable Ca, K and Na content of soil by flame photometer technique. Soil Sci. 1949;67:439-445.
- Lindsay WL, Norvell WA. Development of DTPA soil test for zinc, iron, manganese, and copper. Soil Science Society of America Journal. 1978;42:421-428.

13. Hussain M, Banoo M, Sinha BK, Chand G. Effect of foliar application of zinc and boron on growth, yield and quality attributes in chickpea (*Cicer arietinum* L.). Journal of Pharmacognosy and Phytochemistry. 2022;11(3):270-275.
14. Chintha HB, Swaroop N, Thomas T, Barthwal A, Harit H, Amjad A. Effect of nitrogen, phosphorus, potassium, sulphur and zinc on soil health parameters of chickpea (*Cicer arietinum* L.). The Pharma Innovation Journal. 2021;10(11):1955-1961.
15. Dangi M, Thomas T, David AA, Kumar S, Joshi A. Response of different levels of zinc and FYM on soil health and yield attributes of chickpea (*Cicer arietinum* L.). International Journal of Chemical Studies. 2020;8(4):3559-3563.
16. Parmar PM. Poonia TC. Effect of zinc biofortification on growth, yield and economics of chickpea (*Cicer arietinum* L.). International Journal of Chemical Studies. 2020;8(2):1782-1786.
17. Sahu A, Swaroop N, David AA, Thomas T. Effect of different levels of NPK and Zinc on soil health growth and yield of chickpea (*Cicer arietinum* L.) var. PUSA 362. Int. J. Curr. Microbiol. App. Sci. 2020;9(10):591-597.
18. Deshlahare H, Banjara GP. Impact of biofortification of zinc and iron on growth parameters and yields of chickpea (*Cicer arietinum* L.) through agronomic intervention in Chhattisgarh plains. Journal of Pharmacognosy and Phytochemistry. 2019;8(6):383-386.
19. Pingoliya KK, Dotaniya ML, Lata M. Effect of iron on yield, quality and nutrient uptake of chickpea (*Cicer arietinum* L.). African Journal of Agricultural Research. 2014;9(37):2841-2845.
20. Yadav R, Thomas T, Swaroop N, Kumar T. Response of different levels of phosphorus and zinc on soil health after harvesting chickpea (*Cicer arietinum* L.). The Pharma Innovation Journal. 2021;11(5):402-405.
21. Banjara GP, Majgahe SK. Effect of Biofortification of zinc and iron on yield attributes and yields of chickpea (*Cicer arietinum* L.) through agronomic intervention. The Pharma Innovation Journal. 2019;8(10):45-47.
22. Bouyoucos GJ. The hydrometer as a new method for the mechanical analysis of soils. Soil Science. 1927;23:343-353.
23. Fisher RA. Statistical methods and scientific induction. Journal of the Royal Statistical Society Series. 1955;17:69-78.
24. Munsell AH. Munsell's description of his colour system, from a lecture to the American Psychological Association. American Journal of Psychology. 1971;23(2):236-244.
25. Pongener P, David AA, Thomas T, Hasan A, Reddy IS. Effect of organic and inorganic source of nutrients on physio-chemical properties of soil in black gram (*Vigna mungo* L.) Var. Sekhar-2.

© 2023 Meena et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/101405>