



# Response of *Bacillus megaterium* and *Bacillus mucilaginosus* Strains on Growth and Nutrient Uptake of Soybean

Kavita Solanki <sup>a</sup>, S. K. Choudhary <sup>a</sup>, Aakash <sup>a,b\*</sup>,  
Veer Singh <sup>b</sup>, Hrishikesh Nath <sup>c</sup> and Kumar Anshuman <sup>d</sup>

<sup>a</sup> Department of Agronomy, Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, College of Agriculture, Indore 452001, Madhya Pradesh, India.

<sup>b</sup> Department of Agronomy, R.S.M. (P.G.) College, Dhampur-246761 (Bijnor) (Affiliated to Mahatma Jyotiba Phule Rohilkhand University, Bareilly, Uttar Pradesh), India.

<sup>c</sup> Department of Agronomy, Assam Agricultural University, Jorhat 785013, Assam, India.

<sup>d</sup> Department of Soil Science, KNIPSS, Sultanpur 228001, Uttar Pradesh, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i213973

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/108019>

Original Research Article

Received: 15/08/2023

Accepted: 17/10/2023

Published: 17/10/2023

## ABSTRACT

The current study aimed to investigate the Response of *Bacillus megaterium* and *Bacillus mucilaginosus* strains on growth attributes and uptake of soybean (*Glycine max* (L.) Merrill.) at All India Coordinated Research Project (AICRP) on Integrated Farming System Research at College of Agriculture, Indore, Madhya Pradesh, India during *khari*, 2019 & 20. The study was carried out in randomized block design (RBD) with 8 treatments, viz. Control with 75% RDF (T<sub>1</sub>), Control with

\*Corresponding author: E-mail: akash.agro10@bhu.ac.in;

100% RDF (T<sub>2</sub>), 75% RDF with *Bacillus megaterium* and *Bacillus mucilaginosus* strains as seed treatment (T<sub>3</sub>), 100% RDF with *Bacillus megaterium* and *Bacillus mucilaginosus* strains as seed treatment (T<sub>4</sub>), 75% RDF with *Bacillus megaterium* and *Bacillus mucilaginosus* strains as foliar application at 30 and 45 DAS (T<sub>5</sub>), 100% RDF with *Bacillus megaterium* and *Bacillus mucilaginosus* strains as foliar application at 30 and 45 DAS (T<sub>6</sub>), 75% RDF with *Bacillus megaterium* and *Bacillus mucilaginosus* strains as seed treatment + foliar application of *Bacillus megaterium* and *Bacillus mucilaginosus* strains at 30 and 45 DAS (T<sub>7</sub>) and 100% RDF with *Bacillus megaterium* and *Bacillus mucilaginosus* strains as seed treatment + foliar application of *Bacillus megaterium* and *Bacillus mucilaginosus* strains at 30 and 45 DAS (T<sub>8</sub>). Among all the treatments, growth parameters, NPK uptake by grain and straw and total uptake by crop were observed highest with 100% RDF with *Bacillus megaterium* and *Bacillus mucilaginosus* as seed treatment + foliar application of *Bacillus megaterium* and *Bacillus mucilaginosus* at 30 and 45 DAS.

**Keywords:** Soybean; *Bacillus megaterium*; *Bacillus mucilaginosus*; nutrient uptake.

## 1. INTRODUCTION

Soybeans, scientifically known as *Glycine max* (L.) Merrill, are prominent legume crops with high protein and oil content. This remarkable legume also contains a high oil content of 18% to 22.5%, in addition to its protein content of 40.5%. Furthermore, it contains 20-30% extractable substances and is well-balanced in amino acids [1]. In China and Japan, soybeans are called yellow beans and edamame, while in the United States, they are called miracle beans and golden beans [2]. As a result of its ability to adapt to diverse soil conditions and climates, as well as its nutritional value, soybeans have gained popularity in Central India. Soybean occupies approximately 6% of all agricultural land worldwide as the most widely cultivated oilseed crop [3]. As per [4,] soybean cultivation in India covers approximately 11.33 million hectares, and the potential production in 2019-20 is over 13.79 million tonnes. Madhya Pradesh is known as the "Soya State of India" in Central India.

Synthetic fertilizers are used widely, which helps to address some challenges, but also poses risks to the environment and food production, as highlighted by [5,6]. It is possible to address these concerns by harnessing the potential of bacterial species such as *Azotobacter*, *Azospirillum*, *Bacillus* sp., and *Pseudomonas* sp. In general, these microbes are classified as plant growth-promoting rhizobacteria (PGPR). By utilizing these microorganisms as biofertilizers, synthetic fertilizers can be replaced with an alternative that is sustainable and viable [6]. There are numerous bacteria that exhibit remarkable capabilities, including *Azotobacter* and *Azospirillum*. According to [7] they are capable of fixing atmospheric nitrogen and increasing soil phosphorus solubilization. Through processes like BNF (Biological nitrogen

fixation) and phosphate solubilization, plant growth-promoting bacteria (PGPB) promote plant growth as well as reduce stress by modulating 1-aminocyclopropane-1-carboxylate deaminase expression and producing phytohormones and siderophores [8]. In agriculture, microorganisms can be beneficial, and they are now used to produce sustainable food crops. A number of beneficial microorganisms have been shown to play a role in nitrogen fixation, the decomposition of organic wastes, the detoxification of pesticides, the suppression of plant diseases, the enhancement of nutrient cycling, and the production of bioactive compounds that upregulate plant growth [9,6]. A holistic improvement in agriculture is achieved when organic, inorganic, and bio-fertilizers are combined. By integrating soil productivity, sustainability, reclamation, and crop development, setting, and quality, this integrated approach leads to improvements in soil productivity, sustainability, and reclamation [10]. The production of microbial metabolites, including organic acids, can reduce soil pH, thereby facilitating the solubilization of certain nutrients, thus improving plant availability.

Based on these assumptions, the current study focused on evaluating the effects of *Bacillus megaterium* and *Bacillus mucilaginosus* strains on soybean aiming to achieve the following: (1) Evaluation of the effects of *Bacillus megaterium* and *Bacillus mucilaginosus* strains on growth attributes; (2) To find out the effect of *Bacillus megaterium* and *Bacillus mucilaginosus* strains on nutrient uptake study.

## 2. MATERIALS AND METHODS

The experiment was conducted at Research Farm of Rajmata Vijayaraje Scindia Krishna

Vishwa Vidyalaya, College of Agriculture, Indore, Madhya Pradesh, India during *kharif* season (2019–20) with soybean var. “JS 95-60”. The topography of field was uniform with gentle slope. Indore is situated at an altitude of 555.5 m above mean sea level (MSL). It is located at latitude 22.43 °N and longitude of 75.66 °E. This region enjoys sub-tropical, semi-arid type climate. The maximum temperature varied from

25.36°C - 32.43°C and the minimum temperature varied from 20.4°C - 24.57°C. The soil of experimental site was predominantly clayey in texture, slightly alkaline in reaction (pH 7.5) and low in organic carbon (0.45%) and available nitrogen (210 kg·ha<sup>-1</sup>), low in available phosphorus (11.5 kg·ha<sup>-1</sup>) and high in available potassium (410 kg·ha<sup>-1</sup>).




Fig. 1. Rainfall and other weather parameters during crop growing period of experiment




Fig. 2. Experimental location

**Crop management:** Field preparation began with summer plowing with a tractor-drawn plough followed by cross harrowing to prepare the soil for sowing. After harrowing, the field was leveled by planking before sowing. Before planting the seeds, all fertilizers were applied as basal in the furrows and mixed with soil. The seeds used were healthy and of good quality to ensure perfect germination. At the time of sowing, seeds were treated with Bavistin @ 2 g kg<sup>-1</sup> seeds and then inoculated with *Bacillus megaterium* and *Bacillus mucilaginosus* strains @ 3 g kg<sup>-1</sup> seeds. Plants were spaced 30 cm x 5 cm apart in rows with a seed rate of 80 kg ha<sup>-1</sup> in plots with dimensions of 6 m x 4.5 m. In soybean at early stages, two sprays of Triazophos 40 EC 600 ml ha<sup>-1</sup> were applied at 30 and 45 days of crop growth to protect from insects and pests such as girdle beetles, stem fly caterpillars, blue beetles, etc.

**Collection and preparation of soil sample:** We collected soil samples randomly before sowing and after harvesting from each plot, mixed them thoroughly, dried them in air, crushed them, and sieved them through a 2 mm sieve to determine their fertility status.

**Nutrient uptake estimation:** The final soil samples were collected from the experimental plot with the help of Augur to the depth of 15 cm. The soil samples were composited and preserved in clean polythene bag with proper labeling for the determination of the followings.

**Nitrogen uptake (kg ha<sup>-1</sup>):** The nitrogen content in plant was determined by Kjeldahl's method. The grain and straw were separated and then grinded. The powdered plant sample of 0.5 g was digested in concentrated sulphuric acid using copper sulphate and potassium sulphate mixture as catalyst in digestion unit for 3 hrs and temperature was maintained at 410°C. The digested material was then distilled with 40 percent sodium hydroxide and distilled was collected in boric acid containing the mixture indicator. The content was estimated by titrating the distilled against N/20 sulphuric acid. The nitrogen uptake was calculated by multiplying the dry weight with nitrogen content. In order to get total uptake of nitrogen, the uptake values for grain and straw were added together.

**Phosphorous uptake (kg ha<sup>-1</sup>):** Total phosphorus uptake was determined in the extract by Vandomolybe yellow colour method. The optical density was measured with photoelectric

colorimeter at 470 nm. The content was estimated with calibration curve. The phosphorous uptake by grain and straw per hectare was calculated with the help of percent value of phosphorus and yield of grain and straw. In order to get uptake of phosphorus, the uptake value for grain and straw was added together plot wise.

**Digestion of plant sample:** 0.5 g powdered plant samples were weighed and digested with Diacid (HNO<sub>3</sub>: HClO<sub>4</sub>) mixture at 9:4 ratios in hot plate till clear solution was observed or till white fumes cease to come out. Then the contents were cooled and transferred to 50 ml volumetric flask and volume was made up to the mark by adding distilled water. Then the contents were filtered through Whatman No. 1 filter paper and a known quantity of aliquot were used for further analysis of phosphorus and potassium.

**Potassium uptake (kg ha<sup>-1</sup>):** The potassium uptake was determined with the help of Flame photometer [11] and was estimated with calibration curve. Total uptake of potassium by grain and straw was calculated by multiplying their relative contents with yield and values were added to know the total uptake of potassium in kg ha<sup>-1</sup>.

The uptake of N, P and K at harvest in seed and straw was estimated by using following formula:

$$\text{Nutrient uptake (kg/ha)} = \frac{\text{Nutrient content (\% in seed} \times \text{seed yield (kg/ha)} + \text{Nutrient content (\% in straw} \times \text{straw yield (kg/ha)}}{100}$$

**Analysis of available nutrients in soil:** Available nitrogen (kg ha<sup>-1</sup>): An alkaline permanganate solution was used to determine nitrogen availability [12].

**Available phosphorus (kg ha<sup>-1</sup>):** As referred to by [13], Olsen's extract (0.5 N sodium bicarbonates solution of pH (8.5) was used to estimate available P.

**Available potassium (kg ha<sup>-1</sup>):** In order to determine potassium availability, N-neutral ammonium acetate was used, as described by [11].

**Statistical analysis:** According to [14], an analysis of variance will be performed using a Randomized Block Design and the results will be analyzed at a level of significance of 5%.

### 3. RESULTS AND DISCUSSION

A summary of the results of the present investigation is presented in Tables 1, 2 and 3.

**Growth Parameters:** The data on plant height at 60 DAS was differed significantly among the treatments. The significantly higher plant height was obtained in T<sub>8</sub> (31.33) which was found at par T<sub>7</sub> (30.55). Similarly, plant dry weight also differed significantly and the maximum plant dry matter associated with T<sub>8</sub> (10.40) which was significantly maximum over other treatments but found at par with T<sub>7</sub>. The data on CGR at 40-60 DAS revealed that significantly higher crop growth rate was found at T<sub>8</sub> (0.0019) which was found at par with T<sub>7</sub> and T<sub>6</sub>. This might be due to higher availability and uptake of N, P and other nutrients by the crop and as these inoculations plays important role in improving availability of N and P. Seed inoculation with bio-inoculants also produce fungistatic and growth promoting substances which influenced the plant growth. Similar results were found by [15], [16] and [6].

**Nutrient Uptake study (kg ha<sup>-1</sup>):** Seed inoculation with *Bacillus megaterium* and *Bacillus mucilaginosus* as well as spray *Bacillus megaterium* and *Bacillus mucilaginosus* significantly improved the N, P and K uptake by seed and straw.

**Nitrogen uptake by seed and straw:** It is clear from the data that nitrogen uptake by seed as well as straw differed significantly due to various treatments of RDF and microbial strains. Application of T<sub>8</sub> treatment recorded higher nitrogen uptake by grain (74.57) and straw (18.80) which was statistically at par with T<sub>7</sub> and these treatments were found significantly superior over rest of the treatments. This increase may be related to the supply of P necessary for Rhizobium to fix relatively more nitrogen from soil, resulting in increased plant growth and N uptake by roots and shoots. In plants, phosphorus plays an important role in physiological and developmental processes, and its beneficial effect might have accelerated the growth process that increases N uptake. [17] also found similar results.

**Phosphorus uptake by seed and straw:** The data pertaining to P uptake by seed and straw as affected significantly by different treatments. Application of 100% RDF with *Bacillus*

*megaterium* and *Bacillus mucilaginosus* as seed treatment + foliar application of *Bacillus megaterium* and *Bacillus mucilaginosus* at 30 and 45 DAS recorded significantly higher P uptake by seed (4.11) which was found at par with T<sub>7</sub>, T<sub>6</sub>, T<sub>5</sub>, T<sub>4</sub>, T<sub>3</sub> and T<sub>2</sub>. Similarly, significantly higher P uptake by straw (4.38) was recorded at T<sub>8</sub>. The principal mechanism for mineral phosphate solubilization is the production of organic acids and acid phosphatases play a major role in the mineralization of organic phosphorous in soil. Similar results were concluded by [18] and [19].

**Potassium uptake by seed and straw:** The analysis results on potassium uptake by seed and straw showed significant variation due to various RDF and microbial strains treatments. Among the various treatments, the significantly highest potassium uptake in seed was recorded in T<sub>8</sub> (21.04) which was found at par with T<sub>7</sub>, T<sub>6</sub>, T<sub>5</sub>, T<sub>4</sub> and T<sub>3</sub>. However, straw uptake was recorded significantly higher T<sub>8</sub> (21.15) which was found at par with T<sub>7</sub>. This might be due to the reason that potash mobilizing bacterial strains (*Bacillus mucilaginosus*) produces plant growth promoting substances which benefits the plant growth and development by mobilizing potash and making it available to crops. It also enhances the efficiency of chemical fertilizer as well. Similar findings were reported by [20].

**Total uptake study (kg/ha):** The data on total NPK uptake by crop revealed that NPK uptake differed significantly among different treatments. Among all the treatments, Application of T<sub>8</sub> recorded significantly higher N uptake (93.37) which was found at par with T<sub>7</sub>. Similarly, P uptake was also recorded significantly higher in T<sub>8</sub> (8.49) but was found at par with all treatments except control with 75% and 100% RDF. The K uptake was significantly higher in T<sub>8</sub> (42.19) which was found at par T<sub>7</sub>, T<sub>6</sub> and T<sub>5</sub>.

**Nutrient available in soil after harvesting of crop:** The data revealed that available N did not vary significantly among the treatments. However, the maximum value of available N was found in T<sub>8</sub> (210.0 kg ha<sup>-1</sup>). The available P showed significant variation among the treatments and the significantly higher value was obtained in T<sub>8</sub> (15.9 kg ha<sup>-1</sup>) which was found at par with T<sub>7</sub> and T<sub>5</sub>. Available K also varied significantly and the significantly higher value was recorded in T<sub>8</sub> (480.5 kg ha<sup>-1</sup>) which was found at par with T<sub>7</sub>.

**Table 1. Impact of *Bacillus megaterium* and *Bacillus mucilaginosus* strains on growth parameters of soybean**

Treatment		Plant height (cm)	Dry matter accumulated (g plant <sup>-1</sup> )	CGR (g m <sup>-2</sup> day <sup>-1</sup> )
		60 DAS	60 DAS	40-60 DAS
T <sub>1</sub>	Control with 75% RDF	30.33	8.58	0.0014
T <sub>2</sub>	Control with 100% RDF	29.66	8.90	0.0015
T <sub>3</sub>	75% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment	30.44	9.02	0.0015
T <sub>4</sub>	100% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment	30.66	9.53	0.0017
T <sub>5</sub>	75% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as foliar application at 30 and 45 DAS	30.44	8.68	0.0014
T <sub>6</sub>	100% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as foliar application at 30 and 45 DAS	29.89	9.58	0.0017
T <sub>7</sub>	75% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment + foliar application of <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> at 30 and 45 DAS	30.55	10.23	0.0017
T <sub>8</sub>	100% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment + foliar application of <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> at 30 and 45 DAS	31.33	10.40	0.0019
<b>SEm (±)</b>		0.33	0.10	0.00004
<b>CD at 5%</b>		<b>0.99</b>	<b>0.32</b>	<b>0.00013</b>

**Table 2. Impact of *Bacillus megaterium* and *Bacillus mucilaginosus* strains nutrient uptake of soybean**

Treatment	Nutrient uptake (kg ha <sup>-1</sup> )						
	Grain			Straw			
	N	P	K	N	P	K	
T <sub>1</sub>	Control with 75% RDF	43.31	2.22	12.90	11.92	2.52	13.30
T <sub>2</sub>	Control with 100% RDF	48.42	2.71	14.04	14.56	3.28	16.23
T <sub>3</sub>	75% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment	45.41	2.93	15.61	15.16	3.48	16.74
T <sub>4</sub>	100% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment	54.33	3.00	16.40	15.88	3.42	17.01
T <sub>5</sub>	75% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as foliar application at 30 and 45 DAS	57.27	3.02	16.35	15.46	3.57	17.62
T <sub>6</sub>	100% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as foliar application at 30 and 45 DAS	61.06	3.37	17.78	16.90	3.60	18.80
T <sub>7</sub>	75% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment + foliar application of <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> at 30 and 45 DAS	69.20	4.01	19.48	17.82	3.54	20.50
T <sub>8</sub>	100% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment + foliar application of <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> at 30 and 45 DAS	74.57	4.11	21.04	18.80	4.38	21.15
<b>SEm (±)</b>		3.82	0.58	2.22	0.60	0.15	0.58
<b>CD at 5%</b>		<b>11.58</b>	<b>1.74</b>	<b>6.74</b>	<b>1.83</b>	<b>0.44</b>	<b>1.77</b>

**Table 3. Impact of *Bacillus megaterium* and *Bacillus mucilaginosus* strains on total uptake of soybean**

Treatment		Total uptake crop (kg ha <sup>-1</sup> )		
		N	P	K
T <sub>1</sub>	Control with 75% RDF	55.23	4.74	26.20
T <sub>2</sub>	Control with 100% RDF	62.98	5.99	30.27
T <sub>3</sub>	75% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment	60.57	6.41	32.35
T <sub>4</sub>	100% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment	70.21	6.42	33.41
T <sub>5</sub>	75% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as foliar application at 30 and 45 DAS	72.73	6.59	33.97
T <sub>6</sub>	100% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as foliar application at 30 and 45 DAS	77.96	6.97	36.58
T <sub>7</sub>	75% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment + foliar application of <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> at 30 and 45 DAS	87.02	7.55	39.98
T <sub>8</sub>	100% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment + foliar application of <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> at 30 and 45 DAS	93.37	8.49	42.19
<b>SEm (±)</b>		4.42	0.73	2.80
<b>CD at 5%</b>		<b>13.41</b>	<b>2.18</b>	<b>8.51</b>

**Table 4. Impact of *Bacillus megaterium* and *Bacillus mucilaginosus* strains on nutrient availability after harvest of soybean**

Treatment		Nutrient after crop harvest (kg ha <sup>-1</sup> )		
		N	P	K
T <sub>1</sub>	Control with 75% RDF	202.5	12.1	416.6
T <sub>2</sub>	Control with 100% RDF	205.6	12.2	421.6
T <sub>3</sub>	75% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment	204.6	12.3	430.0
T <sub>4</sub>	100% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment	205.0	12.3	441.6
T <sub>5</sub>	75% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as foliar application at 30 and 45 DAS	205.2	12.4	453.3
T <sub>6</sub>	100% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as foliar application at 30 and 45 DAS	205.6	12.3	465.0
T <sub>7</sub>	75% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment + foliar application of <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> at 30 and 45 DAS	205.8	12.6	476.6
T <sub>8</sub>	100% RDF with <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> as seed treatment + foliar application of <i>Bacillus megaterium</i> and <i>Bacillus mucilaginosus</i> at 30 and 45 DAS	206.0	12.7	485.0
<b>SEm (±)</b>		2.9	0.1	5.3
<b>CD at 5%</b>		<b>NS</b>	<b>0.4</b>	<b>16.2</b>

#### 4. CONCLUSION

According to the research analysis, both organic and biofertilizers applied to soybean plants result in significant variations in nutrient uptake features. It was determined that the most effective approach to increase crop yield and associated characteristics was to apply 100% RDF with *Bacillus megaterium* and *Bacillus mucilaginosus* strains as seed treatment and to apply *Bacillus megaterium* and *Bacillus mucilaginosus* strains at 30 and 45 DAS foliarly. In conclusion, the T<sub>8</sub> module offers a feasible and economical way to increase output and enhance quality over the long term.

#### ACKNOWLEDGEMENT

The Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior, Madhya Pradesh, provided the funding and facilities that were required for the authors to perform this study, and they are grateful for their assistance.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. FAO. Plant protection and production. Paper. 1982;4:1–208.
2. Imran A, Muhammad A, Shah Z, Bari A. Integration of peach (*Prunus persica* L.) residues, beneficial microbes and phosphorous enhance phenology, growth and yield of soybean. Russian Agricultural Sciences. 2020;46:223–30.
3. Muhammad A, Inamullah HK, M Arif. Germination and field emergence potential of soybean land races vs improved varieties under different sowing dates. Pure and Applied Biology. 2017; 6 (1):146–52. DOI: 10.19045/bspab.2017.60007
4. Anonymous. Agricultural statistics at a glance 2019. Government of India, Ministry of Agriculture and Farmers Welfare, Directorate of Economics and Statistics. 2020:74.
5. Dahal BR, Bhandari S. Biofertilizer: A next generation fertilizer for sustainable rice production. International Journal of Graduate Research and Review. 2018; 5(1):1–5.
6. Aakash, Singh MK, Saikia N, Bhayal L, Bhayal D. Effect of integrated nutrient management on growth, yield attributes and yield of green pea in Humid Subtropical Climate of Indo-Gangetic Plains. Ann. Agric. Sci. 2023;44(2):190-196. Available: <https://epubs.icar.org.in/index.php/AAR/article/view/141927>
7. Yasin M, Mussarat, W, Ahmad K, Ali A, Hassan, SA. Role of biofertilizers in flax for ecofriendly agriculture. Science International (Lahore). 2012;24(1):95–99.
8. Souza R, Ambrosini A, Passaglia, LMP. Plant growth-promoting bacteria as inoculants in agricultural soils. Genet. Mol. Biol. 2015;38:401–419.
9. Omotayo OP, Babalola OO. Resident rhizosphere microbiome's ecological dynamics and conservation: Towards achieving the envisioned Sustainable Development Goals, a review. Int. Soil Water Conserv. Res. 2021;9:127–142.
10. Imran A, Khan A, Inam I, Ahmad F. Yield and yield attributes of Mungbean (*Vigna radiata* L.) cultivars as affected by phosphorous levels under different tillage systems. Cogent Food & Agriculture. 2016;2:1151982.
11. Jackson ML. Soil Chemical Analysis. Prentice Hall of India Pvt.Ltd., New Delhi; 1973.
12. Subbiah BV, Asija GC. A rapid procedure for the estimation of available nitrogen in soils. Current Science, 1956;25(7):259-260.
13. Olsen SR, Cole VC, Watanabe FS and Dean LA. Estimation of available phosphorus in soil by extraction with sodium bicarbonate (NaHCO<sub>3</sub>). U.S.D.A. Circular No. 939, Washington; 1954.
14. Panse VG, Sukhatme PV. Statistical Method for Agricultural Workers, ICAR, New Delhi; 1985.
15. Sarawa H, Arma MJ. Effect of biological fertilizer on the growth and nodules formation to soya bean (*Glycine max* (L.) merrill) in ultisol under net house conditions. Journal of Experimental Biology and Agricultural Sciences. 2016;4(6).
16. Umale SM, Thosar VR, Anita BC, Chimote AN. Growth response of soybean to P solubilising bacteria and Phosphorous levels. J. Soils and Crops. 2002;12 (2):258-261.
17. Basir A, Khanzada S, Zahir S. Effect of phosphorus and farm yard manure on

- nitrogen nutrition and grain yield of chickpea. Sarhad J. Agric. 2005;21(1):11-19.
18. Htwe AZ, Moh SM, Moe K, Yamakawa T. Effects of co-inoculation of *Bradyrhizobium japonicum* SAY3-7 and *Streptomyces griseoflavus* P4 on plant growth, nodulation, nitrogen fixation, nutrient uptake, and yield of soybean in a field condition. Soil Sci. Plant Nutr. 2018; 64:222–229.
  19. Ezawa T, Smith SE, Smith FA. P metabolism and transport in AM fungi. Plant Soil. 2002; 244:221-230.
  20. Kumar B, Ismail S, Patil VD. Role of microbial solubilisers on major nutrient uptake - A review. J Pharmacogn Phytochem. 2017;6(5):641-644.

© 2023 Solanki et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/108019>