



# Assessment of Nutrient Status in the Cassava Growing Tracts of Southern Laterites (Agro Ecological Unit 8), Kerala, India

R. A. Aparna <sup>a\*</sup>, G. Rajasree <sup>a</sup>, P. Shalini Pillai <sup>a</sup>,  
Sheeba Rebecca Isaac <sup>a</sup>, Roy Stephen <sup>a</sup>, B. Aparna <sup>a</sup>  
and Pratheesh P. Gopinath <sup>a</sup>

<sup>a</sup> Department of Agronomy, College of Agriculture, Vellayani, Thiruvananthapuram, Kerala, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJECC/2023/v13i113509

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/109283>

**Original Research Article**

**Received: 05/09/2023**

**Accepted: 11/11/2023**

**Published: 18/11/2023**

## ABSTRACT

As part of the study, a survey was conducted to identify the soil fertility constraints of cassava growing tracts of AEU 8, Thiruvananthapuram, which is one of the major cassava growing regions of the State of Kerala in peninsular India. The method of sampling of the survey was random sampling wherein a random sample of size one was taken from each of 50 ha cassava area. Panchayat wise soil samples were collected (77 samples) by geo referenced soil sampling method from major cassava growing areas of the unit. The samples are analyzed for the soil fertility parameters following standard analytical procedures. Results indicated that the soil was very strongly acidic to strongly acidic, normal electrical conductivity, medium organic carbon, medium nitrogen and potassium, high phosphorus, sufficient calcium, sulphur, zinc and copper, deficient magnesium and boron.

\*Corresponding author: E-mail: [aparnara6@gmail.com](mailto:aparnara6@gmail.com);

**Keywords:** Agro ecological unit 8; cassava; soil sampling; available nutrients.

## 1. INTRODUCTION

An agro ecological unit (AEU) is characterized by distinct ecological responses to the macro environmental elements, which are reflected in the vegetation, soils and agricultural land use. The State of Kerala in Indian subcontinent has been divided into 23 distinct AEU's [1]. The AEU 8 (Southern Laterites) extends to the south western part of Thiruvananthapuram district in Kerala covering Thiruvananthapuram city and Nemom, Athiyanthoor, Parassala, Perumkadavila and Vellanadu blocks comprising 24 panchayats and one municipality covering a geographical area of 387.27 square kilometers. Climate is moist sub humid tropical monsoon type. Soils are deep, red, and strongly acid and loam. Land use is dominantly coconut plantations and narrow valleys are cultivated with rice, vegetables, tapioca and banana.

Cassava (*Manihot esculenta* Crantz) is one of the world's most important staple food crops. The crop holds a predominant position in cropping systems of Southern Kerala and occupies an area of 3690 ha in AEU 8 [2]. In the present scenario, rice cultivation in lowlands is being replaced by short duration cassava varieties and increasing cassava production for food diversification would be a way to help achieve food security of the population.

Among the short duration varieties of cassava, Vellayani Hraswa is the most popular variety in Southern Kerala. The nutrient management of Vellayani Hraswa is at present based on the blanket recommendation for varieties with different duration and growth habits. However balanced fertilizer application is possible through assessment of nutrient demand of the crop and innate soil fertility status. Hence, development of scientific nutrient management practices will help the farmers of the AEU 8 to judiciously use the nutrients and to achieve better productivity.

## 2. MATERIALS AND METHODS

### 2.1 Soil Sample Collection and Analysis

The method of sampling followed in the survey was random sampling wherein a sample of size one was taken randomly from each of 50 ha cassava area. Panchayat

wise (23 panchayat) soil samples were collected by geo referenced soil sampling method from major cassava growing areas of the unit.

For soil sample collection, each field was divided into uniform areas and each area was sampled separately. After removing the stubbles and litter, a 'V' shaped cut was made to a depth of 15 cm by using spade. Samples collected from the uniform areas were mixed together to get a composite sample and after the quartering process, 500 g of soil sample was obtained. A total of 77 geo-referenced composite soil samples were drawn at random and the collected soil samples were dried in shade and were analyzed for the soil fertility parameters such as soil reaction, soluble salts, organic carbon, available nitrogen, phosphorus, potassium, calcium, magnesium, sulphur, boron, copper and zinc following standard analytical procedures (Table 1).

## 3. RESULTS AND DISCUSSION

### 3.1 Soil Fertility Status of AEU 8

The soil test data such as pH, EC, organic carbon and available N, P, K, Ca, Mg, S, Fe, Mn, Cu and Zn covering the agro ecological unit 8 was processed and interpreted to arrive at the overall soil fertility status of AEU 8. Distribution of point data over fertility classes of soil reaction, organic carbon and nutrients in cassava growing tracts of AEU 8 were tabulated and given in Table 2.

### 3.2 Soil Reaction

Out of 77 soil samples, 25 samples are very strongly acidic (32.47 percent) and 23 samples are strongly acidic (29.87 percent) with value ranging from 4.78 to 7.20 with a mean value of  $5.55 \pm 0.65$ . According to Nair et al. [3] low content of basic cations and high Al saturation of exchange complex which are responsible for acidity are primarily due to lack of liming. Another reason for the increased acidity is the high rainfall prevalent in the area. The area receives an average annual rainfall of 1884 mm [1]. The problem of soil acidity in the tract was found to have aggravated by the unscientific use of acid producing fertilizers, lack of lime application etc. as identified in the survey.

### 3.3 Electrical Conductivity

Out of 77 soil samples 76 samples are normal (98.70 per cent) with value ranging from 0.042 to 2.60 dS m<sup>-1</sup> with a mean value of 0.16± 0.28 dS m<sup>-1</sup>. Electrical conductivity (EC) is a measure of the soluble salts and is a measure of the cations and anions present in a soil which in turn is a measure of the salinity of the soil.

### 3.4 Organic Carbon

The available organic carbon status of the cassava growing tract of AEU 8 showed that 52 soil samples (67.53 per cent) tested medium range organic carbon with a mean content of (1.45± 0.46) % and a range of 0.51 to 3.18%. Most soils of AEU 8 being the laterite type low activity clay soils, it is essential to maintain high organic matter content in the soil. Cassava is a heavy feeder and requires nutrients in abundant quantity. In spite of the higher requirement of nutrients, the cassava growing tracts of AEU 8 had a medium level of soil organic carbon content in general. The crop cassava is characterized by leaf abscission leading to leaf fall and fallen leaves add a moderate quantity of dry matter to the soil along with major and minor nutrients.

Lack of application of sufficient quantity of organic manures and non-adoption of management practices to protect the soil organic carbon could have seriously affected the organic carbon status of soil. Decline in the organic carbon status of cassava cultivated soils under continuous cultivation was previously reported by Francis et al. [4].

### 3.5 Primary Nutrients

The available nitrogen status of the AEU 8 showed that 42 soil samples (54.55 per cent) tested had medium range nitrogen with value ranging from 125.44 to 489.22 kg ha<sup>-1</sup> with a mean value of 289.81 ± 69.19 kg ha<sup>-1</sup>. Most of the farmers in AEU 8 use poultry manure as organic source in the cassava field and is usually applied in several split doses upto two months before the harvest. Poultry manure has higher nitrogen content (1.12-1.28 per cent) as reported by Limisha [5]. However the late split application of this manure would be resulted in the release of N to the soil after the crop duration which would have reflected on the available N status of the soil.

In general, majority of the study area could be rated as high in available phosphorus, with value ranging from 1.12 to 897.90 kg ha<sup>-1</sup> with a mean value of (232.81±212.58) kg ha<sup>-1</sup>. Out of the total, 63 soil samples (81.82 per cent) tested high phosphorus content. The High P levels in these soils are usually due to over-fertilizing through high analysis complex or straight fertilizers having P. The major portion of Kerala soils has high P availability due to excess use of phosphatic fertilizers through factomphos which is manufactured in the State itself [6]. As explained by Ahmed et al. [7], when exogenous P content is lower, there are abundant adsorption sites on soil colloids and there is a higher chemical adsorption binding capacity for soil P, and lower desorption capacity.

The available potassium status of the AEU 8 showed that 41 soil samples (53.25 per cent) tested had medium range of potassium with values ranging from 79.20 to 986.73 kg ha<sup>-1</sup> with a mean value of (265.90 ± 200.55) kg ha<sup>-1</sup>. The cation exchange capacity of the low activity clay minerals in the soils does not permit retention of significant amounts of potassium in exchangeable form. The lateritic nature of the soil and prevailing humid climate necessitates regular application of potassium fertilizers to the crop plants. Cassava takes up a substantial quantity of potassium from the soil and the harvested roots in particular contain large amounts of K - the NPK ratio in the roots being 5:1:10 in comparison to the typical ratio of 7:1:7 as in other crops [8]. Uptake of K<sup>+</sup> is a driving process for accessing both the exchangeable pool and even a significant part of the nonexchangeable pool in the soil [9].

### 3.6 Secondary Nutrients

In the case of available calcium status, 50 samples (64.94 per cent) of the samples had sufficient Ca content which could be due to the practice of liming by majority of the cassava cultivators. The range of calcium was found to vary from 70 to 610.00 mg kg<sup>-1</sup> with an average of (330.38±133.72) mg kg<sup>-1</sup>. Another possible reason for higher Ca content in soil could be the antagonistic effect of higher soil P content on Ca through competition in the absorption of these nutrients for the same transporters such as non-specific carrier proteins [10]. This antagonistic influence could have affected the crop uptake of Ca, leaving unutilized amount of this nutrient in the soil.

Magnesium was deficient in 52 samples (67.53 per cent) with a mean content of (114.44 ± 66.90) mg kg<sup>-1</sup> and a range of 18 to 396.00 mg kg<sup>-1</sup>. Magnesium is less strongly bound to soil colloids due to its higher hydrated ionic radii and hence easily leached off. The predominance of coarse textured soils, well drained condition and intense leaching losses, and less addition of inputs supplying magnesium to soil [11] could be the reasons for prevalence of magnesium deficiency in the tract.

Analysis of the sulphur status of the soils indicted that 47 soil samples (61.04 per cent) had sufficient sulphur content with a mean value of (15.17± 20.98) mg kg<sup>-1</sup> and a range of 0.5 to 90.00 mg kg<sup>-1</sup>. This is in conformation with the general status of sulphur in Kerala soils. Use of the complex fertilizer factomphos (N:P:K:S @ 20:20:0:13) which is manufactured in Kerala and widely available throughout the State would have resulted in sufficient S status in Kerala soils as reported by Rajasekharan et al. [6].

### 3.7 Micronutrients

The range of boron was found to vary from 0.004 to 0.404 mg kg<sup>-1</sup> with an average of (0.13 ± 0.11) mg kg<sup>-1</sup>. Available boron was deficient in 100 per

cent soil samples. Boron is one of the important micronutrient among all essential micronutrients required for proper growth and development of plants. An estimated annual B requirement of 3.9 thousand tonnes by 2025 is an indication of its emergence as a major limiting nutrient to obtain optimum yield of many crops [12]. Major soils of Kerala, derived from acid igneous rocks are, however, deficient in B [13]. Moreover, being highly mobile in the soil [14], leaching losses further aggravate B insufficiency in the high rainfall zones of Kerala, frequently leading to development of deficiency symptoms in crop plants.

Available zinc was sufficient in 76 samples (98.70 per cent) with a mean content of (9.75 ± 5.20) mg kg<sup>-1</sup> and a range of 0.86 to 29.42 mg kg<sup>-1</sup>. Low Zn status of soil could have been attributed to the antagonistic effect of high P content in soils as reported by Tisdale et al. [14].

Available copper was sufficient in 77 samples (100 per cent) with a mean content of (3.49±2.06) mg kg<sup>-1</sup> and a range of 1.04 to 9.38 mg kg<sup>-1</sup>. Copper is an important ingredient of common fungicides and retention of element in soil organic matter may explain the adequate level of copper in the soil.

**Table 1. Standard analytical procedures adopted for soil samples in Agro Ecological Unit 8**

Sl.no	Parameters	Method used
1	pH	Soil and water suspension of 1:2.5 and read in pH meter [15]
2	Electrical conductivity (dS m <sup>-1</sup> )	Soil and water suspension of 1:2.5 and read in EC meter [15]
3	Organic carbon (%)	Walkley and Black method [16]
4	Available N (kg ha <sup>-1</sup> )	Alkaline permanganate method [17]
5	Available P <sub>2</sub> O <sub>5</sub> (kg ha <sup>-1</sup> )	Ascorbic acid reduced molybdophosphoric blue colour method [18]
6	Available K <sub>2</sub> O (kg ha <sup>-1</sup> )	Neutral normal ammonium acetate extract using flame photometer (Jackson,1958)
7	Available Ca (mg kg <sup>-1</sup> )	Neutral normal ammonium acetate extract using Atomic Absorption Spectrophotometer
8	Available Mg (mg kg <sup>-1</sup> )	Neutral normal ammonium acetate extract using Atomic Absorption Spectrophotometer
9	Available S (mg kg <sup>-1</sup> )	CaCl <sub>2</sub> extract-turbidimetry method [19]
10	Available B (mg kg <sup>-1</sup> )	Azomethine H method
11	Available Zn (mg kg <sup>-1</sup> )	HCl acid extract method using Atomic Absorption Spectrophotometer [20]
12	Available Cu (mg kg <sup>-1</sup> )	HCl acid extract method using Atomic Absorption Spectrophotometer [20]

**Table 2. Distribution of point data over fertility classes of soil reaction, organic carbon and nutrients in cassava growing tract of AEU**

Parameters	Fertility class	Critical range	No of samples	Per cent samples
Soil reaction pH	Extremely acidic	3.5 - 4.4	0	0.00
	Very strongly acidic	4.5 – 5.0	25	32.47
	Strongly acidic	5.1 – 5.5	23	29.87
	Moderately acidic	5.6 - 6.0	6	7.79
	Slightly acidic	6.1 – 6.5	17	22.08
	Neutral		6	7.79
Electrical conductivity	Normal	< 0.8	76	98.70
	Critical for germination	>0.8	1	1.30
	Low	< 0.5	0	0.00
OC	Medium	0.5 – 1.5	52	67.53
	High	>1.5	25	32.47
	Low	< 280	35	45.45
N	Medium	280 - 560	42	54.55
	High	> 560	0	0.00
	Low	< 11	9	11.69
P	Medium	11 - 24	5	6.49
	High	>24	63	81.82
	Low	<116	11	14.28
K	Medium	116 - 275	41	53.25
	High	> 275	25	32.47
	Deficient	<300	27	35.06
Ca	Sufficient	>300	50	64.94
	Deficient	<120	52	67.53
Mg	Sufficient	>120	25	32.47
	Deficient	<5	30	38.96
S	Sufficient	5-10	47	61.04
	Deficient	<0.5	77	100.00
B	Sufficient	>0.5	0	0.00
	Deficient	<1	1	1.30
Zn	Sufficient	>1	76	98.70
	Deficient	<1	0	0.00
Cu	Sufficient	>1	77	100.00

#### 4. CONCLUSION

According to the survey, management practices by most of the farmers for cassava in AEU 8 are poorly related to actual crop requirements and probably lead to large nutrient losses due to over-application and yield losses due to under application. The overall fertility status of agroecological unit 8(AEU 8) indicated that the soil was very strongly acidic to strongly acidic, normal electrical conductivity, medium organic carbon, medium nitrogen and potassium, high phosphorus, sufficient calcium, sulphur, zinc and copper, deficient magnesium and boron and it was used for developing a scientific nutrient management package which could help the farmers in judicious management of costly

chemical fertilizers and to achieve better productivity through increasing the nutrient use efficiencies.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. KSPB [Kerala State Planning Board]. Soil fertility assessment and information management for enhancing crop productivity in Kerala. Kerala State Planning Board, Thiruvananthapuram. 2013;514.

2. GOK [Government of Kerala]. Kerala Soil Health Information System; 2019 [online]. Available:<http://www.keralasoilfertility.net> [01 Dec. 2019].
3. Nair KM, Kumar KSA, Lalitha M, Kumar SCR, Koya A, Parvathy S, Sujatha K, Thamban C, Mathew J, Chandran KP, Haris A. Surface soil and subsoil acidity in natural and managed land-use systems in the humid tropics of Peninsular India. *Current Science*. 2019;116(7):1201–1211.
4. Francis NA, Constant AMF, Ijang TP, Atanga NS, Clarisse MMY, Mapiemfu-Lamare D, Simon NT. Effects of cassava cultivation on soil quality indicators in the humid forest zone of Cameroon. *Greener Journal of Agricultural Science*. 2013;3(6):451-457.
5. Limisha NP Organic nutrition in taro (*Colocasia esculenta* (L) Schott). Ph. D. thesis, Kerala Agricultural University, Thrissur. 2022;252.
6. Rajashekhara P, Nair KM, Susan John K, Suresh Kumar P, Narayanan Kutty MC, Ajith RN. Soil fertility related constraints to crop production in Kerala. *Indian Journal of Fertilizers*. 2014;10(11):56-62.
7. Ahmed W, Jing H, Kailou L, Ali S, Tianfu H, Geng S. Impacts of long-term inorganic and organic fertilization on phosphorus adsorption and desorption characteristics in red paddies in southern China. *PLoS ONE*. 2021;16(1):54-65.
8. Vanlauwe BP, Pypers, Sanginga N. The potential of integrated soil fertility management to improve the productivity of cassava-based systems. In: *Cassava: Meeting of the challenges of the new millennium: Proceedings of the First Scientific Meeting of the Global Cassava Partnership 21-25 July 2008, Ghent, Belgium*. Institute of Plant Biotechnology for Developing Countries (IPBO), Ghent University, Ghent, Belgium.
9. White PJ, George TS, Gregory PJ, Bengough, AG, Hallett PD, McKenzie BM. Matching roots to their environment. *Annals of Botany*. 2013;112:207-222.
10. Rietra RP, Heinen M, Dimkpa CO, Bindraban PS. Effects of nutrient antagonism and synergism on yield and fertilizer use efficiency. *Communications in Soil Science and Plant Analysis*. 2017; 48:1895–1920.
11. Bhindhu PS, Sureshkumar P. Availability indices of calcium and magnesium in soils of Kerala. *Journal of Tropical Agriculture*. 2021;59(1):38-44.
12. Murthy IYLN. Boron studies in major oilseed crops. *Indian Journal of Fertilisers*. 2006;1:11-20.
13. SSO. Benchmark Soils of Kerala. Soil survey organization, agriculture (SC Unit) Department, Govt. of Kerala, Thiruvananthapuram. 2007;623.
14. Subbiah BV, Asija CL. A rapid procedure for the estimation of available N in soils. *Current Science*. 1956;25:259-260.
15. Tisdale SL, Nelson WL, Beaton JD. *Soil fertility and fertilizers*, 4th ed. Macmillan Publishing Company, New York. 1985;754.
16. Jackson ML. *Soil Chemical Analysis*. (Indian Reprint, 1976). Prentice Hall of India, New Delhi. 1958;498.
17. Walkley A, Black IA. An examination of the degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*. 1934;37(1): 29-38.
18. Subbiah BV, Asija CL. A rapid procedure for the estimation of available N in soils. *Current Science*. 1956;25:259-260.
19. Bray RH, Kurtz LT. Determination of total, organic and available forms of phosphorus in soils. *Soil Science*. 1945; 59:39-45.
20. Chesnin L, Yien CR. Tubidimetric determination of available sulphate. *Proceedings of Soil Science Society of America*. 1951;15:149-151.
21. Sims JR, Johnson GV. Micronutrient soil tests. In: Mortvedt, J.J., Cox, F., Rhuman, L.M., and Welch, R.M. *Micronutrient in Agriculture* (2nd ed.), SSSA, Madison, USA. 1991;427-476.

© 2023 Aparna et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/109283>