



# Morphometric Studies of Tobacco Caterpillar, [*Spodoptera litura* (Fabricius)] on Different Host Plants

Rishabh Kumar <sup>a</sup>, S.K. Sharma <sup>a\*</sup>, Tarak Nath Goswami <sup>a</sup>,  
Jigyasa <sup>a</sup> and Nawazish Ahmed <sup>a</sup>

<sup>a</sup> Department of Entomology, Bihar Agricultural College, Bihar Agricultural University,  
Sabour, Bhagalpur – 813210, Bihar, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2024/v36i54587

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/115548>

Original Research Article

Received: 15/01/2024

Accepted: 25/02/2024

Published: 12/04/2024

## ABSTRACT

Tobacco caterpillar, *Spodoptera litura* (Fabricius) is an economically important polyphagous pest inflicting significant economic damage to numerous field and horticultural crops. Research was conducted at the Department of Entomology, Bihar Agricultural University, Sabour, Bhagalpur, Bihar, in laboratory settings to validate the presence of this pest. The investigation involved detailed measurements of the various larval stages, pupa, adult male, and female, of tobacco caterpillar including the wingspan. The first to sixth instars of *S. litura* exhibited significantly greater length and width as well as width of head capsule, when reared on the tomato as a host. The pupae of *S. litura* reached at their maximum length and width when raised on marigold (16.75 mm and 4.36 mm), while the minimum pupal measurements (14.10 mm and 4.42 mm) was observed in case of maize. In regards to the adult moths, both male and female specimens displayed the longest length and wing span when associated with tomato (16.71 mm and 18.22 mm for length, 38.47 mm and 38.73 mm for wingspan, respectively), whereas the smallest measurements were recorded for those

\*Corresponding author: E-mail: drsanjay71@gmail.com;

associated with maize (14.35 mm and 16.95 mm for length, 35.10 mm and 35.12 mm wingspan respectively). The present study confirmed the presence of *S. litura* that require immense attention to prepare the management strategy against it.

**Keywords:** Tobacco caterpillar; larva; morphometry; tomato; marigold.

## 1. INTRODUCTION

*Spodoptera litura* (Fabricius) is a highly polyphagous insect that significant threat to agriculture and many other countries [1]. This insect has a wide range of host plants and has been reported to damage over 112 plant species from 44 different families [2]. In India alone, it has been observed affecting about 40 plant species Chari and Patel [3]. Various crops are known to serve as host plants for *Spodoptera litura*, and researchers [4-6] have documented its presence and damage in several agricultural crops. Some of the crops known to be affected by *Spodoptera litura* viz., maize, castor, cauliflower, tobacco and several grasses etc. *Spodoptera litura*'s ability to infest such a wide range of crops and plant species makes it an important agricultural pest that requires efficient management strategies to mitigate its impact on crop yields. Other hosts include ornamental plants, native vegetation, unwanted plants, and shade trees, such as *Leucaena leucocephala*, which serves as a shade tree in Indonesian cocoa plantations [7].

The damage inflicted by *S. litura* on tomatoes is significantly more pronounced. This pest has been documented to initiate leaf skeletonization in the early stages and extensive defoliation in later stages, thereby diminishing the plants' photosynthetic capability [8]. Entomopathogenic nematodes (EPNs) are beneficial organisms that parasitize insect pests and are increasingly employed as a biopesticide against a wide array of such pests [9]. Their remarkable qualities have sparked considerable commercial interest in utilizing nematodes as biological insecticides, presenting a promising alternative to chemical approaches in integrated pest management (IPM) strategies. EPNs possess numerous attributes that render them effective biocontrol agents. They exhibit behavior akin to insecticides or conventional plant protection chemicals, and they can be seamlessly integrated into IPM programs. Cultivating and applying EPNs is straightforward, utilizing established methods and equipment. Moreover, they demonstrate compatibility with various chemical pesticides, making them a valuable addition to IPM initiatives [10].

## 2. MATERIALS AND METHODS

Mass rearing of Tobacco Caterpillar was initiated under laboratory conditions. Early instars were collected from the field and brought to the laboratory and maintained in plastic container having holes in the lids for their proper growth and development and larvae were fed with fresh chopped cabbage. The paired adult was allowed to mate in separate mating cages. After eggs laying, these were separated and kept under BOD incubator in Petri plates for further development, and to maintain larval population of *Spodoptera litura*.

To initiate the morphometrics study of *S. litura* under laboratory condition, the samples of each instar as well as other stages were preserved in 70 % alcohol. The length and width of larvae, pupae, adults (5 of each stage) and width of head capsule were measured by using stereo zoom binocular microscope having magvision image analysis software and reference scale. Moulting was confirmed by presence of exuviae.

Data on morphometrics of different stages viz. larva, pupa and adult of *S. litura* on different hosts were measured by using stereo zoom binocular microscope having magvision image analysis software and reference scale. To record the length and width, head capsule of larvae, 5 samples (larvae) from each host (maize, marigold, pigeon pea and tomato) were killed and preserved in 70 % alcohol. For further observation, the specimens were transferred to blotting paper for removing moisture then mount on glass slide to study under on stereo zoom binocular and reference scale. Several characters were linearly measured using stereo zoom binocular and on reference scale viz. Head capsule, larval body length, pupal length and width, wing span as well as body length of male and female adults were measured.

## 3. RESULTS AND DISCUSSION

### 3.1 First Instar Larvae

The data in Table 2 and 3 indicates that the maximum length and width of first instar larvae

and usual width of head capsule were observed in tomato, measuring 2.74 mm and 0.45 mm, 0.34 mm respectively. In contrast, when considering maize and marigold, their respective values for first instar larval length were 1.58 mm and 1.38 mm, while their widths were 0.34 mm and 0.26 mm, and their size of head capsule 0.32 mm and 0.31 mm respectively. However, the minimum length and width were recorded in pigeon pea, measuring 1.15 mm in length and 0.30 mm in width along with their head capsule width 0.27 mm

### 3.2 Second Instar Larvae

Table 2 and 3 clearly demonstrates that the second instar larva exhibited its maximum length and width as well as head capsule measuring 8.65 mm and 1.35 mm, 0.53 respectively, on tomato plants. These measurements were similar to those observed on maize, which had lengths of 8.24 mm and widths of 1.13 mm. The average width of head capsule on maize is 0.50. When it comes to marigold, their respective measurements for II instar larval length were 4.14 mm and 0.54 mm, with head capsule width 0.46 mm. Conversely, Pigeon pea displayed notably smaller larval measurements, with a minimum length of 3.55 mm and a minimum width of 0.44 mm with widths of head capsule 0.42 mm.

### 3.3 Third Instar Larvae

The lowest larval length and width measurements as well as head capsule width were observed in case of pigeon pea, measuring 7.75 mm and 0.74 mm, 0.64 mm respectively. On the other hand, marigold and maize displayed varying lengths and widths. Specifically, marigold had larval lengths of 10.58 mm and widths of 1.47 mm, maize had lengths of 11.42 mm and widths of 2.07 mm. Marigold and maize had average width of head capsule 0.71 mm and 0.74 mm respectively. The maximum length and width measurements were obtained from tomato, measuring 16.75 mm and 2.21 mm, respectively with head capsule width 0.87 mm. (Table 2 and 3)

### 3.4 Fourth Instar Larvae

In Table 2 and 3, it is evident that fully grown larvae displayed their maximum body length and width on tomato plants, measuring 26.84 mm and 2.94 mm, respectively. The head capsule width of 1.24 mm was measured on tomato.

These measurements were distinct from those observed in other treatments.

### 3.5 Fifth Instar Larvae

The minimum larval length and width were observed on marigold 23.15 and 3.57 mm respectively. The width of head capsule 1.60 mm was measured on marigold. On maize the respective values were 24.23 mm, 3.68 mm with head capsule 1.65 mm. The maximum larval length and width were recorded from tomato 32.65 and 3.68 mm respectively. The width of head capsule 2.44 mm was measured on tomato. (Table 2 and 3)

### 3.6 Sixth Instar Larvae

The fully grown larvae exhibited their maximum length and width on tomato plants, measuring 37.52 mm and 4.52 mm, respectively. Notably, width of head capsule measured on tomato was 3.35 mm. Conversely, when considering maize and marigold, their respective measurements for fully grown larval length were 29.10 mm and 28.00 mm, with widths of 4.25 mm and 4.19 mm. whereas, measurement of width of head capsule of maize and marigold 1.98 mm and 1.89 mm respectively. These measurements were statistically at par to each other as indicated in Table 2 and 3.

#### 3.6.1 Pupae

Table 1 clearly illustrates that the maximum length and width of fully grown pupae were observed in marigold, measuring 16.75 mm and 4.36 mm, respectively. These measurements were statistically similar to those recorded for tomato, which had a length of 16.72 mm and 4.83 mm width. However, the lowest recorded length (14.10 mm) and width (4.42 mm) were found in maize.

#### 3.6.2 Adult

##### 3.6.2.1 Female moth


The data presented in the table (Table 3) on the maximum body length of female moths and their wing spans. The results indicated that the longest body length was observed in female moths from tomato, measuring 18.22 mm, followed by marigold 17.67 mm and maize 16.04 mm in maize. The maximum wing span of female moths was on tomato (38.47 mm). followed by was 37.74 mm in marigold. These

measurements were similar to each other and slightly smaller wing span of female moths were recorded from maize, which measured 35.10 mm.

### 3.6.2.2 Male moth

Table 3 clearly indicated that maximum body length and wing span of male moth were observed on tomato (16.71 and 38.73 mm), followed by marigold (16.34 and 38.13 mm), and maize (14.35 and 35.12 mm).

The comparative impact of four different host plants viz., maize, tomato, marigold, and pigeon pea were studied on the morphological characteristics of *S. litura*. Notably, the larvae of *S. litura* exhibited the most significant increase in both length and width when reared on tomato plants, followed by maize, marigold and pigeon pea. This observation aligns with the findings of Kumar and Bhattacharya [11], who also noted similar trends in larval development when fed on tomato leaves.



**Fig. 1. Morphometric study of different instars and pupae of *Spodoptera litura***  
 A- Morphometric of early instars larvae. B- Head capsule measurement of early instars C-Morphometric of late instar larvae. D-Head capsule measurement of late instars larvae. E- Morphometric of pupae

Table 1. Morphometric variation of Tobacco caterpillar, *S. litura* larvae and pupae on different hosts

S.No.	Host plants	Tomato		Maize		Marigold		Pigeon pea		
		Instars	Length (mm)	Width (mm)	Length (mm)	Width (mm)	Length (mm)	Width (mm)	Length (mm)	Width (mm)
1	I		2.74±0.49	0.45±0.06	1.58±0.12	0.34±0.02	1.38±0.04	0.26±0.04	1.15±0.09	0.30±0.15
2	II		8.65±0.56	1.35±0.08	8.24±0.12	1.13±0.06	4.14±0.10	0.54±0.13	3.55±0.12	0.44±0.33
3	III		16.75±0.85	2.21±0.08	11.42±0.26	2.07±0.05	10.58±0.23	1.47±0.16	7.75±0.19	0.74±0.04
4	IV		26.84±0.87	2.94±0.24	19.25±0.33	2.36±0.16	18.78±0.28	2.28±0.04	-	-
5	V		32.65±0.69	3.70±0.35	24.23±0.45	3.68±0.02	23.15±0.39	3.57±0.02	-	-
6	VI		37.52±0.61	4.52±0.20	29.10±0.34	4.25±0.29	28.00±0.33	4.19±0.04	-	-
7	Pupae		16.72±0.49	4.83±0.21	14.10±0.07	4.42±0.34	16.75±0.85	4.36±0.01	-	-

Table 2. Width of head capsule and growth rate of *Spodoptera litura* during larval development

S. No.	Host plants	Tomato		Maize		Marigold		Pigeon Pea		
		Instar s	Mean ±SE	Growth Ratio	Mean± SE	Growth ratio	Mean± SE	Growth ratio	Mean± SE	Growth ratio
1	I		0.34±0.03	-	0.32±0.02	-	0.31±0.01	-	0.27±0.01	-
2	II		0.53±0.03	1.55	0.50±0.01	1.56	0.46±0.02	1.48	0.42±0.02	1.55
3	III		0.87±0.02	1.64	0.74±0.02	1.48	0.71±0.01	1.42	0.64±0.01	1.52
4	IV		1.24±0.02	1.24	1.05±0.03	1.41	1.03±0.02	1.45	-	-
5	V		2.44±0.04	1.96	1.65±0.02	1.57	1.60±0.03	1.55	-	-
6	VI		3.35±0.02	1.37	1.98±0.02	1.20	1.89±0.01	1.18	-	-

**Table 3. Morphometric variation in adult moths of tobacco caterpillar, *Spodoptera litura* on different hosts****(A) Female:**

S. No.	Host Plants	Length (mm)			Wing span (mm)		
		Min.	Max.	Mean± SE	Min.	Max.	Mean± SD
1	Tomato	16.40	19.45	18.22±0.34	34.90	41.11	38.73±0.57
2	Maize	14.57	16.95	16.04±0.20	31.20	37.32	35.12±0.40
3	Marigold	16.27	19.05	17.67±0.44	34.39	40.62	38.13±0.58
4	Pigeon pea	-	-	-	-	-	-

**(B) Male:**

S.No.	Host Plants	Length(mm)			Wing span (mm)		
		Min.	Max.	Mean± SE	Min.	Max.	Mean± SD
1	Tomato	15.10	17.97	16.71±0.42	36.10	40.95	38.47±0.80
2	Maize	12.11	15.87	14.35±0.45	32.65	36.87	35.10±0.43
3	Marigold	14.80	17.78	16.34±0.41	35.92	39.77	37.74±0.70
4	Pigeon pea	-	-	-	-	-	-

Average width of head capsule of *S. litura* were measured on four different host, maximum width was observed on tomato 3.35 mm, followed, by maize 1.98 and least development 1.89 mm was recorded on marigold. In terms of pupal development, marigold plants yielded the most substantial pupae, with a recorded length of 16.75 mm and a width of 4.36 mm, followed by tomato, maize. This result is consistent with the research of Ramzan et al. [12], who reported pupal measurements of 14.22 mm in length and 4.10 mm in width on maize. On the contrary, Ramya and Jha [13] observed different pupal dimensions of 16.72 mm in length and 4.83 mm in width when reared on tomato leaves under controlled laboratory conditions.

Regarding the adult stage, tomato proved to be the most favourable host, yielding females with a maximum body length of 18.22 mm and a wing span of 38.73 mm, and males with a body length of 16.71 mm and a wing span of 38.47 mm. Marigold also supported relatively larger adults, while maize was the least suitable host. These findings are in line with those reported by Kumari et al. [14], who found similar measurements for males (15.84 mm in length and 32.12 mm in wing span) and females (14.08 mm in length and 31.64 mm in wing span) when reared *S. litura* on maize in laboratory conditions. Based on the morphological characteristics observed at different developmental stages, it can be concluded that tomato is the most conducive host for the growth of *S. litura*, followed by marigold and maize [15].

**4. CONCLUSION**

The first to sixth instars of *S. litura* exhibited significantly greater length and width as well as

width of head capsule, when reared on the tomato as a host. The pupae of *S. litura* reached at their maximum length and width when raised on marigold (16.75 mm and 4.36 mm), while the minimum pupal measurements (14.10 mm and 4.42 mm was observed in case of maize). In regards to the adult moths, both male and female specimens displayed the longest length and wing span when associated with tomato (16.71 mm and 18.22 mm for length, 38.47 mm and 38.73 mm for wingspan, respectively), whereas the smallest measurements were recorded for those associated with maize (14.35 mm and 16.95 mm for length, 35.10 mm and 35.12 mm wingspan respectively).

**ACKNOWLEDGEMENT**

Author is thankful to The Bihar Agricultural College, Bihar Agricultural University, Sabour, Bhagalpur for providing the financial support and necessary facilities during the study period.

**COMPETING INTERESTS**

Authors have declared that no competing interests exist.

**REFERENCES**

1. Shivayogeshwara B, Mallikarjunaiah H, Krishnaprasad NK and Shetty MVN. Integrated management of *Spodoptera litura* Fabricius (Noctuidae: Lepidoptera) in FCV tobacco crop. Tobacco Research. 1991;17:59-61.
2. Yadav S, Patil J, Sharma HK. Bio-efficacy of *Steinernema carpocapsae* against

- Spodoptera litura under laboratory condition. J Pure Appl Biosci. 2017;5:165-72.
3. Chari MS and Patel NG. Cotton leaf worm Spodoptera litura, its biology and integrated control measures. Journal of Cotton Research & Development. 1983; 13:7-8.
  4. Basu AC. Effect of different food on larval and post larval development of moths of Prodenia litura Fabricius (Lepidoptera: Noctuidae). Bombay natural history society. 1944;44:275-88.
  5. Patel PH, Sisodiya DB, Raghunandan BL, Patel NB, Gohel VR, Chavada KM. Bioefficacy of Entomopathogenic fungi and bacteria against invasive pest Spodoptera frugiperda (JE Smith) under laboratory condition. Journal of Entomology and Zoology Studies. 2020;8(6):716-720.
  6. Thobbi VV. Growth potential of Prodenia litura Fabricius in relation to certain food plants in Surat, Indian Journal Entomology. 1961;22:262-64.
  7. CABI. Invasive Species Compendium: Datasheets, maps, images, abstracts and full text on invasive species of the world. Available: <http://www.cabi.org/isc/datasheet/44520>. Accessed 18 November 2019.
  8. Chauhan D, Srivastava RP. Bioefficacy of three insecticides against tobacco caterpillar, Spodoptera litura (Fab.). Journal of Entomological Research. 2015; 39(3):243-7.
  9. Askary TH, Abd-Elgawad MM. Opportunities and challenges of entomopathogenic nematodes as biocontrol agents in their tripartite interactions. Egyptian Journal of Biological Pest Control. 2021;31:1-0.
  10. Abd-Elgawad MM. Nematode spatial distribution in the service of biological pest control. Egyptian Journal of Biological Pest Control. 2024;34(1):3.
  11. Kumar HD, Bhattacharya S. Biology of Spodoptera litura (Fabricius) on different crop plants. Journal of Entomological Research. 2019;43(2):165-8.
  12. Ramzan M, Sajid Z, Sattar Z, Abbas D, Yaseen T, Mehmood S, Yaseen I. Biological and Morphological Parameters of Armyworm, Spodoptera litura in Cabbage and Maize Plants under Laboratory Conditions in Southern Punjab, Pakistan. Journal of Environmental Issues and Agriculture in Developing Countries. 2020;12(2&3).
  13. Ramya SN, Jha S. Biology and morphometry of Spodoptera litura (Fabricius) on tomato. Journal of Entomology and Zoology Studies. 2018;6 (5):80-82
  14. Kumari K, Kumari K, Saha T. Morphometric studies of fall armyworm, Spodoptera frugiperda (JE Smith) on maize. Journal of Experimental Zoology India. 2022;25(2).
  15. Kumar R, Sharma SK, Kumari JK, Anand A. Comparative Efficacy of Different Biopesticides against the Tobacco Caterpillar [Spodoptera litura (Fabricius)]. Journal of Experimental Agriculture International. 2024;46(5):507-12.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/115548>