



Mycoplasma Pneumonia as a Rare Culprit in Acute Myocarditis

**H.El Ghiati ^{a*}, R.Djafarou ^a, K.Tahani ^b, D.Britel ^a, I.Asfalou ^a,
Z.Lakhal ^a, K. Ennibi ^b and A.Benyass ^a**

^a Cardiology Center, Mohammed V Military Training Hospital, and Mohammed V University, Morocco.

^b Department of Virology, Mohammed V Military Training Hospital, and Mohammed V University, Morocco.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124389>

Case Report

Received: 30/07/2024

Accepted: 27/09/2024

Published: 01/10/2024

ABSTRACT

Mycoplasma pneumoniae (*M. pneumoniae*) is an atypical bacterium transmitted by respiratory droplets through close contact. It is a common cause of community-acquired pneumonia, with an overall mortality rate of approximately 30%. Besides respiratory symptoms, *M. pneumoniae* can cause extrapulmonary complications including myocarditis in 1%-5% of cases leading to arrhythmias and heart failure. We are introducing the case of a 16-year-old boy who presented with acute chest pain, due to myocarditis related to *Mycoplasma pneumoniae*, an unsuspected cause of myocarditis. This case underscores the importance of considering atypical pathogens in myocarditis cases and emphasizes the need for comprehensive diagnostic approaches to guide timely and appropriate therapeutic interventions for rare presentations of cardiac involvement due to *Mycoplasma pneumoniae*.

*Corresponding author: E-mail: elghiati.hanaa@gmail.com;

Cite as: Ghiati, H.El, R.Djafarou, K.Tahani, D.Britel, I.Asfalou, Z.Lakhal, K. Ennibi, and A.Benyass. 2024. "Mycoplasma Pneumonia As a Rare Culprit in Acute Myocarditis". *Asian Journal of Case Reports in Medicine and Health* 7 (1):151-55. <https://journalajcrah.com/index.php/AJCRMH/article/view/203>.

Keywords: Myocarditis; mycoplasma pneumoniae; cardiac MRI; antibiotherapy.

1. INTRODUCTION

The *Mycoplasma pneumoniae* (*M. pneumoniae*) is a pleomorphic organism and a common cause of community-acquired pneumonia, accounting for 10%–40% of cases, with overall mortality around 30%. The population most at risk is children and young people [1, 2]. Additionally, to respiratory involvement, patients are susceptible to developing multiple extra-pulmonary pathologies, including dermatological, gastrointestinal, hematological, neurological, and cardiovascular involvement, such as myocarditis, pericarditis, and other cardiac manifestations [3]. Myocarditis is an unusual complication that occurs in only 1-5% of cases [4]. Herein, we report a case of a 16-year-old boy who presented with chest pain for a few days following respiratory and gastrointestinal symptoms due to *Mycoplasma pneumoniae*-related myocarditis and pneumonia.

2. CASE REPORT

In this abstract, we explore the uncommon occurrence of acute myocarditis attributed to *Mycoplasma pneumoniae*, an unusual culprit bacterium. The study delves into a distinctive case involving a 16-year-old boy who presented to the emergency department with acute myocarditis.

The patient presented to the emergency department with acute chest pain. He had no particular history. He reported experiencing cough, diarrhea, and vomiting four days prior to presentation.

On admission, cardiovascular and pulmonary examinations were normal. Blood pressure was normal at 130/60 mmHg, with a heart rate of 61 beat/minute and pulse saturation of 98% on room air. In laboratory findings, the leukocyte count was normal, while C-reactive protein was slightly elevated at 26.3 mg/L (normal range: <6 mg/L). The electrocardiogram (ECG) showed negative T waves in the inferior leads (Fig. 1).

The ELISA test for *Mycoplasma pneumoniae* IgM was positive, while the nasopharyngeal multiplex PCR panel including adenovirus, coronavirus, rhinovirus, enterovirus, influenza A/B, respiratory syncytial virus, *Chlamydia*, and *Bordetella* were all negative. Hepatitis B and C serology is nonreactive. Two days before admission, the patient's general practitioner prescribed a macrolide, but there was no improvement.

A thoracic scan revealed right pneumonia (Fig. 2). The transthoracic echocardiogram was normal and showed no pericardial effusion (Fig. 3), but troponin levels were elevated at 1061 ng/L (normal range for men: 18-39 ng/L). Cardiac MRI (Fig. 4) indicated signs of myocarditis limited to the basal segment of the inferior (A) and inferolateral (B) walls. A 24-hour ECG recorder did not detect any arrhythmias.

The patient was started with fluoroquinolones therapy (Levofloxacin 500mg bid) combined with adjuvant treatment consisting of vitamin C and paracetamol. He remained under observation at ICU for 48 hours before discharge after good clinicobiological progress. A follow-up at 3 months showed a stable condition.




Fig. 1. ECG showing negative T waves in inferior leads



Fig. 2. Thoracic CT Scan showing right basal pneumonia




Fig. 3. Echocardiogram revealed normal contractility of the heart and no pericardial effusion




Fig. 4. Cardiac MRI showing the presence of patchy late subepicardial gadolinium enhancement in the inferior wall (A) and the inferolateral wall (B) indicative of cardiac edema

3. DISCUSSION

Acute myocarditis is a rare but potentially serious affection involving inflammation of the cardiac muscle. Myocarditis is usually associated with viral infections but can also be caused by

bacterial agents such as *Mycoplasma pneumoniae*. *Mycoplasma pneumoniae* is a major cause of atypical pneumonia, but its association with myocarditis is less frequent and still not fully understood.

The diagnosis of *M. pneumoniae* infection is difficult to determine due to the diversity of manifestations. An estimated 50% of patients present with mild, varied, and non-specific upper respiratory tract symptoms, accompanied by discomfort and low-grade fever. The condition mainly affects children and young adults, most of whom develop mild, benign upper respiratory tract infections. Patients may develop pneumonia in 3-10% of cases and extrapulmonary involvement in 5-10%, which is often clinically more severe than the primary infection [5, 6]. In our case, the patient was presented with cough, gastrointestinal symptoms, without fever, and then chest pain suggesting cardiac involvement.

Cardiac involvement is rare but often severe and occurs more frequently in adults, usually in the form of pericarditis, endocarditis, or myocarditis. The physiopathology of cardiac damage in myocarditis is thought to result from immune modulations such as autoimmunity, allergy, or immune complex formation [7].

Myocarditis is an underdiagnosed cardiac condition, that can be caused by various processes, including immunological, infectious, and toxic. Outcomes may include recovery, sudden deterioration, heart failure, the development of dilated cardiomyopathy, and even death [8]. The main causes of myocarditis are infectious, and viruses such as coxsackievirus, CMV, adenovirus, parvovirus B19, hepatitis C, HIV, varicella, and EBV are the most common [9]. Although *M. pneumoniae* is a rare cause of myocarditis, it should be considered in the differential diagnosis because of the necessity of rapid diagnosis and treatment to reduce morbidity and disabling sequelae [10]. In the study reported by Paz et al., cardiac involvement in *M. pneumoniae* infections was more frequent in patients with severe respiratory involvement; 19% of patients with carditis also had pleural effusion and 43% had pneumonia. The clinical presentation was varied, characterized by chest pain in 38% of cases, as in our patient, and palpitations in 43%. In this study, electrical abnormalities were present in all patients [11].

Further investigations, including cardiac magnetic resonance imaging (MRI) and endomyocardial biopsy (EMB), confirmed the diagnosis of acute myocarditis, with evidence of myocardial edema at MRI and inflammatory cell infiltration, injury to the cardiomyocytes resulting

in fibrosis, necrosis, muscle fiber hypertrophy, or atrophy at EMB [12]. The cardiac MRI that was performed in our patient confirmed the diagnosis of myocarditis in inferior and inferolateral walls. These anomalies may explain the negative T waves observed in inferior leads on ECG.

It is true that the responsibility of *Mycoplasma Pneumoniae* in the occurrence of myocarditis in this case has been difficult to prove, but given the temporal relationship between the respiratory and gastrointestinal symptomatology and the cardiac presentation, as well as the biological, electrical and MRI data, it remains the only possible etiology.

Management of acute myocarditis due to *M. pneumoniae* depends on effective antibiotic therapy. Several studies have highlighted the efficiency of macrolides in the treatment of *M. pneumoniae* infections [1, 13]. However, cases of macrolide resistance have also been reported [14], as in our patient, whose clinical condition worsened on macrolides (Azithromycin). Furthermore, fluoroquinolones can also be used in the treatment of *M. pneumoniae* infections, with comparable efficiency to macrolides [1], as evidenced by the clinical improvement in our patient a few days after the onset of fluoroquinolone therapy.

4. CONCLUSION

This case highlights the importance of considering *Mycoplasma pneumoniae* as a potential trigger for acute myocarditis in patients presenting with cardiac symptoms in the setting of a recent respiratory infection. Clinicians should maintain a high index of suspicion for myocarditis in such cases and promptly initiate appropriate diagnostic and treatment strategies. To improve patient outcomes, further research is required to determine the pathophysiological mechanisms underlying mycoplasma-associated myocarditis and optimize therapeutic approaches.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

CONSENT

As per international standards, parental written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bajantri B, Venkatram S, Diaz-Fuentes G. *Mycoplasma pneumoniae*: a potentially severe infection. *J Clin Med Res.* 2018; 10(7):535–44.
2. Qu J, Yang C, Bao F, Chen S, Gu L, Cao B. Epidemiological characterization of respiratory tract infections caused by *Mycoplasma pneumoniae* during epidemic and post-epidemic periods in North China, from 2011 to 2016. *BMC Infect Dis.* 2018; 18(1):335.
3. Narita M: Classification of extrapulmonary manifestations due to *Mycoplasma pneumoniae* infection on the basis of possible pathogenesis. *Front Microbiol.* 2016;7:23.
4. Gouder M Formosa, M Bailey, C Barbara, C Muscat, and V Grech. *Mycoplasma pneumoniae* – an unusual cause of acute myocarditis in childhood. *Images Paediatr Cardiol.* 2006;8(4): 7–10.
5. Abdulhadi B, Kiel J. *Mycoplasma Pneumonia*. StatPearls Treasure Island (FL): StatPearls Publishing ; 2022. Accessed on : 2022 Jan 24.
6. Waites KB, Talkington DF. *Mycoplasma pneumoniae* and its role as a human pathogen. *Clinical Microbiology Reviews.* 2004;17(4):697–728
7. Fan Q, Meng J, Li P, Liu Z, Sun Y, Yan P. Pathogenesis and association of *Mycoplasma pneumoniae* infection with cardiac and hepatic damage. *Microbiology and Immunology.* 2015;59(7):375–80.
8. Rroku A, Kottwitz J, Heidecker B. Update on myocarditis—what we know so far and where we may be heading. *European Heart Journal Acute Cardiovascular Care.* 2021;10(4):455–67
9. Imazio M, Trincheri R. Myopericarditis: Etiology, management, and prognosis. *International Journal of Cardiology.* 2008; 127(1):17–26
10. Park IH, Oh YK, Kim JD, Yu ST. A case of acute myopericarditis associated with *Mycoplasma pneumoniae* infection in a child. *Korean Circulation Journal.* 2012;42 (10):709–13
11. Paz A, Potasman I. *Mycoplasma*-associated carditis: case reports and review. *Cardiology.* 2002;97(2):83–8
12. Vidusa L, Kalejs O, Maca-Kaleja A, Strumfa I. Role of Endomyocardial Biopsy in Diagnostics of Myocarditis. *Diagnostics (Basel).* 2022;12(9):2104.
13. Blauwet LA, Cooper LT. Antimicrobial agents for myocarditis: Target the pathway, not the pathogen *Heart.* 2010;96: 494-495
14. T Okada, M Morozumi, T Tajima, M Hasegawa, H Sakata, S Ohnari. Rapid Effectiveness of Minocycline or Doxycycline Against Macrolide Resistant *Mycoplasma pneumoniae* Infection in a 2011 Outbreak Among Japanese Children, *Clinical Infectious Diseases.* 2012; 55(12):1642–1649.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/124389>