



Yield and Economics of Dual Purpose Barley as Influenced by Various Nitrogen Dose and Seed Rate

**Amit Sharma^{a++*}, A. K. Dhaka^{a#}, Bhagat Singh^{a†}, Kamal^{a++},
Simran Jasht^{a++} and Mohit Godara^{b++}**

^a Department of Agronomy, College of Agriculture, Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana, India.

^b Department of Agricultural Meteorology, College of Agriculture, Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i82162

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/101135>

Original Research Article

Received: 01/04/2023

Accepted: 03/06/2023

Published: 21/06/2023

ABSTRACT

A field experiment was carried out during *Rabi* season 2019-20 at wheat and barley section research area of CCS Haryana Agricultural University, Hisar. The experiment was comprised of three nitrogen doses (N₁-60, N₂-75 and N₃- 90 kg N/ha) as main plot treatments and four seed rates (S₁- 87.5, S₂- 100, S₃- 112.5 and S₄-125.0 kg/ha) as sub plot treatments. With four replications, experiment was laid out in split plot design. Based on the research investigation, it was found that nitrogen dose and seed rate both had significantly influenced the yield attributes and yield of dual purpose barley. Among nitrogen doses, 90 kg/ha being at par with 75 kg/ha recorded significantly higher grain yield (4895 kg/ha), biological yield (12220 kg/ha) and green fodder yield (3706 kg/ha) which were 9.7, 12.0 and 15.1 percent higher than nitrogen dose of 60 kg/ha, respectively due to

⁺⁺ PhD Research Scholar;

[#] Assistant Professor;

[†] Assistant Scientist;

*Corresponding author: E-mail: amitsharma110297@gmail.com, amitsharma1197@hau.ac.in;

higher yield attributes i.e. number of effective tillers (85.08), number of grains per spike (44.15), number of spikelets per spike (23.10), spike length (7.47) and test weight (37.82) with nitrogen dose 90 kg/ha which were 8.5, 5.1, 7.9, 11.4 and 4.8 percent higher than nitrogen dose of 60 kg/ha, respectively. Among seed rates, 125.0 kg/ha closely followed by 112.5 kg/ha recorded significantly higher effective tillers (85.11), grain yield (4922 kg/ha), biological yield (12181 kg/ha), green fodder yield (3729 kg/ha) which were 8.0, 11.0, 11.8 and 16.7 per cent higher than seed rate of 87.5 kg/ha, respectively.

Keywords: Dual purpose barley; seed rate; nitrogen dose; green fodder yield; grain yield.

1. INTRODUCTION

“Barley (*Hordeum vulgare* L.) is the one of the first domesticated cereal of world agriculture. It stands in fourth place among cereal crops after wheat, rice and maize in the world with a share of 7.0 per cent of the global cereal production. Barley is considered as poor man’s crop that may thrive well in problematic soils and marginal land. It is not only useful for malting, feed and food purpose but also its β – glucanase is useful in lowering the risk of cardio-vascular diseases” [1]. “It is a valuable crop because it is used for food, processed food and feed for livestock. Its straw is used for making hay and silage. Besides these conventional uses, it is an important industrial crop as it is used as raw material for beer, whiskey, baby foods, coca malt drinks, *ayurvedic* medicines and brewing industries” [1].

The global production of barley is amounted to 156.41 million metric tons. In India barley was grown on 0.59 mha area with production of 1.722 mt yielding 29.20 q of grains productivity per ha [2]. The major barley producing states in India are Rajasthan, U.P., M.P. and Haryana. In Haryana, area under barley crop was 12120 ha with production 46500 tons and productivity was 3837 kg/ha during 2019-20 [2].

“Barley possesses very high regeneration capacity compared to other cereals after cutting up to attainment of jointing stage. Lodging close to harvest is usually a major problem in cereals under good nutrition and assured water supply conditions, if winds blow after irrigation. Therefore, it is reasonable to assume that one cutting for green forage at active growth stage will reduce the lodging chances in barley. It will also help in mitigating the fodder shortage. By taking one cutting during the active vegetative growth stage and then leaving the regenerated crop for grain production, barley’s regeneration capacity may be put to use” [3]. “Barley being a fast-growing crop with high biomass in early stages can be utilized as green fodder with very

limited water supply or less rainfall in these areas. In drier parts of northern plains (Rajasthan, Madhya Pradesh, Southern Haryana, South-West Punjab and Western U. P.) during *Rabi*, farmers can grow dual purpose barley over other forage crops because of its dual utilization and less water requirement” [4]. “This suggests an ample scope for growing dual purpose barley for obtaining moderate yield of green fodder as well as grain from the same crop. Growing of dual purpose barley genotypes having a wider adaptability and responsive to inputs has opened a new avenue for exploiting higher grain fodder and subsequent grain yield potential” [5]. Therefore, barley can provide important nutrition to the livestock through its green fodder and grains harvested from regenerated crop.

“Proper fertilization is considered to be one of the most important pre-requisite in this respect. Amongst nutrients, nitrogen plays an important role in synthesis of chlorophyll, amino acids and other organic compounds of physiological significance in plant system” [6]. “As the nitrogen play a vital role in plant nutrient, its application has resulted in significant increase in yield and yield components of all the crops. Increased nitrogen levels improved the number of effective tillers, spike length, number of grains per spike, test weight, and grain production per hectare of barley considerably” [7]. Hence synchronizing N application to barley as per its requirement can be an important technology to obtain reasonable fodder and grain yield for dual purpose barley.

“Optimum seed rate is also considered to be one of the most important prerequisite for realizing higher green fodder as well as grain yields” [8]. “It is suggested that a higher seed rate helps in increasing the crop stand, green forage biomass and stabilize grain yield of cereals in dual purpose systems. Additional applied seed rate produced the maximum forage dry matter and biological yield” [9]. In some countries like Australia and USA, barley is commonly grown for

the dual purpose of producing forage and grain from the same crop, but in India, this kind of experience is still unexploited. So, there is a crucial need to determine the influence of nitrogen dose and seed rate for a balanced green fodder yield and grain production in dual purpose barley.

2. MATERIALS AND METHODS

A field experiment was carried out during *Rabi* season 2019-20 at wheat and barley section research area of CCS Haryana Agricultural University, Hisar which is situated in the sub-tropical region at 29° 10' N latitude and 75° 46' E longitude with an elevation of 215.2 m above mean sea level in Haryana State of India. It lies on the outer margins of the south-west (SW) monsoon region. It has tropical monsoonal climate and is characterized as arid type of climate. The major characteristics of climate in Hisar district are its extremes temperature, dryness and scanty rainfall. The average annual rainfall is around 452 mm. The soil exhibits mixed pattern of Aeolian and Alluvial deposits. During crop growing season 133.5 mm rainfall was received. The mean weekly maximum and minimum temperature ranged between 11.9 to 38 °C and 2.6 to 20 °C, respectively. The mean weekly values of morning and evening relative humidity ranged between 70 to 100 and 20 to 82 per cent respectively, while sunshine ranged between 1.1 to 8.7 hrs during crop season. The experiment was comprised of three nitrogen doses (N₁-60, N₂-75 and N₃- 90 kg N/ha) as main plot treatments and four seed rates (S₁- 87.5, S₂- 100, S₃- 112.5 and S₄-125.0 kg/ha) as sub plot treatments. Experiment was laid out in split plot design with four replications. The recommended crop management practices were applied uniformly for all experimental plots concerning irrigation schemes and weed management. The crop was harvested for green fodder at 55 days after sowing thereafter the crop was left for grain and straw. Net returns and BC ratios were calculated on basis of prevailing market prices of input and produce. Effective tillers per meter row length which have the fully developed spike were counted per meter row length from each plot at maturity. At maturity, the grains were separated from spikelets and the total numbers of grains in each spike were counted. Number of spikelets and spike length were counted from 5 randomly selected spikes and averaged to determine the number of spikelets per spike and length of spike. Green fodder yield (kg/ha) was recorded by cutting the plots raised for dual purpose at 55

days after sowing and weighted. The green forage yield obtained in the plot area and recorded in kg/ha. Grain yield (kg/ha) obtained for individual net plot after threshed, cleaned and grain obtained were recorded and converted into kg/ha. Harvest index was calculated by dividing the grain yield (economic yield) by the total dry matter (biological yield) and multiplied by hundred.

$$\text{Harvest index} = \frac{\text{Economical yield}}{\text{Biological yield}} \times 100$$

Benefit cost ratio was worked out by following formula:

$$B:C = \frac{\text{Gross returns}}{\text{Cost of cultivation}}$$

The experimental data were analyzed by using OPSTAT software available on CCS Haryana Agricultural University home page [10].

3. RESULTS AND DISCUSSION

3.1 Yield Attributes

The data on various yield attributes *viz.* effective tillers, spike length, number of grains per spike, number of spikelets per spike, test weight and as influenced by various seed rate and nitrogen levels are presented in (Table 1).

3.1.1 Effective tiller/mrl

The data pertaining Table 1 showed that number of effective tillers were significantly affected by nitrogen dose and seed rate. Increasing nitrogen dose from 60 to 90 kg/ha progressively increased number of effective tillers but non-significant differences were recorded between 75-90 kg/ha. Nitrogen dose of 90 kg/ha recorded 1.4 and 8.5 percent higher effective tillers over 75 and 60 kg/ha nitrogen dose, respectively. "It might be due to cumulative influence of growth and yield attributing characters owing to fertilization. Greater availability of metabolites (Photosynthates) and nutrients to developing reproductive structures seems to have resulted in increase in all the yield-attributing characters that ultimately improved the yield of the crop" [11].

Number of effective tillers (mrl) progressively increased with the increase in seed rate from 87.5 kg/ha to 125.0 kg/ha. Number of effective tillers obtained with 125.0 kg/ha were found significantly higher over other seed rates except seed rate of 112.5 kg/ha. Number of effective

tillers (85.08) recorded with seed rate of 125.0 kg/ha were significantly higher with a relative advantage of 8.0 and 3.6 per cent over seed rate of 87.5 and 100.0 kg/ha, respectively.

3.1.2 Number of grains per spike

A thorough examination of the data in Table 1 revealed that number of grains per spike was significantly affected by nitrogen dose. Increasing nitrogen dose from 60 to 90 kg/ha progressively increased number of grains per spike but non-significant differences were recorded between 75-90 kg/ha. Nitrogen dose of 90 kg/ha recorded 1.2 and 5.1 percent higher grains per spike over 75 and 60 kg/ha nitrogen dose, respectively. Increased yield attributes due to increasing level of N were due to better growth parameters with the increasing level of N. These findings substantiate the results of O'Donovan et al. [12], Singh et al. [13]. While seed rate failed to produce significant variation in relation to number of grains per spike. Increasing seed rates from 87.5 to 125.0 kg/ha progressively decreased number of grains per spike.

3.1.3 No. of spikelets per spike

A close perusal of data in Table 1 showed that number of spikelets per spike was significantly affected by nitrogen doses. Increasing nitrogen dose from 60 to 90 kg/ha progressively increased number of spikelets per spike but non-significant differences were recorded between 75-90 kg/ha. Nitrogen dose of 90 kg/ha recorded 1.2 and 7.9 per cent higher over 75 and 60 kg/ha nitrogen dose, respectively. The higher performance of yield was associated with higher nitrogen fertilization Oral et al. [14]. While seed rate failed to produce significant variation in relation to number of spikelets per spike. Increasing seed rates from 87.5 to 125.0 kg/ha progressively decreased number of spikelet per spike.

3.1.4 Spike length

A delve to data given in Table 1 presented that length of spike was significantly affected by nitrogen dose. Increasing nitrogen dose from 60 to 90 kg/ha progressively increased spike length but non-significant differences were recorded between 60-75 kg/ha and 75-90 kg/ha. Nitrogen dose of 90 kg/ha recorded 6.6 and 11.5 per cent higher spike length over 75 and 60 kg/ha nitrogen dose, respectively while seed rate failed to produce significant variation in relation to spike length. Increasing seed rates from 87.5 to 125.0

kg/ha progressively decreased spike length. This might be due to length of spike is more of genetic in nature.

3.1.5 Test weight (g)

A delve to data given in Table 1 presented that test weight was significantly affected by nitrogen dose. Increasing nitrogen dose from 60 to 90 kg/ha progressively increased test weight but non-significant differences were recorded between 60-75 kg/ha and 75-90 kg/ha. Nitrogen dose of 90 kg/ha recorded 1.7 and 4.8 per cent higher over 75 and 60 kg/ha nitrogen dose, respectively.

Test weight progressively decreased with the increase in seed rate from 87.5 kg/ha to 125.0 kg/ha. Test weight obtained with 87.5 kg/ha was found significantly higher over seed rate of 125.0 kg/ha. Test weight (37.71) recorded with seed rate of 87.5 kg/ha was significantly higher with a relative advantage of 4.4 per cent over seed rate of 125.0 kg/ha.

3.2 Yield Studies

A delve to data exhibited in Table 2 showing yield studies *i.e.* biological yield, grain yield, green fodder yield, straw yield and harvest index at various nitrogen doses and seed rates in dual purpose barley.

3.2.1 Biological yield (kg/ha)

The perusal of data in Table 2 revealed that progressive significant increase in biological yield with increase in nitrogen dose from 60 to 90 kg/ha was recorded. Biological yield with nitrogen dose of 90 kg/ha was found significantly higher over other nitrogen doses. Nitrogen dose of 90 kg/ha recorded 3.8 and 12.0 per cent higher over 75 and 60 kg/ha nitrogen dose, respectively. "Biological yield is a function of grain and straw yield representing reproductive and vegetative growth of the crop. The profound influence of nitrogen dose on both of these characters mediated *via* increased photosynthetic efficiency and nutrient accumulation might have ultimately led to production of higher biological yield under its application" [15]. The results of present investigation indicated higher production of dual purpose barley under influence of higher fertility levels.

The biological yield progressively increased with the increase in seed rate from 87.5 kg/ha to

125.0 kg/ha. Among the seed rates, biological yield obtained with 125.0 kg/ha was found significantly higher over other seed rates except 112.5 kg/ha. Biological yield (12181 kg/ha) recorded with seed rate of 125.0 kg/ha was significantly higher with a relative advantage of 11.8 and 5.3 per cent over seed rate of 87.5 and 100.0 kg/ha, respectively.

3.2.2 Grain yield (kg/ha)

A disquisition to data given in Table 2 and Fig. 2 exhibited that grain yield was significantly affected by nitrogen doses. Increasing nitrogen dose from 60 to 90 kg/ha progressively increased grain yield but non-significant differences were recorded between 75- 90 kg/ha. Nitrogen dose of 90 kg/ha recorded 1.2 and 7.9 percent higher grain yield over 75 and 60 kg/ha nitrogen dose, respectively. "It has been well emphasized that N fertilization play vital role in improving three major aspects of yield determination *i.e.*, formation of vegetative structure for nutrient absorption, photosynthesis and strong sink strength through development of reproductive structure and production of assimilates to fill economically enhanced sink (source strength). Thus, cumulative influence of N application seems to have maintained balanced source sink through improving both the events of crop development (vegetative and generative), ultimately resulted in increased grain yield per hectare" [16,17].

Grain yield progressively increased with the increase in seed rate from 87.5 kg/ha to 125.0

kg/ha (Fig. 3). Grain yield obtained with 125.0 kg/ha was found significantly higher over other seed rates except seed rate of 112.5 kg/ha. Grain yield (4895 kg/ha) recorded with seed rate of 125.0 kg/ha was significantly higher with a relative advantage of 11.0 and 4.8 per cent over seed rate of 87.5 and 100.0 kg/ha, respectively. Higher grain yield was recorded with seed rate 125.0 kg/ha this was due to higher number of effective tillers recorded with seed rate of 125.0 kg/ha which were significantly higher with a relative advantage of 8.0 and 3.6 per cent over seed rate of 87.5 and 100.0 kg/ha, respectively. Results are close conformity with findings of Singh et al. [18].

3.2.3 Green fodder yield (kg/ha)

A disquisition to data given in Table 2 exhibited that green fodder yield was significantly affected by nitrogen doses. Increasing nitrogen dose from 60 to 90 kg/ha progressively increased green fodder yield but non-significant differences were recorded between 75-90 kg/ha. Nitrogen dose of 90 kg/ha recorded 3.3 and 15.1 per cent higher over 75 and 60 kg/ha nitrogen dose, respectively. The increase in green fodder yields were due to taller plants having more effective tillers due to increased nitrogen uptake. This might be due to N plays important role for providing nutrient to the plants during vegetative stage. These improvement in availability of nutrients also increased the supply of metabolites that's synchronized to demand for overall improvement and development of vegetative structure.

Table 1. Effect of different nitrogen dose and seed rate on yield attributes of dual purpose barley

Treatments		Yield attributes			
Nitrogen dose	No. of effective tiller/mrl	No. of grain per spike	No. of spikelet per spike	Spike length (cm)	Test Weight (g)
Nitrogen dose					
N ₁ - 60 kg/ha	78.42	42.01	21.41	6.70	36.09
N ₂ - 75 kg/ha	83.92	43.62	22.77	7.14	37.19
N ₃ - 90 kg/ha	85.08	44.15	23.10	7.47	37.82
SEm ±	0.57	0.44	0.43	0.18	0.40
CD at 5%	1.71	1.33	1.29	0.55	1.20
Seed rate					
S ₁ -87.5 kg/ha	78.78	43.89	22.85	7.24	37.71
S ₂ -100.0 kg/ha	82.11	43.67	22.45	7.12	37.45
S ₃ -112.5 kg/ha	84.27	42.90	22.32	7.08	36.70
S ₄ -125.0 kg/ha	85.11	42.59	22.01	7.00	36.13
SEm ±	0.67	0.51	0.40	0.28	0.29
CD at 5%	2.38	NS	NS	NS	1.02

Table 2. Effect of different nitrogen dose and seed rate on yield of dual purpose barley

Treatments	Yield (kg/ha)				Harvest index
	Biological yield	Grain yield	Green fodder yield	Straw yield	
Nitrogen dose					
N ₁ - 60 kg/ha	10907	4463	3221	6444	40.96
N ₂ - 75 kg/ha	11773	4788	3586	6985	40.70
N ₃ - 90 kg/ha	12220	4895	3706	7325	40.05
SEm ±	108	69	54	99	0.54
CD at 5%	328	210	164	299	NS
Seed rate					
S ₁ -87.5 kg/ha	10892	4435	3197	6457	40.79
S ₂ -100.0 kg/ha	11571	4696	3482	6875	40.59
S ₃ -112.5 kg/ha	11890	4810	3610	7085	40.48
S ₄ -125.0 kg/ha	12181	4922	3729	7259	40.43
SEm ±	122	57	67	97	0.38
CD at 5%	429	201	238	335	NS




Fig. 1. Mean weekly weather condition during cropping period




Fig. 2. Effect of different nitrogen dose on grain yield of dual purpose barley

Green fodder yield progressively increased with the increase in seed rate from 87.5 kg/ha to 125.0 kg/ha. Green fodder yield obtained with 125.0 kg/ha was found significantly higher over other seed rates except seed rate of 112.5 kg/ha. Green fodder yield (3729 kg/ha) recorded with seed rate of 125.0 kg/ha was significantly higher with a relative advantage of 16.6 and 7.1 per

cent over seed rate of 87.5 and 100.0 kg/ha, respectively.

3.2.4 Straw yield (kg/ha)

The perusal of data in Table 2 revealed that progressive significant increase in straw yield with increase in nitrogen dose from 60 to 90

kg/ha. Straw yield with nitrogen dose of 90 kg/ha was found significantly higher over other nitrogen doses. Nitrogen dose of 90 kg/ha recorded 4.9 and 13.7 per cent higher over 75 and 60 kg/ha nitrogen dose, respectively. The significant increase in straw yield with additional N fertilization seems to be due to its direct effect in improving biomass per plant at successive growth stages as well as in plant part at harvest of the crop, while the indirect effect might be on account of increase in various morphological parameters viz., plant height and number of tillers.

Straw yield progressively increased with the increase in seed rate from 87.5 kg/ha to 125.0 kg/ha. Straw yield obtained with 125.0 kg/ha was found significantly higher over other seed rates except seed rate of 112.5 kg/ha. Straw yield (7259 kg) recorded with seed rate of 125.0 kg/ha was significantly higher with a relative advantage of 12.42 and 5.6 per cent over seed rate of 87.5 and 100.0 kg/ha, respectively.

3.2.5 Harvest index

Data mentioned in Table 2 reflects that nitrogen dose and seed rate failed to produce significant

variation in relation to harvest index. Increasing nitrogen dose and seed rate from 60 to 90 kg/ha and 87.5 to 125.0 kg/ha, respectively showed the progressive decreasing trend of harvest index. However, numerical variation was recorded among different nitrogen dose and seed rate for harvest index. Higher harvest index (40.96) and (40.79) was recorded with nitrogen dose 60 kg/ha and seed rate 87.5 kg/ha, respectively, while minimum harvest index (40.05) and (40.43) recorded with nitrogen dose 90 kg/ha and seed rate 125.0 kg/ha, respectively. Nitrogen dose and seed rate of 60 kg/ha and 87.5 kg/ha recorded 0.5 and 2.3 and 0.5, 0.8 and 0.9 per cent higher over 75 and 90 kg/ha nitrogen dose and 100, 112.5 and 125.0 kg/ha seed rates, respectively.

3.3 Economics

The data pertaining to cost of cultivation, gross returns, net returns and B:C is given as under various seed rate and nitrogen dose are presented in Table 3.

Data given in Table 3 revealed that increasing nitrogen dose and seed rates from 60 to 90 kg/ha and 87.5 to 125.0 kg/ha, respectively increased cost of cultivation, gross returns, net returns and




Fig. 3. Effect of different seed rate on grain yield of dual purpose barley

Table 3. Effect of nitrogen dose and seed rate on economics (₹ ha⁻¹) on dual purpose barley

Treatment	Economics (₹ ha ⁻¹)			
	Cost of cultivation	Gross return	Net return	B:C ratio
Nitrogen dose				
N ₁ - 60 kg/ha	62968	88787	25819	1.41
N ₂ - 75 kg/ha	63408	97602	34194	1.54
N ₃ - 90 kg/ha	63848	100437	36589	1.57
Seed rate				
S ₁ -87.5 kg/ha	62315	90166	27851	1.45
S ₂ -100.0 kg/ha	63043	95456	32412	1.51
S ₃ -112.5 kg/ha	63773	97374	33601	1.53
S ₄ -125.0 kg/ha	64503	99439	34937	1.54

B: C ratio progressively. Among different nitrogen doses, maximum cost of cultivation (₹63,848), gross returns (₹1,00,437), net returns (₹36,589) and B:C ratio (1.57) recorded with nitrogen dose 90 kg/ha, which was 1.4, 13.1, 41.7 and 11.3 per cent, respectively higher over 60 kg/ha. These results are in conformity with those of Singh et al. (2013) who reported that the higher net profit was obtained with higher dose of nitrogen in dual purpose barley over lower doses.

Among different seed rates, maximum cost of cultivation (₹64,503), gross returns (₹99,439), net returns (₹34,937) and B:C ratio (1.54) recorded with seed rate 125.0 kg/ha which was 3.5, 10.3, 25.4 and 6.2 per cent higher over 87.5 kg/ha, respectively. These findings are confirmed by Stougaard and Xue, [19].

4. CONCLUSION

Based on field research experiment, it is concluded that increasing nitrogen dose from 60 to 90 kg/ha progressively increased plant population, phenology, yield attributes and yield of dual purpose barley. Among nitrogen doses, 90 kg/ha being at par with 75 kg/ha recorded significantly higher grain yield (4895 kg/ha), biological yield (12220 kg/ha), green fodder yield (3706 kg/ha) which were 9.7, 12.0, 15.1 percent higher than nitrogen dose of 60 kg/ha, respectively due to higher yield attributes i.e. number of effective tillers (85.08), number of grains per spike (44.15), number of spikelets per spike (23.10), spike length (7.47) and test weight (37.82) with nitrogen dose 90 kg/ha which were 8.5, 5.1, 7.9, 11.4 and 4.8 percent higher than nitrogen dose of 60 kg/ha, respectively. Increase in number of days taken to booting, heading and maturity was recorded with increase in nitrogen doses from 60 to 90 kg/ha. Among seed rates, 125.0 kg/ha closely followed by 112.5 kg/ha recorded significantly higher effective tillers (85.11), grain yield (4922 kg/ha), biological yield (12181 kg/ha), green fodder yield (3729 kg/ha) which were 8.0, 11.0, 11.8 and 16.7 per cent higher than seed rate of 87.5 kg/ha, respectively. So, to obtain higher yield and economic returns for dual purpose barley (green fodder cut at 55 DAS and left after that for grain production), variety BH 946 should be sown using optimum seed rate of 125.0 kg/ha and nitrogen dose of 90 kg/ha.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kharub AS, Verma RPS, Kumar D, Kumar V, Selvakumar R, Sharma I. Dual purpose barley (*Hordeum vulgare* L.) in India: Performance and potential. J. Wheat Res. 2013;5(1).
2. Anonymous; 2020. Available: <https://www.indiastat.com>
3. Mishra BN, Kumar S. Barley: Textbook of field crop production. In: Rajendra Prasad (Tech. ed.) Directorate of Information and Publication of Agricultural, Indian Council of Agricultural Research, New Delhi. 2002:228.
4. Verma RPS, Kharub AS, Sharma RK, Singh R, Mishra B. Jau Anusandhan-Paramparik se Vayasayik Upyog ki Aur. Directorate of Wheat Research, Karnal. Res. Bull. 2007; 23:36.
5. DWR. Nitrogen scheduling for higher grain and fodder yield in dual purpose barley in north western plains zone and north eastern plains zone. Progress Report, All India Coordinated Wheat Improvement Project, Directorate of Wheat Research, Karnal, Haryana. 2010;6:4.28-4.31.
6. Havlin JL, Beaton JD, Tisdale SL, Nelson WL. Soil fertility and fertilizer, IV Edition, Pearson Education Pvt. Ltd. Indian Branch, FIE. 2003;482.
7. Narolia GP. Response of malt barley (*Hordeum vulgare* L.) to levels and scheduling of nitrogen application under varying growing environments. Ph.D. (Agron.) Thesis, Department of Agronomy, SKRAU, Bikaner; 2009.
8. Thomason WE, Brooks WS, Griffey CA, Vaughn ME. Hulled barley seeding rate effect on grain yield and yield components. Crop Sci. 2009;49:342-46.
9. Khalil SK, Khan F, Rehman A, Muhammad F, Amanullah, Khan AZ, Wahab S, Akhtar S, Zubair M, Khalil IH, Shah MK, Khan H. Dual purpose wheat for forage and grain yield in response to cutting, seed rate and nitrogen. Pak. J. Bot. 2011;43(2):937-947.
10. Sheoran OP, Tonk DS, Kaushik LS, Hasija RC, Pannu RS. Statistical software package for agricultural research workers. Recent advances in information theory, Statistics & Computer Applications by Hooda DS and Hasija RC, Department of Mathematics Statistics, CCS HAU, Hisar 1998:139-143.
11. Singh P, Singh KN, Singh R, Aga FA, Bahar F, Raja W. Evaluation of wheat

- (*Triticum aestivum* L.) genotypes for productivity and economics under graded levels of nitrogen in temperate Kashmir. Indian J. Agric. Sci. 2010;80:380-384.
12. O'Donovan JT, Turkington TK, Edney MJ, Clayton GW, McKenzie RH, Juskiw PE, Lafond GP, Grant CA, Brandt S, Harker KN, Johnson EN, May WE. Seeding rate, nitrogen rate and cultivar effects on malting barley production. Agronomy Journal. 2011;103:709-716.
 13. Singh D, Singh DR, Nepalia V, Amina Kumari. Agro-economic performance of dual purpose barley (*Hordeum vulgare* L.) varieties under varying seed rate and fertility levels. Ann. Agric. Sci. 2013;34(3): 325-229.
 14. Oral E, Kendal E, Dogan Y. Influence of nitrogen fertilization levels on grain yield and its components in barley (*Hordeum vulgare* L.). Biotechnical Faculty of the University of Montenegro; 2018.
 15. Sharma RK, Verma RPS. Effect of irrigation, nitrogen and varieties on the productivity and grain malting quality in barley. Cereal Res. Commun. 2010;38: 419-28.
 16. Shaktawat RPS, Shekhawat PS. Soil fertility status as affected with and farmyard manure in *Kharif* crops and fertilizer levels in barley (*Hordeum vulgare* L.). Indian J. Agric. Sci. 2010;80:791-4.
 17. Meena LR, Mann JS, Meena SL. Effect of levels and mode nitrogen application on dual purpose barley (*Hordeum vulgare* L.) under semi-arid condition. Indian J. Agron. 2012;57:168-170.
 18. Singh RP, Tripathi HP, Yadav AS. Effect of stages of cutting and nitrogen levels on grain and fodder yield of barley (*Hordium vulgare*). Indian J. Agric. Sci. 2009;79(1): 78-79.
 19. Stougaard RN, Xue Q. Quality versus quantity: Spring wheat seed size and seeding rate effect on wild oat interference, economic returns and economic thresholds. Weed Research. 2005;45:351-360.

© 2023 Sharma et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/101135>