



Influence of Organic Seaweed Pelleting on Seed Quality Parameters of Onion Seeds

R. Elamparithi ^{a++*} and K. Sujatha ^{a#}

^a *Department of Seed Science and Technology, Agriculture College and Research Institute, TNAU, Madurai-625104, Tamil Nadu, India.*

Authors' contributions

This work was carried out in collaboration between both authors. Author RE designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Author KS managed the analyses of the study. Author RE managed the literature searches. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i113508

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107752>

Original Research Article

Received: 04/09/2023

Accepted: 11/11/2023

Published: 18/11/2023

ABSTRACT

Aim: Small and irregular seed shapes make precision sowing difficult. Seed pelleting, which helps to make seed handling easier during sowing, is converting such seeds into bold, spherical shapes with a smooth surface. Seaweeds are a rich source of nutrients that promote growth, such as IAA, kinetin, zeatin, and gibberellins; auxins and cytokinins; metabolic enhancers; macro and micro elements; amino acids; and vitamins. Their use in crop plants has been shown to have positive effects, including early seed germination and establishment, improved crop performance and yield, increased resistance to biotic and abiotic stress, and increased seed shelf life (Bluden).

Study Design: The experiment was undertaken with three replications in a completely randomized block design. Eight treatment combinations were used in this study. Statistical analysis was carried out by AGRES software. The Critical Differences (CD) was calculated at 5 per cent probability level.

⁺⁺ *Ph.D. Scholar;*

[#] *Professor and Head;*

^{*}*Corresponding author: E-mail: elamparithi27798@gmail.com;*

Place and Duration of Study: Department of Seed Science and Technology, Agriculture College and Research Institute, TNAU, Madurai, during the period of December 2020 and November 2021.
Methodology: Organic seaweed pellet combinations can be successfully implemented to provide improved field emergence, crop establishment and production. Seeds were pelleted with different mixtures of *Sargassum* sp., *Kappaphycus* sp., *Bacillus subtilis* and Talc powder.
Results: The results exposed that between the treatments seed pelleting with combination of T₇- *Sargassum* sp + *Kappaphycus* sp + *Bacillus subtilis*+ Talc powder was found to be more efficient in comparison with other combinations.
Conclusion: Pelleting with, *Kappaphycus* sp, *Bacillus subtilis* and Talc powder was found to be better in all aspects of seed quality and biochemical parameters.

Keywords: *Bacillus subtilis*; onion; *Kappaphycus* sp.; *Sargassum* sp.; pelleting.

1. INTRODUCTION

“Onion (*Allium cepa* L.) comes below the family of Amaryllidaceae. Onion is an important vegetable crop, extensively cultivated in India and other part of Asian countries like Bangladesh, Pakistan and Philippines. The major onion producing countries are China, Turkey, Japan, Egypt, Indonesia, Iraq, Italy, Syria and Spain” [1] “India contributes 24 million tons to the global production of onion and ranks second after China with an area of 7.80 Lakh hectares” (NHB, 2020). In India, major onion producing states are Maharashtra, Madhya Pradesh, Karnataka, Rajasthan, Bihar, Gujarat, Andhra Pradesh, Haryana, West Bengal and Uttar Pradesh.

Onion seeds are small and inappropriate for mechanical sowing, seed pelleting must be used to enlarge the seeds' size. One method of treating seeds that increases the needed seed size is called seed pelleting. It involves coating the seeds with appropriate chemicals, botanicals, micronutrients, and biocontrol agents using an adhesive. For ease of handling and to aid in mechanical sowing, pelleting is mostly used on small, irregularly shaped seeds [2] and (Rajeswari et al., 2020).

“In order to facilitate precise planting, a method known as “seed pelleting” involves encasing a seed in small amounts of inert material that are just big enough to produce globular units of a specified size” [3]. The inert substance gives young seedlings a small quantity of nutrients and produces natural water-holding medium [4]. The performance of the seeds is the main emphasis of pelleting them with adhesive, fillers and bioactive compounds. This helps in reaching the desired population, which is essential for producing crops and seeds successfully. During germination and early crop growth, seed pelleting with botanicals (or) organics is the least

expensive, non-toxic method that protects against pests and diseases [5,6]. Nutrient-pelleted seeds provide better seedling emergence and early development [7]. The ability of organic pellets to absorb water is higher than that of mineral or organic-mineral pellets. Hence, the main objective of the current study was to determine which low-cost organic pelleting materials would be best for standardizing and improving seaweed pelleting combination procedures and their impact on onion seed quality.

2. MATERIALS AND METHODS

Genetically pure seeds of Onion (*Allium cepa* L.) CO (On) 5 obtained from the Department of Vegetable crops, Horticultural College and Research Institute, Tamil Nadu Agricultural University, Periyakulam was used as base material and the experiment was carried out in the Department of Seed Science and Technology, Agricultural College and Research Institute, Tamil Nadu Agricultural University, Madurai during 2021. The seeds were sprayed with adhesive (4%) after being spread out thin layer for pelleting. After transferring the wet seeds into a container, a predetermined quantity of pelleting mixture was added. Seeds were pelleted with along the following treatments viz., T₀- Control (unpelleted seeds), T₁- Talc powder, T₂- *Bacillus subtilis*, T₃- *Bacillus subtilis* + Talc powder, T₄ - *Sargassum* sp + Talc powder, T₅ - *Kappaphycus* sp + Talc powder, T₆ - *Sargassum* sp + *Kappaphycus* sp + Talc powder, T₇- *Sargassum* sp + *Kappaphycus* sp + *Bacillus subtilis* + Talc powder. The experiment was undertaken with three replications in a completely randomized block design and evaluated for germination [8], shoot length (cm), root length (cm), dry matter production 10 per seedlings (g), vigour index values [9]. After the pelleted material was removed, the remaining

parameters were assessed. The electrical conductivity was calculated in duplicate with slight modification of Presley [10] by soaking 25 seeds in 50ml water for duration of 20hand expressed as dSm⁻¹. Following the EC measurement, the seed leachate was used to measure leachate sugars (g g⁻¹) with some modifications based on Somogyi [11] and leachate free amino acids (g g⁻¹) in duplicate using a method outlined by Moore and Stein [12] with minor modifications. Also measured the activities of dehydrogenase [13], catalase [14], and antioxidants (Blois 1958).

3. RESULTS AND DISCUSSION

Between the treatments seeds pelleted with *Sargassum sp + Kappaphycussp + Bacillus subtilis + Talc powder* recorded higher germination (86%) Seedling length (17.43 cm), dry matter production (0.036g/10 seedlings) andvigour index I (1499), compared to other treatments. Unpelleted seeds recorded the lowest germination percentage (76%), Seedling length (14.86cm), dry matter production (0.022g/10 seedlings) andvigour index I (1129) (Table 1). Seaweed coating improved the parameters of physiological quality. Growth enhancing compounds such IAA, kinetin, zeatin, and gibberellins [15], auxins and cytokinins [16], metabolic enhancers, and macro and micro elements [17] are abundant in seaweeds [18]. These substances may also improve the

parameters related to seed quality. Nitrogen fixation, vegetative growth, and root development are all enhanced by biofertilizers. They release compounds that promote growth and vitamins, help in preserving soil fertility, enhance the physical characteristics of the soil, enhance soil health in general, and support the biocontrol of disease [19]. According to Suma et al. [20], seed pelleting with *Bacillus subtilis* in *Sesamum indicum* resulted in higher seedling characteristics than the control. Seeds pelleted with *Sargassum sp + Kappaphycussp+ Bacillus subtilis+ Talc powder* recorded lowest EC (0.273dSm⁻¹), leachate sugars (23.80µg g⁻¹) and leachate free amino acids (37.90µg g⁻¹) compared to other treatments (Table 2, Figs. 1,2). The highest dehydrogenase (0.072), peroxidise (1.37) and catalase activities (1.49 µmol of H₂O₂ min⁻¹ gram⁻¹) observed in T₇ treatment. The unpelleted seeds recorded higher EC (0.294dSm⁻¹), leachate sugars (28.65µg g⁻¹) and leachate free amino acids (42.25µg g⁻¹). The electrical conductivity of seed leachate was low in pelleted seed. The results were in conformity with the findings of Sujatha [21] in blackgram, redgram and cowpea and Vethanayagi *et al.*, (2009) in bhendi. Unpelleted seeds also have recorded the lowest dehydrogenase (0.060), peroxidise (1.24) and catalase activities (1.42 µmol of H₂O₂ min⁻¹ gram⁻¹). The term "antioxidant" refers to the group of vitamins, minerals, polyphenols, and carotenoids that work

Table 1. Effect of pelleting treatments on physiological quality of onion

Treatment Details	Germination (%)	Seedling length (cm)	DMP (mg/10seedlings)	Vigour index I
T ₀	76(66.67)	14.86	0.022	1129
T ₁	76(66.67)	14.97	0.023	1138
T ₂	80(63.44)	15.67	0.026	1254
T ₃	78(62.03)	15.15	0.025	1182
T ₄	83(65.65)	16.30	0.030	1353
T ₅	81(64.16)	15.88	0.029	1286
T ₆	84(66.42)	16.87	0.033	1417
T ₇	86(68.03)	17.43	0.036	1499
Mean	81(64.16)	15.89	0.028	1282
SEd	1.625	0.376	0.0008	25.004
CD (0.05)	3.444**	0.796**	0.0016**	53.007**

(Figures in parentheses indicate arc sine value)

NS - Non Significant ** - Highly Significant

SEd- Standard error of the difference between means, CD -Critical Differences

T₀- Control T₁- Talc powder T₂- *Bacillus subtilis* T₃- *Bacillus subtilis + Talc powder*

T₄- *Sargassumsp + Talc powder* T₅- *Kappaphycussp + Talc powder*

T₆- *Sargassumsp + Kappaphycus sp + Talc powder*

T₇- *Sargassum sp + Kappaphycussp + Bacillus subtilis + Talc powder*

Table 2. Effect of different pelleting treatments on biochemical parameters of onion

Treatment Details	Dehydrogenase (OD value)	Peroxidase (units/gram)	Catalase (units/gram)	Electrical conductivity (ds/m)	Leachate free amino acids (µg/g)	Leachate free sugars (µg/g)
T ₀	0.060	1.24	1.42	0.294	42.25	28.65
T ₁	0.060	1.25	1.42	0.290	42.10	28.10
T ₂	0.063	1.29	1.44	0.283	40.65	26.80
T ₃	0.061	1.27	1.43	0.287	41.22	27.50
T ₄	0.067	1.33	1.46	0.278	39.18	25.24
T ₅	0.066	1.32	1.45	0.281	39.85	26.11
T ₆	0.069	1.35	1.47	0.275	38.74	24.43
T ₇	0.072	1.37	1.49	0.273	37.90	23.80
Mean	0.065	1.30	1.45	0.283	40.24	36.33
SEd	0.001	0.028	0.033	0.004	0.746	0.418
CD (0.05)	0.002**	0.060**	0.071*	0.009**	1.582**	0.887**

NS - Non Significant ** - Highly Significant

SEd Standard error of the difference between means, CD Critical Differences

T₀– Control T₁- Talc powder T₂– Bacillus subtilis T₃- Bacillus subtilis + Talc powder

T₄- Sargassumsp + Talc powder T₅- Kappaphycussp + Talc powder

T₆- Sargassumsp + Kappaphycus sp + Talc powder

T₇- Sargassum sp + Kappaphycussp + Bacillus subtilis + Talc powder




Fig. 1. Effect of different pelleting treatments on biochemical parameters of onion

together to stop free radical damage. By eliminating free radical intermediates, antioxidants stop these chain events. They also prevent further oxidation processes by being oxidized themselves, which improves seed performance [22-26]. The enhanced antioxidant capability of seaweed-treated seeds was shown by analysis of their DPPH free radical scavenging activity. Because it breaks down hydrogen peroxide into water and oxygen and protects against the accumulation of peroxides,

peroxidase is a useful tool in the assessment of seed quality [16]. Pelleted seed had reduced levels of sugar, electrical conductivity, and free amino acids and increased levels of dehydrogenase and peroxidase. Seed pelleted with *Sargassum* sp+ *Kappaphycus* sp + *Bacillus subtilis*+ Talc powder recorded higher seed quality as well as biochemical parameters viz., dehydrogenase, peroxidase, catalase, and lowers values of EC, leachate free amino acids and leachate free sugars [27-30].




Fig. 2. Effect of different pelleting treatments on biochemical parameters of onion

4. CONCLUSION

This study revealed that seeds pelleted with different seaweed pelleting mixture *Sargassum* sp + *Kappaphycus* sp + *Bacillus subtilis* + Talc powder (T₇) were found suitable to get fine, round and smooth seed pellets in onion. The germination was not affected by pelleting. Pelleting mixture increased the seed germination, seedling growth, dry matter production and vigour index. And also increased the biochemical parameters of seed. T₇ in onion was found to be the best combinations for improving the seedling vigour. Further research and development are essential to optimize these combinations for maximum efficiency and sustainability.

ACKNOWLEDGEMENTS

Authors would like to thank University for providing Laboratory to facilitate research throughout the course of investigation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Elamparithi R, Sujatha K, Menaka C, Senthil K. Impact of organic Seaweed pelleting on seed quality and biochemical parameters in brinjal seeds. *Pharma Innov.* 2021;10:1466-9.
2. Halmer P. Enhancing seed performance for better yield and quality. *Asian Seeds Plant Mater.* 2003;10:4-7.
3. Talha Javed, Irfan Afzal, Rubab Shabbir, Kamran Ikram, Muhammad Saqlain Zaheer, Muhammad Faheem, Hafiz Haider Ali, Javaid Iqbal, Seed coating technology: An innovative and sustainable approach for improving seed quality and crop performance. *Journal of the Saudi Society of Agricultural Sciences.* 2022;21(8):536-545
4. Krishnasamy V. Seed pelleting principles and practices. *ICAR short course on seed hardening pelleting technologies for rainfed/garden land ecosystems;* 2003.
5. Tengfei X, Nanda S, Fengliang J, Qingsheng L, Xia F. Control efficiency and mechanism of spinetoram seed-pelleting against the striped flea beetle *Phyllotreta striolata*. *Scientific Reports.* 2022;12(1):9524.
6. Kavitha M, Deshpande VK, Vyakaranahal BS, Awakkanavar JS, Yashoda H, Mathad JC. Seed pelleting with organic and inorganic inputs for vigour and viability in chilli seeds. *Karnataka J. Agric. Sci.* 2009;22(2):15-19.
7. Roos EE. Storage behavior of pelleted, tableted and taped lettuce seed. *ROOS.* 1979.
8. ISTA. International Rules for Seed Testing, Rules. International Seed Testing Association. Zurich, ISTA Secretariat; 2013
9. Abdul-Baki AA, Anderson JD. Vigour determination in soybean seed by multiple criteria. *Crop Sci.* 1973;13:630-633.
10. Presley JT. Relationship of protoplast permeability of cotton seed viability and predisposition of seedling disease. *Pl. Dis. Repr.* 1958;42(7):582.
11. Somogyi, M. Notes on sugar determination. *J. Biol. Chem.* 1952;200:245-247.
12. Moore, S. and W. H. Stein. Photochromic ninhydrin method for use in chromatography of amino acids. *J. Biol. Chem.,* 1948;176:367-388.
13. Kittock DL, Law AG. Relationship of seedling vigour, respiration and tetrazolium chloride reduction by germination of wheat seeds. *Agron. J.* 1968;60:286-288.
14. Luck, H. Assay of catalase. In: *Methods in Enzymatic Analysis 2* (Ed Bermeyer), Academic Press, New York: 1974;885.
15. Zodape S, Mukhopadhyay S, Eswaran K, Reddy M, Chikara J. Enhanced yield and nutritional quality in green gram (*Phaseolus radiata* L) treated with seaweed (*Kappaphycus alvarezii*) extract. *Crop Science;* 2010.
16. Zhang, X. and E. H. Ervin. Cytokinin-containing seaweed and humic acid extracts associated with creeping bentgrass leaf cytokinins and drought resistance. *Crop Sci.* 2004;44:1737- 1745.
17. Strik WA, Novak MS, Van Staden J. Cytokinins in macroalgae. *Plant Growth Regul.* 2003;41:13 - 24.
18. Sylvia S, Baluswami M, Vijaya Partha Sarathy MD, Krishnamurthy V. Effect of liquid seaweed fertilizers extracted from *Gracilaria edulis*, *Sargassum wightii* and *Ulvalactuca* on the growth and yield of *Abelmoschus esculentus*. *Indian Hydrobiology.* 2005;7: 69- 88.

19. Iswariya, S., K. Sujatha and Subhashini, R. Enhancement of seedling vigour through bio-priming for barnyard millet var. MDU 1. Int. J. Curr. Microbiol. App. Sci. 2019; 8(04):2254-2259.
20. Suma N. Studies of seed quality enhancement techniques in sesamum (*Sesamum indicum* L.) cv. Co-1 in sesamum. M.Sc. Thesis, Tamil Nadu Agricultural University, Coimbatore; 2005.
21. Sujatha K, Vethanayagi G, Ramamoorthy K. Studies on the effect of seaweed extracts on storage potential in bhendi seeds. Seaweed Res. Utiln. 2011; 33(1&2):125-131.
22. Butkhup L, Samappito S. *In vitro* free radical scavenging and antimicrobial activity of some selected thai medicinal plants. Res. J. Med. Plant., 2011;5:254–265.
23. Ambika S, Sujatha K. Organic seaweed nano powder effect on growth and yield attributes of pigeonpea. Legume Research. 2016;40(4):731-734.
24. Balaji DS, Sathitya Narayana G. Effect of various bio priming seed enhancement treatment on seed quality in certain minor millets. Plant Archives. 2019;19(1):1727-1732.
25. Bharathi A, Nateshan P, Vanangamudi K, Sherin PS, Ramya M, Thangavelu P. Conceptual and utility differences among seed enhancement technologies viz., seed pelleting, seed coating and seed coloring. ICAR short course on seed hardening and pelleting technologies for rainfed/garden land ecosystems. Tamil Nadu Agricultural University, Coimbatore. 2003;131.
26. Blunden G. Agricultural uses of seaweeds and seaweed extracts. In: Seaweed resources in Europe: Uses and potential. M. D. Guiry and G. Bluden (eds). Wiley, Chicester. 1994;65- 81.
27. Chen, Zijun. "Effects of biochar-based seed coating binders on seed germination and radicle extension; 2021.
28. Rathore SS, Chaudhary DR, Boricha GN, Ghosh A, Bhatt BP, Zodape ST, Patolia JS. Effect of seaweed extract on the growth, yield and quality of soybean (*Glycine max*) under rain fed conditions. South African Journal of Botany, 2009; 75:351-355.
29. Saletnik B, Bajcar M, Zagula G, Saletnik A, Tarapatsky, M, Puchalski C. biochar as a stimulator for germination capacity in seeds of Virginia mallow (*Sida hermaphrodita* (L.) Rusby). Applied Sciences. 2019;9(16):3213.
30. Shashibhaskar MS. Influence of seed pelleting on field performance and storability of tomato (*Lycopersicon esculentum* Mill.) cv. PKM-1. M.Sc. (Ag.) Thesis, University of Agricultural Sciences, Dharwad; 2008.

© 2023 Elamparithi and Sujatha; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/107752>