



A Study on the Hepatoprotective Potential of Methanol Seed Extract of (*Carica papaya*) Pawpaw in CCl₄ Induced Hepatic Damage in Wistar Rats

Elendu Melford Uche ^{a*}, Ejiofor Dominic Chinedu ^a,
Azunna Uchenna ^b, Ogwo Elisha Uko ^b
and Ubani Clement Dimgba ^b

^a Department of Human Physiology, Faculty of Basic Medical Sciences, Imo State University, Owerri, Imo State, Nigeria.

^b Department of Human Physiology, Faculty of Basic Medical Sciences, Abia State University, Uturu, Abia State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc. are available here: <https://www.sdiarticle5.com/review-history/113679>

Original Research Article

Received: 26/12/2023

Accepted: 02/03/2024

Published: 12/03/2024

ABSTRACT

The liver performs a variety of crucial metabolic processes, including being a hub for the metabolism of nutrients and the excretion of waste products. The liver plays a vital role in life. Therefore, the purpose of this study was to evaluate the hepatoprotective efficacy of *Carica papaya*

*Corresponding author: E-mail: Melfordue123@gmail.com;

seed extract on hepatic damage induced with CCl₄. *Carica papaya* seeds obtained from ripe pawpaw fruit were processed into fine powder and subsequently extract. Thirty adult male Wistar rats were divided into five groups of five rats each. Groups II-VI were administered carbon tetrachloride (2 mL/kg body weight) and olive mixed in the ratio of 1:1 to induce acute liver injury. Group I was the normal control and was administered 1 mL/kg of distilled water. Group II was not administered CP extract. Group III was administered 100 mg/kg bw of CP extract. Group IV was administered 200 mg/kg bw of CP extract. Group V was administered 300 mg/kg bw of CP extract. Group VI was administered 100 mg/kg bw of standard drug (Silymarin). Animal body weight was determined twice i.e., at the commencement the study and at the completion of the study. Administration of carbon tetrachloride significantly ($P < 0.05$) raised the activities of the liver enzymes i.e., Aspartate aminotransferase (AST), Alanine transaminase (ALT) and Alkaline phosphatase (ALP). However, oral administration of CP seed extract significantly ($P < 0.05$) lowered the activities of the said enzymes though to level which were significantly ($P < 0.05$) higher than that reported for the normal control. A contrary observation was however made on the activity of AST. The body weight of rat at the completion of the experiment was significantly ($P < 0.05$) higher than that reported at the commencement of the study. Meanwhile, a contrary observation was made on group II. In conclusion, it can be deduced from this study that that methanol seed extract of *Carica papaya* may have the ability to ameliorate chemically induced hepatic damage.

Keywords: Liver; enzymes; *Carica papaya*; hepatoprotective; silymarin.

1. INTRODUCTION

The liver is the most vital and largest solid organ of the body saddled with the task of nutrient metabolism as well as evacuation of metabolic waste products [1]. Primarily, it is concerned with the control of the flow as well as the safety of substances absorbed from the gastrointestinal tract (GIT) prior to distribution of substances to systemic circulation [2]. Thus, a complete loss of liver function could translate to death within a short time, thereby demonstrating the significance of liver to life" [3].

The harmful effects of ambient particulate matter or carbon black on the liver are well documented. Several animal models have clearly shown that exposure to PM or CB can cause direct hepatotoxicity [4]. Incidence of hepatic damage resulting from exposure to particulate matter is more prevalent in developing countries owing to the ever-increasing human population and consequent demand to meet human need through industrialization in a system with a wobbling environmental laws and attendant weak adherence which culminate to an alarming level of air pollution and its attendant consequences notably liver damage among others [5].

Plant-based therapeutic options remain the pivot upon which trado-medical practices rest since prehistoric times. In fact, an estimated 80% of the global population rely on plant-based therapeutic substances to meet their basic health care needs [6]. This could be attributed to the

fact that they are relatively safe for consumption in addition to being affordable among numerous other advantages [7].

Carica papaya popularly known as papaya has been used in the treatment of diverse human ailments. It is a member of the *Caricaceae* family. Its cultivation predominates in the tropics, sub-tropical and temperate zones such as Australia, Brazil, China, Hawaii, Nigeria etc [8];[9]. It owes its therapeutic potential to its rich phytochemical contents. Major phytochemicals present in the plant include tannins, steroids, terpenoids, saponins, phenols, flavonoids etc. [10]. Evidently, *Carica papaya* fruit pulp extract has been relied upon to protect against CCl₄ induced hepatic toxicity in rats [11]. Similarly, *C. papaya* root extract had also been used to protect against sodium arsenate induced hepatic damage in mice [12]. Unfortunately, these plant parts are not sustainable as the fruit serves mainly as food while the root plays key role to the survival of the plant itself. Thus, the need to evaluate the *Carica papaya* fruit seed which could be considered irrelevant to human as food or are not required by the plant to survive becomes imperative.

2. MATERIALS AND METHODS

2.1 Collection of Plant Material

Ripe fruits of *Carica papaya* (pawpaw) were procured from Eke Market, Afikpo, South Eastern Nigeria. They were identified and authenticated

at the herbarium unit of the Department of Forestry, Micheal Okpara University of Agriculture, Umudike, South Eastern University.

2.2 Extraction

Pawpaw fruits were dissected and the seeds collected were washed using clean water get rid of debris. The clean seeds were subsequently sun-dried for 48 hr and then oven-dried at 40°C for 1 week to ensure complete dryness. With the aid of an electric blender, dried seeds were thoroughly ground into fine power. Exactly 500 g of the papaw seed powdered sample was wrapped in a thimble and placed in a 500 cm³ Soxhlet extractor (M&G Scientific Co., England). The sample was Soxhlet extracted following standard analytical laboratory method at 60°C in methanol for 72 h. The extract was evaporated to a paste form at 40°C for 8 hr to produce 36% yield of raw sample of *C. papaya* seed extract.

2.3 Animals

Adult male Wistar rats weighing between 150 and 180 g were purchased from the Animal House of the Department of Science Laboratory Technology, Akanu Ibiam Federal Polytechnic Unwana, Afikpo. The animals were housed in transparent plastic cages and fed rat pellets. They were allowed access to water *ad-libitum* with a 12-h dark-light cycle. The animals were allowed an acclimatization period of 14 days prior to the commencement of extract administration.

2.4 Acute Toxicity Test

Acute toxicity test was determined in line with the description of Lorke [13]. In this test, two phases were involved. At the initial phase, nine (9) adult male rats were divided into three (3) groups of three rats each and were subsequently administered 10, 100 and 1000 mg/kg b.w. of extract respectively. This was followed by a close monitoring of the animals for manifestations of signs of toxicity for a period of 24 hrs in the absence of which the second phase was initiated with three adults male Wistar rats which were divided into three groups of one rat per group and were administered 1600, 2900 and 5000 mg/kg b.w of extract orally respectively. the animals were observed for 24 hours for clinical signs of toxicity. The LD₅₀ was calculated as the geometric mean of the non-lethal dose and lowest lethal dose.

2.5 Induction of Hepatic Damage

Carbon tetrachloride (2 mL/kg body weight) and olive were mixed in the ratio of 1:1 to induce acute liver injury in rats [14]. Olive oil was used as the vehicle and treatment lasted for 28 days.

2.6 Animal Grouping

Group I (Normal Control): Animals were orally administered 1 ml of distilled water daily and olive oil (2 ml/kg body weight)

Group II: Animals were induced hepatic damage without treatment

Group III: Animals with hepatic damage treated with 100 mg/kg of CP extract and olive oil (2 ml/kg body weight)

Group IV: Animals with hepatic damage treated with 200 mg/kg of CP extract and olive oil (2 ml/kg body weight).

Group V: Animals with hepatic damage treated with 300 mg/kg of CP extract and olive oil (2 ml/kg body weight).

Group VI: Animals with hepatic damage treated with 100 mg/kg bodyweight of silymarin and olive oil (2 ml/kg body weight).

2.7 Determination of Body Weight

With the aid of an electronic weighing scale, the weights of the animals were determined at the beginning of the experiment and at its termination after which the animals were humanely sacrificed and samples collected for analysis.

2.8 Biochemical Analysis

Sample preparation: In order to perform liver function tests, exactly 2 mL of blood introduced into an EDTA tube and subsequently centrifuged at 4,000 rpm for 15 min and the plasma obtained was used for biochemical analysis. Kits were used to determine the activity of aspartate aminotransferase (AST), alanine transaminase (ALT), alkaline phosphatase (ALP).

2.9 Histopathological Studies

Liver tissue harvested was preserved in 4% formaldehyde solution, embedded in paraffin wax, and subsequently sliced. Sliced sections were fixed on slides and stained with H&E

staining. The tissue sections were observed under a microscope (×400) to identify changes in the hepatocyte [15].

2.10 Data Analysis

Data generated were expressed as Mean ± Standard Deviation using SPSS (Ver. 23). Data were analysed using one way Analysis of Variance (ANOVA). Differences in mean were compared using Turkey Test. *p-values* less than 0.05 was considered statistically significant.

3. RESULTS AND DISCUSSION

The liver is one the critical organs of the body saddled with the task of detoxification of xenobiotics. It is also highly susceptible to damage through exposure to diverse environmental pollutants. Table 1 shows the hepato-protective effect of methanol seed extract of *Carica papaya* in Wistar rats. Exactly 2 ml of CCl₄ was orally administered to induce hepatic damage. However, following oral administration *Carica papaya* seed extract, there was a significant (*P*<0.05) reduction in the activity of aspartate amino transaminase compared to the activity reported for the untreated control (group II). It is important to note that groups III and IV administered 100 and 200 mg/kg of extract of *Carica papaya* seed was not significantly (*P*>0.05) different from that reported for the standard control (group VI) administered 100 mg/kg of silymarin (standard drug). On the

activity of ALT, the reported activity for the normal control (group 1) was not significantly (*P*>0.05) different from that reported for the standard control (group VI) but significantly (*P*<0.05) lower than that reported for the negative control (group II). However, the activity of ALT reported for groups III, IV and V administered 100, 200 and 300 mg/kg of *Carica papaya* extract was not significantly (*P*<0.05) different but was significantly (*P*<0.05) lower than the activity of ALT reported for group II and higher than that reported for the normal control (group I). Similar observation was made on the activity of the alkaline phosphatase (ALP). Observed decline in the activity of the liver enzymes following oral administration of extract could be attributed to the presence of phytochemicals inherent in the seed. This observation is consistent with the finding of Shuban et al. [11] who established the prophylactic and restorative effect of *C. papaya* Linn pulp water extract against CCl₄ induced hepatic damage. It is also in tandem with the outcome of a work done by Ojo et al. [12] which showed that 150 mg/kg of aqueous extract of pawpaw root protected against arsenate induced damage. Table 2 shows the body weight of rats with hepatic damage administered with *Carica papaya* seed extract indicating that although hepatic damage could cause a marked reduction in the weight of the experimental animals, treatment with the aforementioned extract resulted in a significant (*P*<0.05) increase in the body weight of rats.

Table 1. Effect of *Carica papaya* seed extract on serum hepatomarkers

Treatment	AST (U/l)	ALT (U/l)	ALP (U/l)
Group I (Normal control)	9.90±2.30 ^a	15.00±5.97 ^a	55.00± 3.24 ^a
Group II (Negative control)	42.10±3.10 ^d	29.04±3.20 ^d	115.00±6.20 ^c
Group III (100 mg/kg.)	10.00±3.02 ^a	21.00±1.21 ^c	66.00±2.20 ^b
Group IV (200 mg/kg)	10.20±1.32 ^a	20.01±1.23 ^c	64.00± 1.43 ^b
Group V (300 mg/kg)	10.00±4.23 ^a	19.00±2.32 ^c	60.00± 1.65 ^b
Group VI Std drug (silymarin)	10.05±2.10 ^a	15.2.00±2.10 ^a	56.00±9.74 ^a

Results are expressed as mean ± standard of five determination. Values with the same superscript in a column are not significantly (*p*<0.05)

Table 2. Body weight of rats with hepatic damage treated with *Carica papaya* extract

Treatment	Initial wt (g)	Final wt (g)
Group I (Normal control)	125±21.87 ^a	157±21.00 ^b
Group II (Negative control)	161±2.52 ^b	135±2.56 ^a
Group III (100 mg/kg.)	153±2.18 ^a	165±2.05 ^b
Group IV (200 mg/kg)	165±2.26 ^a	177±5.36 ^b
Group V (300 mg/kg)	163±2.36 ^a	178±21.65 ^b
Group VI Std drug (silymarin)	130±4.20 ^a	135±4.55 ^b

Results are expressed as mean ± standard of five determination. Values with the same superscript in a column are not significantly (*p*<0.05)

4. CONCLUSIONS

In order to maximize the benefits of pawpaw consumption, and to shift attention from dependence on certain vital parts of the plant for use as medicine, it is imperative to explore the efficacy of some of its relegated parts such as the seed. Thus, it can be deduced from this study that that methanol seed extract of *Carica papaya* may have the ability to ameliorate chemically induced hepatic damage.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ozougwu JC, Eyo JE. Hepato protective effects of *Allium cepa* extracts on paracetamol-induced liver damage in rat. African Journal of Biotechnology. 2014; 13(26):2679 -2688.
2. Allen SE. The liver: Anatomy, Physiology, Disease and Treatment. North Eastern University Press, USA; 2002.
3. Ozougwu JC. Comparative hepatoprotective and antioxidant effects of *Allium cepa*, *Allium sativum* and *Zingiber officinale* methanolic extracts against paracetamol-induced liver damage in *Rattus norvegicus*. Ph.D Research Thesis, Department Of Zoology and Environmental Biology, University of Nigeria, Nsukka. 2014;222.
4. Bourdon JA, Saber AT, Jacobsen NR, Jensen KA, Madsen AM, Lamson JS, Wallin H, Møller P, Loft S, Yauk CL, Vogel UB. Carbon black nanoparticle instillation induces sustained inflammation and genotoxicity in mouse lung and liver. Part. Fibre Toxicol. 2012;9:5.
5. Son JY, Lee JT, Kim KH, Jung K, Bell ML. Characterization of fine particulate matter and associations between particulate chemical constituents and mortality in Seoul, Korea. Environ. Health Perspect. 2012;120:872-878.
6. Singh SP, Kumar S, Mathan SV, Tomar MS, Singh RK, Verma PK, Kumar A, Kumar S, Singh RP, Acharya A. Therapeutic application of *Carica papaya* leaf extract in the management of human diseases, Daru. 2020;28(2):735–744. DOI:10.1007/s40199-020-00348-7
7. Santana LF, Inada AC, Espirito Santo BLSD, Filiú WFO, Pott A, Alves FM, Guimarães RCA, Freitas KC, Hiane PA. Nutraceutical potential of *Carica papaya* in metabolic syndrome, Nutrients. 2019; 11(7): 1608. DOI:10.3390/nu11071608
8. Nguyen TT, Shaw PN, Parat MO, Hewavitharana AK. Anti-cancer activity of *Carica papaya*: A review. Mol. Nutr. Food Res. 2013;57(1):153–164. DOI:10.1002/mnfr.201200388
9. Afzan A, Abdullah NR, Halim SZ, Rashid BA, Semail RH, Abdullah N, Jantan I, Muhammad H, Ismail Z. Repeated dose 28-days oral toxicity study of *Carica papaya* L. leaf extract in Sprague Dawley rats, Molecules. 2012;17(4): 4326–4342. DOI:10.3390/molecules17044326
10. Ghaffarilaleh V, Fisher D, Henkel R. *Carica papaya* seed extract slows human sperm. J. Ethnopharmacol. 2019;241:111972. DOI:10.1016/j.jep.2019.111972
11. Shaban NZ, Awad OM, Fouad GM, Hafez AM, Abdul-Aziz AA, El-Kot SM. Prophylactic and curative effects of *Carica papaya* Linn. pulp extract against carbon tetrachloride-induced hepatotoxicity in male rats, Environ. Sci. Pollut. Res. Int. 2023;30(10):27815–27832. DOI:10.1007/s11356-022-24083-5
12. Ojo OA, Ojo AB, Awoyinka O, Ajiboye BO, Oyinloye BE, Osukoya OA, Olayide II, Ibitayo A. Aqueous extract of *Carica papaya* Linn. roots potentially attenuates arsenic induced biochemical and genotoxic effects in Wistar rats, J. Tradit. Complement. Med. 2017; 8(2):324–334. DOI:10.1016/j.jtcme.2017.08.001
13. Lorke D. A new approach to practical acute toxicity testing, Arch. Toxicol. 1983;54:275-287. Available:https://doi.org/10.1007/BF01234480
14. Dmitry Frank, Shiri Savir, Benjamin F. Gruenbaum, Israel Melamed, Julia Grinshpun, Ruslan Kuts, Boris Knyazer, Alexander Zlotnik, Max Vinokur, Matthew boykoinducing acute liver injury in rats via carbon tetrachloride (ccl4) exposure

- through an orogastric tube. J Vis Exp; 2021 April 28.
15. Justin T, Raja A, TR, Janakiraman U, Manivasagam T. Neuroprotective effect of hesperidin on aluminium chloride induced Alzheimer's disease in wistar rats. Neurochemistry Research. 2015;40(4): 767–776.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/113679>