



Unlocking Nature's Secret: Microbiological and Nutrient Profiling of Fermented Apple

**Omodara, Tolani Rachael^a, Ajayi, Olayemi Elizabeth^{a*}
and Kotila, Oluwasefunmi Omolabake^a**

^a *Department of Microbiology, Faculty of Science, Ekiti State University, Ado-Ekiti, P.M.B. 5363, Ado-Ekiti, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/BJI/2024/v28i2714

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/114242>

Original Research Article

Received: 08/01/2024

Accepted: 12/03/2024

Published: 26/03/2024

ABSTRACT

Aim: This study aimed to investigate the microbiological and nutrient composition of naturally fermented apple cider.

Place and Study Duration: Department of Microbiology, between June and August 2023.

Methodology: The purchased apples were weighed, cut into dice, cleaned, and incorporated into distilled water. Subsequently, they were allowed to ferment spontaneously at room temperature in a 500 mL conical flask. The sample was inoculated onto sterilized glucose yeast carbonate agar and subsequently incubated at 30°C for 48 hours. The colony characteristics of the isolates on glucose yeast carbonate agar were observed. Gram's staining and other standard biochemical tests were performed to identify the isolates. During the fermentation period, the pH of the sample was determined. The proximate content (moisture, protein, crude fat, and carbohydrate) of the apple cider was also analyzed.

*Corresponding author: E-mail: oluwagbohunmi.ao@gmail.com;

Results: Microscopy revealed the isolates to be Gram-negative bacteria with rod-shaped morphology. The growth characteristics of the isolates on selective media and the results of biochemical tests suggested the presence of *Acetobacter* sp. The pH of the sample tends towards 4.0, suggesting the production of acidic compound. The apple cider produced had a high moisture content (78.04%), and the protein content was 20.04%. The results showed low concentrations of ash content (0.49%), carbohydrate (0.44%), and lipid (0.99%).

Conclusion: The presence of macro and micronutrients in apple cider has been associated with some positive health effects, which are enhanced through fermentation, making them a valuable option for health-conscious consumers.

Keywords: Apple cider; *Acetobacter* sp; acetic acid bacteria; fermentation.

1. INTRODUCTION

Apples (*Malus sylvestris*) are a significant economic fruit species belonging to the Rosaceae family, widely consumed globally for their appealing taste, juiciness, vibrant color, pleasing texture, and nutritional value [1]. Apples, being easily preserved, remain available year-round at affordable prices, and their versatility allows for the production of various apple products through different processing technologies [2,3]. These products include juices, dehydrated, canned, or purées, as well as fermented items like cider and fermented apple juices, all utilizing apple pomace as industrial by-products [4]. Fermentation, a transformative process facilitated by microorganisms under specific conditions, is pivotal in the production of fermented apple products. The success of fermentation is influenced by factors including the microorganism(s) utilized, culture medium, processing conditions, and stages of product recovery [5,6]. Control of pH, temperature, humidity, aeration, medium bed thickness, and agitation speed is essential for effective fermentation [7,8]. Apples are recognized for their high content of bioactive compounds, particularly flavonoids like quercetin and its derivatives, with antioxidant, anti-inflammatory, antimicrobial, anti-depressive, and anticarcinogenic properties. Fermentation impacts the properties of apple products, inducing changes in composition, nutritional value, and organoleptic characteristics. According to findings from the study conducted by [9], certain fermented apple products undergo considerable changes in chemical composition, sensory profiles, and bacterial counts. These observations imply their potential to serve as functional, non-dairy probiotic beverages [10]. The concentration and variety of bioactive molecules in apples may vary depending on factors such as species, cultivar, climatic conditions, agronomic practices, and processing methods. Fermentation further enhances the

health benefits of apple products, making them a valuable option for health-conscious consumers. This study aims at the bench-scale production of apple cider and the proximate analysis of the naturally fermented cider.

2. MATERIALS AND METHODS

Apples were purchased from the Erekesan market in Ado-Ekiti, Ekiti state, Nigeria. Glucose yeast carbonate (GYC) agar, formulated with glucose 4%, yeast extract 1%, CaCO₃ 1%, and agar 1.5%, was prepared according to [11].

2.1 Preparation of Apple Cider

Twenty grams (20 g) of the apple were weighed, diced, washed, and soaked in distilled water. Subsequently, they were left to undergo natural fermentation at room temperature within a 500 mL conical flask. Additionally, at each stage, 1 ml of the sample was serially diluted and cultured in GYC agar medium for further analysis.

2.2 Isolation of Acetic Acid Bacteria from Naturally Produced Cider

Aseptically, 0.1 ml of diluted samples were transferred and spread onto GYC agar plates using a sterile bent rod and incubated at 30°C for 48 hours. Well-formed colonies with a single morphology were isolated and sub-cultured using the streak plate technique on GYC agar plates and then incubated at 30°C for 48 hours.

2.3 Microscopic and Biochemical Characterization of Acetic Acid Bacteria

The colony characteristics of the isolates on GYC agar were observed. Gram staining was performed for morphological characteristics of isolated strains. For the characterization of isolates, Gram staining, sugar fermentation tests, catalase tests, and motility tests were conducted.

2.4 Determination of pH

The pH was determined according to methods of [12]. 1 mL of the cider was introduced on pH paper and the colour change was compared with the standard. pH measurements were taken at intervals of 0, 24, 48, and 72 hours, as well as on day 5, week 1, week 3, week 4, and week 5.

2.5 Proximate Analysis of Natural Fermented Apple Cider

The proximate components of the Cider, including protein, moisture, crude fiber, crude fat, and ash, were analyzed using standard analytical procedures. Specifically, the protein content was determined following [12] method 960.52, while ash, moisture content, and total fat were determined according to [12] methods 923.03, 934.01, and 963.15, respectively. The percentage of carbohydrates was calculated as follows:

$$\begin{aligned} \% \text{Carbohydrate} &= 100 \\ &- (\% \text{protein} + \% \text{moisture} \\ &+ \% \text{crude fat} + \% \text{ash}) \end{aligned}$$

3. RESULTS AND DISCUSSION

3.1 Cultural and Microscopic Characterization of Isolates

Five colonies were chosen from the culture plate based on their cultural characteristics,

and designated as FAC1, FAC2, FAC3, FAC4, and FAC5. The cultural characteristics of isolated strains are described and represented in Table 1. FAC1, FAC2, FAC3, and FAC4 were filamentous, whereas FAC5 was irregular in shape. All isolates exhibited a creamy colony color and a viscid consistency. The isolates were identified as Gram-negative or Gram-variable (isolates FAC3 and FAC5) small rods, exhibiting straight or slightly curved shapes.

3.2 Biochemical Characterization

The selected isolates tested positive for catalase and were initially identified as Acetic acid bacteria following the standard guidelines outlined in Bergey's Manual of Determinative Bacteriology [13]. Table 2 also illustrates the utilization of selected carbohydrates by the five isolates presumed to be acetic acid bacteria (AAB). All isolates demonstrated the ability to ferment glucose with acid production, whereas none utilized sucrose.

3.3 pH Value during Fermentation

The pH decreased from 7.0 to 5.0 after day 3, maintaining this value until week 4, at which point it dropped to 4.0 as shown in Fig. 1. The pH value of 4.0 was sustained in week 5.

Table 1. Cultural and microscopic characteristics of isolates

Isolates	Shape	Consistency	Colony color	Gram Stain	Shape	Motility
FAC 1	Filamentous	Viscid	Creamy	-ve	Small rods	+
FAC 2	Filamentous	Viscid	Creamy	-ve	Rods	+
FAC 3	Filamentous	Viscid	Creamy	-ve	Curved rod	+
FAC 4	Filamentous	Viscid	Creamy	-ve	Rods	+
FAC 5	Irregular	Viscid	Creamy	-ve	Rods	+

Table 2. Biochemical characterization of fermented apple cider isolates

Isolates	Catalase	Utilization of sugar			Organisms
		Glucose	Lactose	Sucrose	
FAC 1	+	+	-	Acid and gas produced	<i>Acetobacter</i> sp
FAC 2	+	+	-	Acid and gas produced	<i>Acetobacter</i> sp
FAC 3	+	+	-	Acid and gas produced	<i>Acetobacter</i> sp
FAC 4	+	+	-	Acid and gas produced	<i>Acetobacter</i> sp
FAC 5	+	+	-	Acid and gas produced	<i>Acetobacter</i> sp




Fig. 1. pH of fermented apple cider at different days of fermentation

Table 3. Proximate composition of fermented apple cider in percentage

Proximate	%
Moisture	78.04
Crude fat	0.99
Protein	20.04
Ash	0.49
Carbohydrate	0.44

3.4 Proximate Composition of the Apple Cider

The apple cider produced showed a high concentration of moisture (78.04%). The protein content was 20.04%. The results showed a low concentration of ash content (0.49%), carbohydrate (0.44%), lipid (0.99%), and no fiber content as outlined in Table 3.

3.5 Discussion

The microscopic and biochemical characterization revealed that the isolates are Acetic acid bacteria (AAB) and identified as *Acetobacter* spp. The main species responsible for apple cider production belong to the genera *Acetobacter*, *Gluconacetobacter*, *Gluconobacter*, and *Komagataeibacter*, owing to their capacity to oxidize ethanol and their tolerance to acetic acid released during fermentation [14]. The pH is a crucial parameter in fermentation processes as it affects the growth and activity of microorganisms. The gradual increase in the acidic condition of the fermented apple suggests the production of acidic compounds. The acidity

of cider primarily stems from the production of organic acids, including acetic acid and tartaric acid, by the fermenting microorganisms [15]. These fastidious organisms and strains tend to lose their capacity to produce high concentrations of acetic acid when sub-cultured [16]. This can be attributed to the reduction in the oxygen concentration available and decrease in pH due to continuous acetic acid production, a primary metabolite [14].

The analysis of the proximate content of the apple cider as shown in Table 3, suggests that apple cider contains high moisture content, this aligns with the findings of [17]. High moisture content is indeed a significant factor that affects the flavor of juices and can result in reduced shelf stability [18, 6]. Ash content signifies the mineral content of the food sample, as minerals do not combust and are left behind in the form of ash. Therefore, by measuring the ash content, one can gain insights into the types and quantities of minerals present in the food product [18]. The ash content of cider produced in this study shows similar result to that of [19] who recorded the ash content of a given apple juice

sample in the range of 0.045-0.17%. The lower ash content also indicates the minimized quantity of apple fruit used to prepare this cider. Long storage at different temperatures also affects the quality parameters of the juices [20].

The percentage of carbohydrate was recorded to be low with a value of 0.44%. Carbohydrates account for the principal food value of apples. They are starches, sugars, pectin, cellulose, and hemicelluloses [21]. The study [22] also reported (3.6%) total sugar in apple juices. Pectin is a component of carbohydrates in apple cider that has shown additional health benefits on digestive systems [17]. Differences in the result obtained from this study and those of other researchers may be attributed to variations in pectin content. The pectin content of different apple cultivars grown in diverse locations influences the carbohydrate composition of cider and other apple juices [23]. The lipid content in the apple cider sample was low which is common for fruits. The elevated protein content in the cider could stem from the presence of apple pomace. Apple pomace serves as a source from which valuable components like pectin, aroma compounds, edible fibers, and antioxidant polyphenols can be extracted. It can also be utilized to obtain protein-enriched feeds, synthesize pectolytic enzymes, or produce natural aroma compounds through fermentation processes [6].

4. CONCLUSION

Acetobacter sp. have been associated with the fermentation of apples based on their cultural and biochemical characteristics. The pH of the sample tends toward 4.0, suggesting the production of acidic compound. The apple cider proximate contents are moisture (78.04%), protein content 20.04%, ash content (0.49%), carbohydrate (0.44%), and lipid (0.99%). The presence of these macro and micronutrients have been linked to the positive health effects of apple-fermented products therefore, further research is needed to explore the specific mechanisms behind the health benefits of cider and establish optimal consumption recommendations.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Food and vegetable processing (FAO Agricultural Services Bulletin No. 119).

2. Bilková A, Bad'urová K, Svobodová P, Vávra R, Jakubec P, Chocholouš P, et al. Content of major phenolic compounds in apples: Benefits of ultra-low oxygen conditions in long-term storage. *J. Food Compos. Anal.* 2020;92:103587.
3. Li Y, Zhang X, Nie J, Bacha S, Yan Z. et al. Occurrence and co-occurrence of mycotoxins in apple and apple products from China. *Food Control.* 2020;118:107354.
4. Madrera, R, Bedriñana, R, Valles, B. Enhancement of the nutritional properties of apple pomace by fermentation with autochthonous yeasts. *LWT-Food Sci. Technol.* 2017;79:27–33.
5. García C, Rendueles M, Díaz L. Liquid-phase food fermentations with microbial consortia involving lactic acid bacteria: A review. *Food Res. Int.* 2019;119:207–220.
6. Guiné R, Barroca M, Coldea T, Bartkiene E, Anjos O. Apple fermented products: An overview of technology, properties and health effects. *Processes.* 2021;9(223):1-25.
7. Peris M, Escuder-Gilabert L. On-line monitoring of food fermentation processes using electronic noses and electronic tongues: A review. *Anal. Chim. Acta.* 2013;804:29–36.
8. Reihani S, Khosravi-Darani K. Influencing factors on single-cell protein production by submerged fermentation: A review. *Electron. J. Biotechnol.* 2019;37:34–40.
9. Peng B, Yue T, Yuan Y. Analysis of key aroma components in cider from Shaanxi (China) Fuji apple. *Int. J. Food Sci. Technol.* 2009;44:610–615.
10. Roberts G, Spadafora N. Analysis of apple flavors: The use of volatile organic compounds to address cultivar differences and the correlation between consumer appreciation and aroma profiling. *Journal of Food Quality.* 2020;8497259:1-7.
11. Ezemba A, Osuala E, Orji M, Ezemba C, Anaukwu C. Production and comparative physicochemical analysis of vinegar from locally grown fruits in Nigeria and industrial produced vinegar. *American journal of microbiological research.* 2021;9(1):25-33.
12. Association of analytical chemists (AOAC). Official methods of analysis of AOAC. 17th ed. Washington DC; 2000.

13. De Ley J. Genus *Acetobacter*. Bergey's manual of systematic bacteriology. 1984;1:268-274.
14. Mathew B, Agrawal S, Nashikkar N, Bundale S, Upadhyay A. Isolation of acetic acid bacteria and preparation of starter culture for apple cider vinegar fermentation. *Advances in Microbiology*. 2019;9:556-569.
15. Ahmad I, Khalique A, Shahid M, Rashid A, Faiz F, Ikram, et al. Studying the influence of apple peel polyphenol extract fortification on the characteristics of probiotic yoghurt. *Plant*. 2020; 9(77):2-15
16. Mas A, Torija MJ, García Parrilla M, Troncoso A. Acetic acid bacteria and the production and quality of wine vinegar. *The Scientific World Journal*; 2014. Article ID: 394671, 3.
17. Esselen S, Fellers W. Analytical characterization of the aroma of five premium red wines. Insights into the role of odor families and the concept of fruitiness of wines. *Journal Agriculture Food Chemistry*. 2016;511:4501–4510.
18. Ekanem J, Ekanem O. Proximate analysis and sensory evaluation of freshly produced apple fruit juice stored at different temperatures and treated with natural and artificial preservatives. *Global journal of pure and applied sciences*. 2020;25:31-37.
19. Khan B, Ali S, Shahzad S, Basharat M, Ali J, Jabeen Z, et al. Postharvest quality of apple (*Malus domestica* Borkh) as influenced by storage conditions. *International journal of advanced research in biological sciences*. 2017;4(7):154-163.
20. Rai T, Saxena E. Effects of different factors on sensory attributes, overall acceptance and preference of Rooibos (*Aspalathus lineares*) tea. *Journal of Sensory Studies*. 2018;20:228–242.
21. Hayes S, Haddad R. Principles and practices of winemaking, Chapman and Hall, New York. 2016;455:56-67.
22. Yılmaz A, Gerçekçioğlu, Atasver Ö. Determination of pomological and chemical properties of some medlar (*Mespilus germanica* L.) genotypes. *J. New Results Sci*. 2016;11:118–124.
23. Bray U. Sensory and analytical study of rose sparkling wines manufactured by second fermentation in the bottle. *Journal of Agricultural and Food Chemistry*, 2014;52:6640–6645.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/114242>