



# Standardization of Quick Seed Viability Protocol for *Pinus roxburghii* Sarg. Using Tetrazolium Assay

Ravindra Kumar Dhaka <sup>a++\*</sup>, Priya Thakur <sup>a#</sup>,  
Navjot Singh Kaler <sup>b†</sup>, Som Dutt Sharma <sup>b‡</sup>,  
Chaman Negi <sup>c++</sup> and M. K. Brahmi <sup>d‡</sup>

<sup>a</sup> Department of Tree Improvement and Genetic Resources, COHF-Neri, Dr. Y. S. Parmar University of Horticulture and Forestry Nauni, Solan (HP), India.

<sup>b</sup> Department of Silviculture and Agroforestry, COHF-Neri, Dr. Y.S. Parmar University of Horticulture and Forestry Nauni, Solan (HP), India.

<sup>c</sup> Department of Social Sciences, COHF-Neri, Dr. Y. S. Parmar University of Horticulture and Forestry Nauni, Solan, (HP), India.

<sup>d</sup> Department of Forest Products, COHF-Neri, Dr. Y. S. Parmar University of Horticulture and Forestry Nauni, Solan (HP), India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/JABB/2024/v27i5760

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/114321>

Original Research Article

Received: 15/01/2024  
Accepted: 20/03/2024  
Published: 26/03/2024

## ABSTRACT

Chir pine (*Pinus roxburghii*) is one of the important tree species among five indigenous pine species of the Himalayan region and is widely distributed throughout the Himalaya. It is used as a commercial timber, fuel wood and pulpwood and also provide resin for various utilities. For the

<sup>++</sup>Assistant Professor;

<sup>#</sup> Research Scholar;

<sup>†</sup> Scientist;

<sup>‡</sup> Associate Professor;

\*Corresponding author: E-mail: [ravindrakrdhaka@gmail.com](mailto:ravindrakrdhaka@gmail.com);

ecological, silvicultural and economical significance, the regeneration of this species is required for restoration and conservation purposes. Before seed sowing in nursery, there is need to know the actual seed vigour and quality of seeds. Therefore, the present experiment was conducted at Dr. Y. S. Parmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh of India to standardized a quick seed viability testing protocol using tetrazolium (TZ) and confirmed with laboratory seed germination testing. The tetrazolium test and germination test were conducted with 100 seeds in 4 replications in CRD design as per the procedure devised by ISTA and analyzed with OPSTAT statistical software. After pre-moistening, chir pine seeds were cut longitudinally into two parts away from the radicle end to expose the embryo. The seeds of the species stained with 0.5, 1.0, 1.5 and 2.0 % solution of TZ and incubated at 25, 30 and 35 °C temperature in the darkness for 12, 24 and 36 hours. The result of TZ test showed that the seeds of chir pine soaked in 0.5 % tetrazolium solution at 25 °C temperature for 24 hours found to be the best one for the seed vigour and seed quality testing. For predicting the accurate seed viability, RMS (root mean square) method was used and recognized six staining seed categories distinguished 1 to 6. Seed category 1 denoted as viable seeds and category 6 as non-viable seeds. Result showed that the topographical staining categories 1, 2 and 3 provided least root mean square value and recommended these as viable seed categories for TZ seed viability test. Furthermore, it is also compared with laboratory seed germination test to standardize the seed viability protocol and result in line with the conformity. Overall, the results of the present study fulfilled all the gaps and criteria of quick seed viability testing in chir pine species and standardized protocol could be used for future seed vigour testing within a short period of time for the forest restoration and conservation purpose.

**Keywords:** *Chir pine; Himalaya; seed viability; tetrazolium test; seed quality; root mean square; germination test; seed vigour.*

## 1. INTRODUCTION

Chir pine (*Pinus roxburghii* Sarg.) is one of the important tree species of five indigenous pine species of the Himalaya, which is widely distributed an altitudinal range between 450-2300 m elevation in Afghanistan, Pakistan, India, Nepal and Bhutan between latitude range 26°N-36°N and longitude range 71°E-93°E [1]. It is of great ecological, silvicultural and economical significance tree species in the Himalayan region. It provides commercial timber, resin, fuel wood and pulpwood for muciferous end uses. Resin obtained from the trees by taping method is used in soap, pharmaceutical, paper and paint industries in India [2]. It is a fire resistant, strong light demander tree species and successful pioneer on exposed sites which can be planted on bare lands where it can withstand frost, drought, snow and free from serious pests and diseases. It has a wide variation in germination, growth, stem forming, resin yield and timber yield with grain angle throughout its natural forests in relation to altitude and site [3]. The artificial and natural regeneration of the species takes place through seeds. The pollination takes places in the spring season by air in this monoecious species and fertilization occurs in the next coming spring season and mature cones disperse their seeds in next or third spring season after pollination. Thus, all most the time

taken between pollination and seed maturity is 2 years [4]. Good seed year of chir pine is about 4-5 years, so seed can be collected only that year for the regeneration purpose. The seed germination and its attributes are not only irregular but also takes prolonged germination period which resulted uneven seedling growth and poor-quality seedlings [5]. The precise determination of seed viability prior to nursery germination and field sowing is of great importance for the regeneration of the species in the forest. The accurate quantity of seeds to be sown in nursery is mainly dependent upon the quality or viability of the seeds.

Seed viability testing in tree species is very important to know the quality of seeds [6] which has a great effect on the quality of planting stock. Therefore, it is imperative to test the viability of the seeds before commencing seed sowing in nursery or field. Seed germination test for most of the forest tree species takes places from 30 to 60 days after seed sowing [7]. Moreover, the germination test of pines species is also inadequate and unknow their true seed viability [8]. There is a need to know the non-viable seeds from viable seeds which is sometimes visually undifferentiated from dead seeds. Various tests have been used to estimate the seed viability, but they are laborious enough in workability [9]. One of the most accepted and reliable method of

seed viability is the tetrazolium test (TZ) which is also referred as “quick test” [10]. A topographical method of seed viability assessment was developed first for cereals [10] and later on used for seed viability test in coniferous species [11]. The tetrazolium test is also one most rapid and useful methods to evaluate seed quality of many species [12]. The historical aspects of the tetrazolium test from its development [13] to the applicability of 2,3,5-Triphenyl Tetrazolium Chloride (TTC) to the quick viability test in conifers species [14] for the restoration [15] and conservation ecology [16] as well as in endangered pine species [17]. Seed dormancy is common in most of the forest tree species [18,19]. Tetrazolium test (TZ) also provides a quicker knowledge to obtain seed viability even when a seed in a state of dormancy [20]. However, there are several factors which contribute to the loss in seed quality, viability and vigor may be either physical or physiological phenomenon [21]. Savonen (1999) developed an improved method of topographic tetrazolium testing in Scots pine to overcome the patchily stains in seeds [22] due to the effect of storage time, temperature and container on the viability of the seed of some pine species [23]. Daws et

al. (2006) also tried pressure with time dependency of vacuum degassing in 17 *Pinus* species to rapid the method for viability assessment using tetrazolium chloride [24]. Keeping in mind all the knowledge, the present experiment is conducted to standardize a quick seed viability testing protocol for the chir pine species in Himachal Pradesh of India.

## 2. METHODOLOGY

### 2.1 Seed Collection and Extraction

The cones of mature aged chir pine tree were collected from the five different tree seed stands in the Hamirpur Forest Division, Himachal Pradesh, India during the year of 2023 as geolocation presented in Fig. 1. After the collection of cones, they were sun dried for 2-3 days and followed the extraction, cleaning and processing of the seeds. Further, chir pine seeds were mixed thoroughly to maintain homogeneousness. These five seed lots were designated from A to E and stored in well labelled air tight plastic containers at room temperature ( $25^{\circ}\text{C}\pm 2$ ).




Fig. 1. Geolocation of cone collection of *Pinus roxburghii* during the course of study


## 2.2 Tetrazolium Seed Viability Testing

The procedure of the tetrazolium test (TZ) was followed as per devised by ISTA [13]. Four replications of 100 seeds were taken from the random samples after mixing some quantity of seeds from each seed lot of *Pinus roxburghii* and employed for each treatments given below to standardize optimum concentration of TTC solution, incubation temperature and time. These samples were soaked in distilled water for 24 hours at ambient room temperature before TZ staining to allow complete moistened of all the seed tissues. After pre-moistened, chir pine seeds were cut longitudinally into two parts away from the radicle end to expose the embryo. Seeds of the species were then stained with 0.5, 1.0, 1.5 and 2.0 % solution of TZ prepared in the double distilled water. The chir pine seeds were placed in the Petri dishes on the moistened double sheets of Watman filter paper No.1 to incubate with TZ solution at 25, 30 and 35 °C temperature in the darkness for 12, 24 and 36 hours. The 2,3,5-Triphenyl Tetrazolium Chloride (TTC) salt solution enters into the living tissue of the seed which produce a reddish or dark purple colour due to water insoluble compound called formazon by the activity of dehydrogenase enzyme [25]. After completion of incubation period, the TZ solution was drained out and chir pine seeds were washed in distilled water. Then, red coloured seeds were counted as viable or germinable seeds and non-coloured seeds as nonviable or non-germinable seeds. Then, the concentration of TZ solution, incubation temperature and time standardized on the basis of optimum seed viability results presented in Table 1. After that, the optimised concentration of TTC solution, incubation temperature and time was employed for all the seed lots of chir pine seed separately to standardize the stain topography of viable and non-viable seeds. Seed viability was evaluated according to the topographical staining pattern of the presence, location, extent, intensity of the colouration and nature of tissue staining in all the seed structures with the help of magnifying glass [8, 13]. The staining pattern of each seed structure was recorded under 6 staining categories depicted in Fig. 2 and details presented in Table 2.

## 2.3 Seed Germination Testing

Seed germination test was conducted in laboratory condition with 100 seeds in four replications taken from each seed lot to know the viability of seeds by performing the seed germination test. These chir pine seeds were

placed in the Petri dishes on the moistened double sheets of Watman filter paper No.1 as per divested by ISTA [13]. Before to sowing the seeds, they were soaked in distilled water for 7 days to overcome the seed physiological dormancy which is mainly found in conifers [26]. The germination data were recorded to calculate the germination percentage of chir pine seeds to test the viability or quality of seeds. The emergence of radicle i.e., newly developed root portion of the seedling reached upto the equal length of the seed was taken as the criterion for seed germination [27].



**Fig. 2. Staining topography category (1-6) for tetrazolium viability test in *Pinus roxburghii***

## 2.4 Statistical Analysis

The experimental data were subjected to OPSTAT statistical software package [28] for the statistical analysis and appropriate interpretation. ANOVA (Analysis of Variance) was derived using completely randomized design (CRD) with four replications in the laboratory condition. All the treatments were considered to be random and compared at 5 % level of significance. For predicting the accurate seed viability, the RMS (root mean square) method was used as described by Kuo et al. (1996) [29]. This method had been used for predicting the seed viability in *Pinus wallichiana* seeds [6] and recognized six staining categories distinguished 1 to 6. Seed category 1 denoted as viable seeds and category 6 as non-viable seeds. Then, total 15 combinations possibilities were found within the remaining 4 categories i.e., 2, 3, 4, 5 for the evaluation criteria. Root mean square (RMS) differences between actual germination percentage as determined by laboratory germination test (G) and potential viability percentage as determined by Tetrazolium test (P) for each evaluation as per the following equation:

$$RMS = \sqrt{\frac{1}{n} \sum_{i=1}^n (G_i - P_i)^2}$$

Where,  $G_i$  = Actual germination percentage of seed lots from 1 to n.  $P_i$  = Potential viability

percentage as determined by TZ staining of seed lots from 1 to n. n = Total number of seed lots in the study. The evaluation criterion used in the study that created least root mean square values was recommended for the Tetrazolium test of the species.

### 3. RESULTS AND DISCUSSION

#### 3.1 Standardization of Optimum Conditions for TZ Staining

The experimental data is presented in Table 1 with respect to know the optimum concentration of tetrazolium solution, soaking temperature and soaking time needed for maximum staining. All the treatments significantly ( $P < .05$ ) affected the

staining pattern in *Pinus roxburghii* seeds. The chir pine seed staining pattern decreased when the concentration of tetrazolium increased from 0.5% to 2.0%, however seed less stained when incubated at high temperature i.e. 35°C and for a longer period (36 hours). The light colour staining pattern is very difficult to differentiate viable and non-viable seeds. The seeds of chir pine soaked in a 2.0 and 1.5% TZ solution and incubated at 35° C temperature for 36 hours showed only 30 and 30.50% seed viability, respectively. The chir pine seed soaked at 25°C for 24 hours in 0.5% TZ solution indicated significantly high viability of 64.75% which is followed by 57.50% in seeds incubated at 25°C in 0.5% TZ solution for 36 hours.

**Table 1. Effect of incubation in concentration of TZ solution, temperature and soaking time on seed staining pattern in *Pinus roxburghii***

Sr. No.	Treatments			Seed viability percentage (%)	
	TZ conc. (%)	Soaking temp. (°C)	Soaking time (hours)	Mean	±S.E.
1	0.5	25	12	50.50	0.65
2	1.0	25	12	43.75	0.85
3	1.5	25	12	40.50	0.96
4	2.0	25	12	34.50	0.65
5	0.5	30	12	39.50	1.04
6	1.0	30	12	35.00	0.41
7	1.5	30	12	33.50	0.65
8	2.0	30	12	30.00	0.71
9	0.5	35	12	27.50	0.29
10	1.0	35	12	26.00	0.41
11	1.5	35	12	24.00	0.41
12	2.0	35	12	23.00	0.41
13	0.5	25	24	64.75	1.25
14	1.0	25	24	57.00	1.29
15	1.5	25	24	53.50	1.19
16	2.0	25	24	47.75	0.85
17	0.5	30	24	53.00	1.29
18	1.0	30	24	48.50	0.65
19	1.5	30	24	46.25	0.85
20	2.0	30	24	43.00	1.08
21	0.5	35	24	40.50	0.65
22	1.0	35	24	39.50	0.65
23	1.5	35	24	38.25	0.48
24	2.0	35	24	37.25	0.48
25	0.5	25	36	57.50	0.65
26	1.0	25	36	50.75	0.85
27	1.5	25	36	47.00	0.91
28	2.0	25	36	41.50	0.65
29	0.5	30	36	46.50	1.04
30	1.0	30	36	41.75	0.48
31	1.5	30	36	39.75	0.85
32	2.0	30	36	36.50	0.87

Sr. No.	Treatments			Seed viability percentage (%)	
	TZ conc. (%)	Soaking temp. (°C)	Soaking time (hours)	Mean	±S.E.
33	0.5	35	36	33.75	0.48
34	1.0	35	36	32.75	0.48
35	1.5	35	36	30.50	0.65
36	2.0	35	36	30.00	0.41
<b>Overall Mean of TZ test (%)</b>				<b>40.70</b>	
				C.D.	2.20
				±SE(m)	0.78
				C.V.	3.85
<b>Overall Mean of Germination test (%)</b>				<b>71.25</b>	

There was a big difference between the seed viability percentage by TZ staining pattern (mean= 40.70%) and laboratory germination (mean= 71.25%) with four replications of 100 seeds. On the other hand, chir pine seed soaked at 25°C for 24 hours in 0.5% TZ solution showed 64.75 % seed viability which is nearby the laboratory germination (71.25%). Thus, the above concentration of TZ solution, incubation temperature and time could be considered as an optimum for TZ staining in *Pinus roxburghii* seed and standardized for the future use for the seed viability testing in this species. The results obtained from the present study are configured with the tetrazolium test in *Pongamia pinnata* [30]; in *Moringa oleifera* [31]; in *Acer caesium* [27]; *Ulmus wallichiana* [27] and in *Pinus wallichiana* [8]. This is also corroborated with the ISTA rules of tetrazolium test [13]. Masullo et al. (2017) optimized the tetrazolium tests to assess the viability and quality of *Platymiscium floribundum*, *Lonchocarpus muehlbergianus* and *Acacia polyphylla* seeds for different concentration of tetrazolium solution, incubation time and temperature [32]. Thus, TZ solution of 0.5%, incubation at 25°C temperature for 24 hours standardized as an optimum condition for TZ staining in *Pinus roxburghii* seeds for the seed viability and quality testing.

### 3.2 Standardization of Staining Topography of Viable and non-Viable Seed

Tetrazolium (TZ) solution is a colourless before staining of seeds and when seeds are soaked in TZ, then it imbibed by the seed tissues of chir pine to interferes in the reduction progression of living cells and accept hydrogen ions from dehydrogenase which resulting in the production of red stable colour and non-diffusible substance i.e., Triphenyl Formazan in the living tissues. Then, it is possible to distinguish the red coloured living tissues of the seed from the colourless dead tissues. Based on this type of phenomenon, the topographical staining pattern of the presence, location, extent, intensity of the colouration and nature of tissue staining, the tetrazolium (TZ) stained seeds were classified into six staining categories as shown in the Fig. 2 and detailed in Table 2. Data presented in Table 2 showed that the seed viability ranging from 1.25 to 62.25% in the five staining topographical categories (1-5) on the basis of the stained area of radicle axes and cotyledons. The comparison of staining category 1 alone (mean= 55.15%) with laboratory germination (73.35 %) among all the seed lots showed that there was big difference in viability test. The similar results also observed by Aslam et al. (2010) in *Pinus wallichiana* seed viability test [8].

**Table 2. Staining topography of TZ test results and laboratory germination percentage for five seed lots of *Pinus roxburghii***

Staining topography category	Details of staining category	TZ viability percentage in five seed lots (%)					Mean
		A	B	C	D	E	
1	Embryo and cotyledon fully stained.	48.50	54.25	62.25	52.50	58.25	55.15
2	Embryo fully stained and minor unstained areas (i.e. less than 3/4) on cotyledon.	10.50	8.25	5.25	10.25	9.25	8.70

Staining topography category	Details of staining category	TZ viability percentage in five seed lots (%)					Mean
		A	B	C	D	E	
3	Embryo fully stained and less than 1/2 portion of the cotyledon unstained.	8.25	7.25	4.50	7.50	8.25	7.15
4	Embryo stained and stained patches on cotyledon.	5.25	3.25	2.25	4.75	2.75	3.65
5	Embryo unstained and stained patches on cotyledon.	4.25	2.50	1.25	3.50	2.25	2.75
6	Embryo and cotyledon unstained or stained in very small patches.	23.25	24.50	24.50	21.50	19.25	22.60
<b>Laboratory germination (%)</b>		<b>70.25</b>	<b>73.50</b>	<b>76.25</b>	<b>72.25</b>	<b>74.50</b>	<b>73.35</b>

However, this is an under estimate and to overcome it with the procedure of RMS (Root Mean Square) determination [27]. There were 15 possible combinations (Table 3) for RMS values of staining categories 2 to 5 and together with the fully viable staining category 1. Staining category 6 was clearly showed non-viable seeds as the radicle axes and cotyledons were fully colourless or unstained. The experimental results showed that staining categories from 1 to 3 included as viable seeds because of the category 2 showed a fully stained embryo and minor unstained areas on cotyledon, and staining category 3 showed a fully stained embryo and less than 1/2 portion of the cotyledon unstained. After pooling together, the RMS values of staining categories 1, 2 and 3 provide least root mean square value (18.89) and therefore, it is recommended to include staining categories 2 and 3 in the viable categories. However, the lowest RMS value of 13.47, 13.76, 4.43 and 14.60 showed when staining category 1 pooled with either on one of the categories 2 or 3 or 4 or 5. Staining category 4 stained only embryo and stained some patches on cotyledon. However, staining category 5 not stained embryo and stained only some patches on cotyledon. In the other combinations, the mean root square values were much higher as compared to the actual germination percentage which leading to faulty determination of seed viability.

Anatomically, staining categories 2 and 3 also represent normal cotyledons and TZ staining as described earlier by several authors [8,27]. Aslam et al. (2010) categories six different staining patterns to distinguish the difference between colouration, radicle germination and surface area of staining of blue pine (*Pinus wallichiana*) seeds for potential or real seed viability testing. They also reported that the pooling together the root mean square (RMS) values of staining categories 1, 2 and 3 contributed the least RMS value which is justified to include staining categories 2 and 3 as in viable categories. These staining patterns are also anatomically visible [8]. The staining category 4 and 5 included in non-viable seeds due to they are anatomically non-visible. Phartyal et al. (2003) also categories six staining topography in *Acer caesium* seed viability test and results showed that staining categories 1 to 3 considered as viable due to least RMS values in this combination [27]. Santos et al. (2014) also standardized the preparation and staining procedure and seed viability classes in *Pinus taeda* seeds using tetrazolium test [33]. Thus, topographical staining categories 1, 2 and 3 provided least root mean square value and recommended these as viable seed categories for TZ seed viability testing in *Pinus roxburghii*.

**Table 3. RMS (Root Mean Square) determination for depiction of viable seed categories (1-5) in *Pinus roxburghii* seeds after tetrazolium staining**

Sr. No.	Staining categories considered as viable	Root mean square
1	1,2,3,4,5	27.41
2	1,2,3,4	23.49
3	1,2,3,5	23.60
4	1,2,4,5	24.00
5	1,3,4,5	24.16
6	1,2,3	18.89

Sr. No.	Staining categories considered as viable	Root mean square
7	1,2,4	19.39
8	1,2,5	19.52
9	1,3,4	19.60
10	1,3,5	19.72
11	1,4,5	20.20
12	1,2	13.47
13	1,3	13.76
14	1,4	14.43
15	1,5	14.60

When the seed viability percentage of first three categories (1,2 & 3) was pooled together, the potential seed viability was comparable to the actual laboratory germination in all five seed lots (Table 4). The relationship between potential seed viability and actual viability for all the five seed lots showed that potential viability was positively comparable with laboratory germination for *Pinus roxburghii* seeds. This is also confirmed the standardised protocol for quick seed viability test in *Pinus roxburghii* in the stipulated time period. Aslam et al. (2010) reported that the seed viability percentage in tetrazolium test (79.82 %) did not vary significantly than the germination percentage (82.69 %) in seed germination test in *Pinus wallichiana* [8]. Gagliardi et al. (2011) advocated that the tetrazolium test (TZ) proved to be a definite test for seed viability than germination viability test which takes longer time in days to complete it, hence tetrazolium test is efficient for evaluating the physiological potential seed viability in short time of period [34]. Thus, the standardized protocol was confirmed with the laboratory seed germination test in *Pinus roxburghii*.

**Table 4. Relationship between laboratory germination and potential viability as determined by TZ test for five seed lots of *Pinus roxburghii* seeds**

Sr. No.	Seed lots	Laboratory germination (%)	Potential viability (%)
1	A	70.25	67.25
2	B	73.50	69.75
3	C	76.25	72.00
4	D	72.25	70.25
5	E	74.50	75.75
<b>Mean</b>		<b>73.35</b>	<b>71.00</b>

When seed vigor test was compared with germination test both in the laboratory and in nursery beds for loblolly and slash pines then the correlation was found between tetrazolium staining and leachate conductivity with seed germination [35]. The basic objective of seed vigor or quality testing is to provide a precise or accurate physiological potential among different seed lots before seed sowing as similar to germination percentage mostly to identify good seeds lots [36] for higher germinability as well as seed storage. The results of the present study fulfilled all the gaps and criteria of quick seed viability testing in *Pinus roxburghii* and the standardized protocol could be used for future seed vigour and quality testing within a stipulated time period.

#### 4. CONCLUSION

From the present experiment, it can be concluded that the seeds of *Pinus roxburghii* should be soaked in 0.5 % tetrazolium solution at

25°C temperature for 24 hours to test the viability and quality of seeds before sowing in nursery with the counting of seed staining categories 1, 2 and 3 within a short period of time for the forest restoration and conservation purposes. There is a need to test this standardized seed viability protocol for different seed storage periods for future research thrust.

#### ACKNOWLEDGEMENTS

The present experiment is formulated by RKD and carried out by Priya Thakur at Tree Seed Technology Laboratory, Department of Tree Improvement and Genetic Resources, College of Horticulture and Forestry, Neri, Hamirpur which is a continent College of Dr. Yashvant Singh Parmar University of Horticulture and Forestry, Naini, Solan, Himachal Pradesh, India. All the authors helped to prepare manuscript in the publication format. The authors are thankful to authorities of the university for providing



necessary facilities and funding during the course of present study.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Roy SM, Thapliyal RC, Phartyal SS. Seed source variation in cone, seed and seedling characteristic across the natural distribution of Himalayan low level pine *Pinus roxburghii* Sarg. *Silvae Genetica*. 2004;53(1-6):116-23.
2. Ghildiyal SK, Sharma CM, Gairola S. Additive genetic variation in seedling growth and biomass of fourteen *Pinus roxburghii* provenances from Garhwal Himalaya. *Indian Journal of Science and Technology*. 2009;2(1):37-45.
3. Troup RS. The silviculture of Indian trees. Clarendon Press; 1921.
4. Khanduri VP, Sharma CM. Wind pollination in *Pinus roxburghii*. *Progress in Natural Science*. 2007;17(1):32-8.
5. Ghildiyal SK, Sharma CM, Khanduri VP. Effect of pre-soaking and pre-chilling treatments on seed germination of *Pinus roxburghii* provenances from western Himalaya, India. *Journal of Forestry Research*. 2009;20:323-30.
6. Moore P. TZ checks your seed quality. *Crops and Soils*. 1962;15:10-12.
7. Baskin CC, Baskin JM. Seeds: ecology, biogeography, and, evolution of dormancy and germination. Elsevier; 1998.
8. Aslam M, Reshi ZA, Siddiqi TO. Standardization of seed viability protocol for *Pinus wallichiana* AB Jackson in Kashmir, India. *Int. J. Pharm. Sci. Rev. Res*. 2010;4(3):93-8.
9. Baalbaki R, Elias S, Marcos Filho J, McDonald MB. Seed vigor testing handbook. Ithaca: Association of Official Seed Analysts. 2009. 341p. (AOSA. Contribution, 32).
10. Lakon G. Topographical detection of viability of cereal seeds with tetrazolium salts. *Ber. Deut. Bot. Ges*. 1942;60:299-305.
11. Lakon G. The topographical tetrazolium method for determining the germinating capacity of seeds. *Plant physiology*. 1949;24(3):389-94.
12. Wood CB, Miles S, Rix C, Terry J, Daws MI. The effect of seed oil content on viability assessment using tetrazolium: a case study using 171 species. *Plant Genetic Resources Newsletter*. 2005; 143:17.
13. ISTA. International Seed Testing Association. International rules for seed testing. Rules-1999. 1999.
14. On D. 2, 3, 5-triphenyl tetrazolium chloride as a viability indicator of certain coniferous seeds.
15. García Quintana Y, Arteaga Crespo Y, De Decker M, Castelo MD. Seed viability, germination and seedling quality patterns of three forest species for restoration in Amazonian conditions. *Enfoque UTE*. 2023;14(1):18-32.
16. Bertiller MN, Aloia DA. Seed bank strategies in Patagonian semi-arid grasslands in relation to their management and conservation. *Biodiversity & Conservation*. 1997;6:639-50.
17. Xiang X, Zhang Z, Wu G. Effects of seed storage conditions on seed water uptake, germination and vigour in *Pinus dabeshanensis*, an endangered pine endemic to China. *Seed Science and Technology*. 2019;47(2):229-35.
18. Viswanath S, Singh RP, Thapliyal RC. Seed germination patterns in a Himalayan moist temperate forest. *Tropical Ecology*. 2002;43(2):265-73.
19. Dhaka RK, Jha SK. Evaluation of five teak (*Tectona grandis* LF) provenances for germination test to find out reasons of low germination. *International Journal of Pure and Applied Bioscience*. 2017;5(5):1420-6.
20. Gaspar S, Nagy J. A new method for tetrazolium analysis of wheat seeds without embryo excision. *Seed Science and Technology*. 1981;9:553-56.
21. Chhabra R, Singh T. Seed aging, storage and deterioration: An irresistible physiological phenomenon. *Agricultural Reviews*. 2019;40(3):234-8.
22. Savonen EM. An improvement to the topographic tetrazolium testing of Scots pine (*Pinus sylvestris* L.) seeds. *Seed Science and Technology*. 1999;27:49-57.
23. Donald DG, Jacobs CB. The effect of storage time, temperature and container on the viability of the seed of four pine species. *South African Forestry Journal*. 1990;154(1):41-6.
24. Daws MI, Cousins C, Hall J, Wood CB. Pressure–time dependency of vacuum

- degassing as a rapid method for viability assessment using tetrazolium chloride: a comparative study of 17 *Pinus* species. *Seed Science and Technology*. 2006;34(2): 475-83.
25. Wortmann S, Kutta RJ, Nuernberger P. Monitoring the photochemistry of a formazan over 15 orders of magnitude in time. *Frontiers in Chemistry*. 2022;10: 983342.
  26. Willan RL. A guide to forest seed handling with special reference to tropics. Danida Forest Seed Center, DK-3050, Humlebaek, Denmark, 1985.
  27. Phartyal SS, Thapliyal RC, Nayal JS, Joshi G. Assessment of viability of *Acer caesium* and *Ulmus wallichiana* seeds through the tetrazolium test. *Seed science and technology*. 2003;31(3):691-700.
  28. Sheoran OP, Tonk DS, Kaushik LS, Hasija RC, Pannu RS. Statistical software package for agricultural research workers. Recent advances in information theory, statistics & computer applications by DS Hooda & RC Hasija Department of Mathematics Statistics, CCS HAU, Hisar. 1998;8(12):139-43.
  29. Kuo WHJ, Yan AC, Leist N. Tetrazolium tests for the seeds of *Salvia splendens* and *S. farinavea*. *Seed Science and Technology*. 1996;24:17-21.
  30. Jerlin R, Vanangamudi K, Rajasekaran R. Standardising methods of quick viability test for Pungam (*Pongamia pinnata* L. Pierre) seeds. *Tropical Agricultural Research and Extension*. 2007;7:261-63.
  31. Pallavi HM, Madalageri MB, Vishwanath K, Biradar IB, Thattimani M. Decortication to enhance seed value and standardization of quick viability test in drumstick (*Moringa oleifera* L.). *Environment & Ecology*. 2015;33(4):1558-61.
  32. Masullo LS, Piña-Rodrigues FC, Figliolia MB, Américo C. Optimization of tetrazolium tests to assess the quality of *Platymiscium floribundum*, *Lonchocarpus muehlbergianus* and *Acacia polyphylla* DC. seeds. *Journal of Seed Science*. 2017;39:189-97.
  33. Santos FS, Vieira ES, Panobianco M. Tetrazolium test for *Pinus taeda*: preparation, staining, and seed viability classes. *Pesquisa Agropecuária Brasileira*. 2019;54.
  34. Gagliardi B, Marcos Filho J. Assessment of the physiological potential of bell pepper seeds and relationship with seedling emergence. *Revista Brasileira de Sementes*. 2011;33:162-70.
  35. Bonner FT. Measurement of seed vigor for loblolly and slash pines. *Forest Science*. 1986;32(1):170-8.
  36. Marcos Filho J. Seed vigor testing: an overview of the past, present and future perspective. *Scientia agricola*. 2015;72: 363-74.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/114321>