



Efficacy of Methyl Eugenol and Cuelure Traps for Monitoring of Cucurbit Fruit Fly (*Bactrocera spp.*) in different Doses in Pumpkin

Puneet Kumar ^{a*}, Umesh Chandra ^b, Rishabh Mishra ^c,
Ashutosh Singh Aman ^c and Sova Yadav ^a

^a Department of Entomology, Banaras Hindu University Varanasi, U.P., India.

^b Department of Entomology, Acharya Narendra Deva University of Agriculture and Technology, Ayodhya, U.P., India.

^c Department of Entomology, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, U.P., India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARJA/2024/v17i2431

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/115131>

Original Research Article

Received: 24/01/2024

Accepted: 27/03/2024

Published: 30/03/2024

ABSTRACT

The present study on the "Efficacy of methyl eugenol and cue lure traps for monitoring of cucurbit fruit fly (*Bactrocera spp.*) in different doses in pumpkin" was conducted in the Department of Entomology, Acharya Narendra Deva University of Agriculture & Technology, Kumarganj, Ayodhya, during 2020-21. Among the 3 different doses tested, methyl eugenol rakshak trap with 1.5ml +2 ml (methyl eugenol + malathion 50 % E.C. were significantly superior in attracting highest number of fruit flies *B. zonata* and *B. dorsalis* with trap catches of 69.39 fruit flies /trap/week followed by 2ml

*Corresponding author: E-mail: puneet22025@bhu.ac.in;

+2ml doses (49.36 fruit flies/trap/week). 1ml + 2ml (methyl eugenol +malathion 50% E.C. performance was significantly lower to other doses (42.47 fruitflies/traps/week) and other three doses tested, cue lure rakshak trap with 2 ml + 2ml (cue lure + malathion 50% E.C) were significantly superior in attracting highest number of fruit flies *B. cucurbitea* and (19.92 fruit flies /trap/week) followed by 1.5 ml +2ml doses (13.69 fruit flies/trap/week). 1ml + 2ml (cue lure +malathion 50% E.C. 50 % E.C performance was significantly lower to other doses (7.42 fruitflies/traps/week).

Keywords: Methyl eugenol; cue lure; malathion; fruitflies; rakshak trap.

1. INTRODUCTION

Pumpkin (*Cucurbita moschata*) is one of the main vegetable crop cultivated since prehistoric times and is now the most common cucurbit in Asia and the United States of America. *C. moschata* is grown in approximately in all the regions of India [1]. The fresh pumpkin fruit contains 92.2 % moisture, 0.15 % fat, 0.98 % protien, 0.76% ash, 0.56 % crude fiber and 5.3% corbohydrates [2] Pumpkin fruit is rich in carotenoid, vitamins, minerals, and dietary fibers. Dhiman *et al* [3]. The pumpkin fruit contains 78-86 % edible portions. The seed of pumpkins contains 3.1% of total fruit weight, are abundant in protein 33%, high in S-containing amino acids, and low in phytic acids and trypsin inhibitors. Two compounds, methyl eugenol (4-allyl-1, 2-dimethoxybenzene) and cue-lure [4-(pacetoxyphehyl)- 2-butane one, play significant roles as attractants for male tephritid fruit flies Metcalf *et al*. [4]. These pheromones are plant-related products derived from phenylpropanoid and related compounds. Methyl eugenol occurs mostly as a natural product in the plant kingdom, is a highly dominant attractant for the males of *B. dorsalis*. Due to its strong attractant properties, it has been used as a pheromone that captures male fruit flies for monitoring and management. Outbreak of fruit fly is happening regularly in Eastern Uttar Pradesh, India. For monitoring of insect pests in crop, various techniques are used but pheromone traps are most effective. Methyl eugenol and cue-lure were used by Khoo and Tan [5] for monitoring the fruit flies. Hardy et. al. [6] also reported that at least 90 percent of the Dacinae species were strongly attracted to either methyl eugenol or to cue-lure-raspberry ketone. All these findings relate to foreign studies, and very meager work is done in India on semiochemicals-based management of fruitflies in pumpkin crops. These pheromones are very expensive, so it is important to know how much dose is most effective in one trap; so, this study was designed for field evaluation of traps, methyl eugenol, and cue-lure with the objective of

finding the most efficient dose of the two attractants for the attract-and-kill strategy of the pest.

2. MATERIALS AND METHODS

This experiment was conducted in RBD with 7 treatment and 3 replications at Students' Instruction Farm, Acharya Narendra Deva University of Agriculture & Technology, Kumarganj, Ayodhya during February to June 2020. A distance of 2 m was maintained between each treatment. Mainly Rakshak fruit fly traps were used during the study. Commercial grade methyl eugenol, cuelure, malathion 50 EC were procured. These pheromones and insecticide were impregnated in cotton wicks with the help of a 5ml syringe. In each traps cotton wicks (size: 3cm x 1cm) impregnated with different amounts of methyl eugenol and cuelure, viz., 2 ml, 1.5 ml and 1 ml along with and 2ml malathion were put. These traps were installed in the field at the time of crop germination by hanging with the help of string on the branches of trees. The old pheromone traps were replaced with fresh traps every month. Data on attracted and trapped male pumpkin fruit fly were collected on daily basis. The trapped male pumpkin fruit flies were identified on the basis of morphological character.

3. RESULTS AND DISCUSSION

The data of efficacy of methyl eugenol and cue lure in different doses has been given in Table -2

3.1 Response of *B. dorsalis* + *B. zonata* in different Dose of Methyl Eugenol Rakshak Trap

During the first Month of test period fruitfly captures varied significantly among different dose of methyl eugenol rakshak traps with greater catches in 1.5ml (methyl eugenol) + 2ml Malathion 50% EC Rakshak traps (68.00

fruitfly/trap/week) it was significant difference with 2ml (methyl eugenol) + 2ml Malathion 50% EC Rakshak traps (49 fruitflies/trap/week) whereas, the lowest catches was recorded in 1ml (methyl eugenol) +2ml Malathion 50% EC Rakshak traps (41 fruitflies /trap/week). The same trend was observed at the second month of test period fruitfly captures varied significantly among different dose of methyl eugenol rakshak traps with greater catches in 1.5 ml (methyl eugenol) +2 ml Malathion 50% EC Rakshak traps (59.25 fruitfly/trap/week) it was significant difference with 2 ml (methyl eugenol) +2 ml Malathion 50% EC Rakshak traps (45.50 fruitflies/trap/week) whereas, the lowest catches was recorded in 1 ml (methyl eugenol) +2 ml Malathion 50% EC Rakshak traps (37.33 fruitflies /trap/week). At third month of test period, 1.5ml (methyl eugenol) + 2ml Malathion 50% EC Rakshak traps attracted significantly more fruit flies with highest trap catches of 80.0 fruitfly/trap/week. 1.5 ml dose was superior than 2 ml dose of methyl eugenol Rakshak traps. 2ml (methyl eugenol) + 2ml Malathion 50% EC Rakshak traps was attracted 53.00 fruitflies/trap/week whereas, the lowest catches were recorded in 1 ml (methyl eugenol) + 2ml Malathion 50% EC Rakshak traps (48.92 fruitflies /trap/week). The similar research was done by Shankar Mandal *et al.* [7] that is Ten replicate analyses were done for both methyl eugenol quantity of the pheromones in formulated products was found to be 0.55 ± 0.11 g/lure (Mean \pm SD), respectively as well as Khan *et al.* [8] observed that methyl eugenol catches (81.69 fruitfly/trap/week).

3.2 Response of *B. cucurbitae* + *B. tau* in different Dose of Cue Lure Rakshak Traps

During the first Month of test period fruitfly captures varied significantly among different dose of cue lure rakshak traps with greater catches in 2 ml (cue lure) + 2ml Malathion 50% EC Rakshak traps (6.50 fruitfly/trap/week) it was at par with 1.5ml (cue lure) +2ml Malathion 50% EC Rakshak traps (4.67 fruitflies/trap/week) whereas, the lowest catches was recorded in 1ml (cue lure) + 2ml Malathion 50% EC rakshak traps (3.42 fruit flies /trap/week). During the second Month of test period fruitfly captures varied significantly among different dose of cue lure rakshak traps with greater catches in 2 ml (cue lure) +2ml Malathion 50% EC Rakshak traps (21.42 fruitfly/trap/week) which was significant difference with 1.5ml (cue lure) + 2ml Malathion

50% EC Rakshak traps (15.33 fruitflies/trap/week) whereas, the lowest catches was recorded in 1ml (cue lure) + 2ml Malathion 50% EC Rakshak traps (8.0 fruitflies /trap/week). At third month of test period, 2 ml (cue lure) +2ml Malathion 50% EC Rakshak traps attracted significantly more fruit flies with highest trap catches of 31.83 fruitfly/trap/week. 2 ml dose was superior than 1.5 ml dose of cue lure Rakshak traps. 1.5 ml (cue lure) + 2ml Malathion 50% EC Rakshak traps was attracted 21.08 fruitflies/trap/week whereas, the lowest catches was recorded in 1ml (cue lure) + 2ml Malathion 50% EC Rakshak traps (10.83 fruitflies /trap/week). Khan *et al.* [8] observed that cue lure was more effective, resulting in higher male catches of fruit flies (171.82/trap/week), compared to methyl eugenol (81.69/trap/week). They also identified ecological conditions as the reason for population fluctuation, with maximum temperature accounting for 33.80% of the variation, while average temperature and wind speed contributed 12.70% collectively. Due to these reasons, as well as another factor, which was the onset of fruiting in fruit plants near the experimental plot during the time of experimentation, fluctuations occurred in the number of fruit fly catches in both methyl eugenol and cue lure traps.

3.3 Effect of Pumpkin Fruits Yield in Various Dose of Methyl Eugenol and Cue Lure

The data are presented in Table 3 revealed that the all treatment produce significantly higher yield than control (144q/ha). The Treatment 6 (T6) where cue lure 2ml + malathion 2ml was trapped produced maximum yield (236 q/ha) it was significant difference with T5 where cue lure 1.5 ml +malathion 2ml was trapped produced yield (223 q/ha). Followed by T2 where was methyl eugenol 1.5 ml + malathion 2 ml was trapped produced yield (217q/ha) it was at par with T3 where methyl eugenol 2 ml +malathion 2ml was trapped produced yield (212 q/ha). Followed by T4 where was cue lure 1 ml + malathion 2 ml was trapped produced yield (209 q/ha) it was significant difference with T1 where methyl eugenol 1 ml +malathion 2ml was trapped produced yield (196 q/ha). According to Raghuvanshi *et al.* [9] The minimum yield produced 144 q/ha. In *Kharif*, maximum bitter gourd fruit damage (62.70%) occurred in the 45 SW. A second peak, with 49.70% fruit damage was observed during the 15 SW period.

Table 1. Details of treatments for the efficacy of methyl eugenol and cuelure

Treatment	Chemical mixture	Dose
T1	Methyl Eugenol +Malathion50EC	1ml+2ml
T2	Methyl Eugenol +Malathion50EC	1.5ml+2ml
T3	Methyle Eugenol + Malathion 50EC	2ml+2ml
T4	Cuelure+Malathion50EC	1ml+2ml
T5	Cue lure + Malathion 50EC	1.5ml+2ml
T6	Cue lure + Malathion 50EC	2ml+2ml
T7	Untreated Control	

Table 2. Efficacy of methyl eugenol and cue lure in different doses

Treatments	Chemicals	Dose	Fruitflies/trap/week			
			First month	Second month	Third month	Mean
T1	Methyl Eugenol+Malathion 50% EC	1ml+2ml	41.17 (6.45)	37.33 (6.15)	48.92 (7.03)	42.47
T2	Methyl Eugenol+Malathion 50% EC	1.5ml+2ml	68.00 (8.27)	59.25 (7.73)	80.92 (9.02)	69.39
T3	Methyl Eugenol+Malathion 50% EC	2ml+2ml	49.58 (7.07)	45.50 (6.78)	53.00 (7.30)	49.36
T4	Cue lure+ malathion 50% EC	1ml+2ml	3.42 (1.98)	8.00 (2.98)	10.83 (3.37)	7.42
T5	Cue lure+ malathion 50% EC	1.5ml+2ml	4.67 (2.27)	15.33 (3.98)	21.08 (4.64)	13.69
T6	Cue lure+ malathion 50% EC	2ml+2ml	6.50 (2.65)	21.42 (4.68)	31.83 (5.69)	19.92
T7	Untreated Control		5.92 (2.53)	5.92 (2.53)	6.00 (2.53)	5.94
	SEm±		0.12	0.08	0.17	0.37
	C.D. at 5%		0.39	0.27	0.54	1.17

Table 3. Variation of yield in different doses

No. of Treatments	Treatments	dose/trap	Concentration	Damage percent	yield/hect
T1	Methyl Eugenol+Malathion 50% EC	1ml+2ml	12 trap /hec	34.67	196
T2	Methyl Eugenol+Malathion 50% EC	1.5 ml+2ml	12 trap /hec	27.67	217
T3	Methyl Eugenol+Malathion 50% EC	2 ml+ 2ml	12 trap /hec	29.33	212
T4	Cue lure+ malathion 50% EC	1ml+2ml	12 trap /hec	30.33	209
T5	Cue lure+ malathion 50% EC	1.5 ml+2ml	12 trap /hec	25.67	223
T6	Cue lure+ malathion 50% EC	2 ml+ 2ml	12 trap /hec	21.33	236
T7	Untreated Control			52.00	144
	SEm±			1.10	3.37
	CD at 5 %			3.44	10.52




Fig. 1. Efficacy of Methyl eugenol and cuelure in different doses

4. CONCLUSIONS

The present research on utilizing the Rakshak trap to attract fruit flies, it was observed that a higher dose of methyl eugenol proved to be less effective compared to a lower dose. Furthermore, in the case of cue lure, a lower dose did not attract more fruit flies but exhibited similar efficacy to a higher dose. Methyl eugenol+ malathion (1.5 ml +2ml) standard dose caught 69.39 fruitflies (*Bactrocera dorsalis* + *Bactrocera zonata*). whereas cuelure+ malathion (2ml+2ml) standard dose caught 19.92 fruitflies (*Bactrocera cucurbitae*+ *Bactrocera tau*). According to Divya, S. et al. [10] the mean of overall captured of *B. cucurbitae* in Jar trap + Cue lure + ME (DISC) combination were significantly greater than all other treatments (208.5 numbers) and less in treatments Jar trap + AA (Disc) combination (5.75 numbers) respectively fruit fly. Hence, it is imperative for farmers to be cognizant of the recommended dosage of any chemical or insecticide before application. Excessive amounts not only yield inferior results but also pose environmental hazards.

ACKNOWLEDGEMENT

The author acknowledges the Department of Agricultural Entomology, Acharya Narendra Dev

University of Agriculture and Technology, Kumarganj Ayodhya, India for providing the laboratory facilities and instruments to carried out the research programme successfully.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Nath P, Bhushan S. Screening of cucurbit crops against fruit fly. Annals of Plant Protection Science. 2006;14:472-473
- See EF, Wan NWA, Noor AAA. Physico-chemical and sensory evaluation of breads supplemented with pumpkin flour. Asian Food J. 2007;14:123-130
- Dhiman AKK, Bavita SA, Ramachandran P. Preparation of pumpkin powder and pumpkin seed kernel powder for supplementation in weaning mix and cookies. Int. J. Chem. Stud. 2018;6: 167-175.
- Metcalf RL. Chemical ecology of Dacinae fruit flies (Diptera: Tephritidae). Entomological Society of America. 1990;83:1017-1030.

5. Khoo CCH, Tan KH. Attraction of both sexes of melon fly, *Bactrocera cucurbitae* to conspecific males- a comparison after pharmacophagy of cue-lure and a new attractant Zingerone. *Ent. Experim. Appl.* 2000;97:317-320.
6. Hardy DE. Contribution of taxonomic studies to Integ. Pest management. Fruit flies with emphasis on the Asia pacific region. In:Proc. First Intern. Symp. 1991;430.
7. Mandal S, Rahman AMH, Mamun MIR, Shoeb M. Method development and validation for estimation of commercially produced sex pheromones in lure, *Journal of Bangladesh Chemical Society.* 2012;25(2):180-185.
8. Khan MA, Gogi DA, Ali A. Efficacy of methyl eugenol and cue-lure traps for monitoring melon fruit fly in relation to environmental conditions in bitter gourd, *J. Agric. Res.* 2010;48(4).
9. Raghuvanshi AK, Satpathy S, Mishra DS. Role of abiotic factors on seasonal abundance and Infestation of fruit fly, *Bactrocera cucurbitae* (COQ.) on bitter gourd. *Journal of Plant Protection Research.* 2012;2-52.
10. Divya A, Sukumaran S. An Efficient Vector Quantization Based Image Compression using Fruit Fly Algorithm, *Digital signal processing.* 2019;12(1).

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/115131>