



Comprehensive Review on the Bio-ecology and Integrated Management Strategies for Brinjal Shoot and Fruit Borer (*Leucinodes orbonalis* Guenee)

Sakshi Sharma ^{a*}, Ankush Mohapatra ^a
and Khushboo Kathayat ^a

^a Department of Horticulture, School of Agriculture, Lovely Professional University, Phagwara, Punjab- 144411, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i71029>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119027>

Review Article

Received: 15/04/2024

Accepted: 19/06/2024

Published: 20/06/2024

ABSTRACT

The most important and commonly grown vegetable for both raw and cooked purposes is brinjal, or *Solanum melongena* Linnaeus. It is a member of the solanaceae family and is also known as eggplant or baingan. Nevertheless, it faces significant threat from a prominent pest known as the eggplant shoot and fruit borer, scientifically termed *Leucinodes orbonalis* Guenee, capable of inflicting damage ranging from 37% to 100%. This pest can also diminish both the quantity and

*Corresponding author: E-mail: sakshisharma5713@gmail.com;

Cite as: Sharma, Sakshi, Ankush Mohapatra, and Khushboo Kathayat. 2024. "Comprehensive Review on the Bio-Ecology and Integrated Management Strategies for Brinjal Shoot and Fruit Borer (*Leucinodes Orbonalis* Guenee)". *Journal of Advances in Biology & Biotechnology* 27 (7):692-708. <https://doi.org/10.9734/jabb/2024/v27i71029>.

quality of eggplant produced. Farmers persist in depending on pesticides to address this problem; nevertheless, excessive pesticide application has resulted in negative impacts on the environment, unintended beneficial organisms, phytotoxicity, pesticide resistance, pest resurgence, bioaccumulation, and secondary pest outbreaks. In different regions of the world, it has been discovered that a number of insects, including Various pests such as the Fruit and Shoot Borer, White Fly, Leaf Hopper, Thrips, Mites, Leaf Roller, and Red Spider Mite contribute to losses in eggplant. Moreover, this insect can also cause severe harm to other vegetables within the Solanaceae family, acting as an alternative host. The adult insect can eventually withstand the problems of chemical pesticides and find it challenging to control the insect population in standing crops due to the larva's unique ability to subsist on a monophagous diet supported by homing and tunneling behavior. It results in a decrease in both yield and vitamin C content. This is due to the fact that high humidity and moderate temperatures encourage the population growth of the Brinjal Fruit and Shoot Borer, which results in significant losses in hot, humid weather. Farmers primarily use chemical insecticides, which they apply carelessly to manage this pest. A lot of farmers also employ biological control techniques and home-based remedies like marigold barriers, cow urine, ashes, and so forth. Farmers are unable to totally manage the infestation, though, and the measures cost more to produce than they really bring.

Keywords: *Brinjal; shoot and fruit borer; integrated pest management.*

1. INTRODUCTION

The most prominent vegetable crop and tender perennial plant cultivated for its tasty fruit is the eggplant (*S. melongena* L.), belonging to the Solanaceae/Nightshade family and the Solanoideae subfamily. It is referred to as brinjal in South Africa and Southeast Asia, eggplant in the United States, Australia, New Zealand, and Canada, aubergine or guinea squash in the United Kingdom, Ireland, and Quebec, and garden egg in Quebec [7]. Because of its widespread use and adaptability, brinjal, also known as baingan, is referred to be the "King of vegetables" and is utilized in Indian cuisine on both regular and festive occasions. Brinjal fruits are widely utilized in various culinary dishes, including sliced bhaji, packed curry, bertha, chutney, vangnibath, and pickles. Commercially produced brinjal fruit comes in a variety of shapes, sizes, and colors, including round, rectangular, pendulum, egg shaped, green, white, and yellow, as well as striated tones [36]. Its skin exhibits a smooth and shiny texture. Brinjal stands out as a leading vegetable in terms of oxygen radical absorption capacity and serves as a significant source of vitamins, minerals, proteins, cancer- preventive agents, dietary fiber, and factors conducive to weight training [54]. Nutritionally speaking, 100 grams of cooked fruit has a very low-calorie value of 25.0%, 92.7% moisture, 8.29 grams of carbohydrates (of which 3.04 are sugar), 0.2 grams of fat, 1 gram of protein [8], 21.1 µg of beta-carotene [10], and 3.4 grams of fiber. Other elements include 213.0 mg of potassium, 10.6 mg of magnesium, 13.0 mg of

sodium, and 0.7 mg of iron [11-13]. The ripened fruit, per 100g, also contains 12.0 mg of calcium, 26.0 mg of phosphorus, 8.93 mg of choline, 13.4g of folate, 5.0 mg of ascorbic acid, and 27 International Units of vitamin A. Additionally, it contains 0.89 mg of vitamin B, 2.2 mg of vitamin C, 0.30 mg of vitamin E, and 3.5 µg of vitamin K [109,116,117,118,119].

The peel of brinjal types with rich blue or purple colors has a substantial number of anthocyanins, which are phenolic flavonoid phytochemicals that help prevent neurological illnesses, aging, and cancer [73,106]. Apart from serving as a popular appetizer, aphrodisiac, cardiac tonic, laxative, and anti-inflammatory, brinjal has also been cited in Ayurveda as a remedy for treating diabetes [62]. It is also a great therapy for people with liver issues. While often associated with Middle Eastern or Mediterranean cuisines, brinjal has been cultivated in the region for the past 4,000 years. A warm weather crop, brinjal is grown in subtropical areas worldwide. Nonetheless, it is extensively grown throughout the world in tropical and temperate climates, mostly during the warm season [82]. Although less renowned, the Gboma eggplant (*S. macrocarpon* L.) and the scarlet eggplant (*S. aethiopicum* L.), two other cultivated eggplant species, hold significant importance locally in Sub-Saharan Africa [20]. Brinjal is grown in outdoor fields, polyhouses, net houses, kitchen gardens, and commercial gardens across the globe during the Rabi and Kharif seasons. It ranks as the fifth most important solanaceous crop economically, trailing behind tobacco, tomato, potato, and

pepper. According to Gautam [30], eggplant is among the top five vegetable crops cultivated in Asia and the Mediterranean.

Additionally, frozen or fresh brinjal is exported. With a productivity of 29 tons per hectare and a production of 54077210 tons, eggplant is grown over 1864556 hectares worldwide. Region wise, Asia accounts for the largest portion of eggplant production (93.6%), with Africa coming in second (3.8%), Europe in second place (1.8%), America in third place (0.7%), and Oceania placing last (0%). In terms of both area and global brinjal production, India is ranked second only to China. With a productivity of 17.43 tons/hectare, eggplant is grown on 736000 hectares of land in India, where it is produced in 12826000 tons. From the nursery stage to harvesting, a number of insect pests and mites attack eggplants. These pests include *Thrips palmi* (Karny), *Eublemma olivacea* (Walker), *Leucinodes orbonalis* (Guenee), *Bemisia tabaci* (Gennadius), *Amsasca biguttula biguttula* (Ishida), *Henosepilachna vigintioctopunctata* (Fab.), *Amsasca biguttula biguttula biguttula* (Ishida), and *Tetranychus macfarlanei* (Baker and Pritchard) [105], and *Tetranychus macfarlanei* (Koch) [60]. *L. orbonalis* is the harmful pest found in Asia, [68,108] among them [50,17, and 91]. According to reports, this infamous pest limits the growth of brinjal in India, resulting in losses of 37–63% [25], up to 90% [39], as high as 70–92% [71], and up to 100% damage if management measures are not implemented [82]. Losses of up to 67% have been observed in Bangladesh [24], 31 to 90%, and 50–70% have been reported in Pakistan. Because of the borer's severe infestation and the reduced yields, many farmers are reluctant to cultivate brinjal [67]. Because moderate temperatures and high humidity encourage population growth and result in significant losses during hot and humid conditions, the losses in agricultural yield decrease caused by pests vary from season to season and from place to location [95,14,33]. Unpredictable weather, such as sudden drops in temperature, droughts, or floods, can also lower fruit quality and production [107,64] and [110,111]. Presently, farmers apply numerous insecticides, sometimes up to 140 times or more during a cropping season, which typically spans 6–7 months and incurs costs amounting to 32% of all agricultural production [4].

In Bangladesh, a significantly high amount of pesticides, approximately 180 times the usual, were utilized in a single year to protect brinjal against the Brinjal Fruit and Shoot Borer (BFSB),

as reported in an insecticide survey (40). There exists an economic threshold level for shoot and fruit borer in brinjal, indicating that 0.5% shoot damage, 5% fruit damage, and 8–10 moths per day per trap are considered significant [23]. To combat BFSB, farmers in Bangladesh often resort to applying broad-spectrum insecticides two or three times a week, and occasionally even twice a day. Insecticide usage is common, yet farmers still lose between 30 and 60 percent of their crop production to BFSB. The results of using 25–30 insecticidal sprays by farmers to control this pest are not good enough. Consequently, a season may see the application of more than 100 sprays, leaving heavy residues on the fruit. Thirty-five to forty percent of the entire expense of cultivating brinjal is incurred in pesticide treatments. Such an insecticide-dependent approach raises issues for farmers' and consumers' health and the environment [21]. Furthermore, improper insecticide treatment is leading to the emergence of secondary pests, the devastation of natural enemies, and a pest rebound. The current focus is on developing alternative management methods for this insect in order to prevent these issues. The larval stage of this insect is the only dangerous phase, as it feeds inside the fruit and creates large exit holes for the pupae once it has completed development. This reduces the fruit's market value and makes it unsuitable for human eating [5].

Damage starts when the seedlings are planted and lasts until the fruit is harvested. During the early stages of the plant's life, larvae pierce the petioles and midribs of large leaves and early shoots. This caused the entrance pores to close with their frass and the shoot to begin feeding inside [15], which ultimately caused the shoot to droop and wither [1]. The larva pierces the fruit and flower buds through the calyx during the last stages of fruit formation. The fruits have one or more sizable circular exit holes. Fruits that are impacted become internally rotten and lose their market value [27].

2. BIOLOGY OF BRINJAL SHOOT AND FRUIT BORER

2.1 Egg

A single female may deposit anywhere between 5 and 242 eggs over her lifetime, according to studies by Alam [2] and [45]. Most of the time, eggs were placed individually, albeit occasionally in groups of two or four. The bottom surface of fragile leaves, plant twigs, blooms, or fruit calyces

was the favored location for females to deposit their eggs. Before hatching, the creamy white, oval shaped or slightly elongated eggs became orange with a noticeable black mark [35,102]. Alpuerto [6], [40], [57], [85], and [102] have all documented the pre- oviposition, oviposition, and post-oviposition periods, which are, respectively, 1.1 to 2.1 days, 1.4 to 4.0 days, and 1.0 to 2.0 days. In contrast, the incubation period was reported by Mallik [61], to be 3 to 4 days. The highest hatching rate (38.2%) was observed on the third day after oviposition, followed by significant rates on the fourth and fifth days, respectively [85].

2.2 Larva

According to Kavitha [41], [35], [70], [85], and [102], larvae progressed through five instars before reaching the pupal stage. The typical durations of the first, second, third, fourth, and fifth larval instars were found to be 1-2, 2-3, 2-3, 2-4, and 2-4 days, respectively. The newly hatched larva was small, creamy or dirty white in color, with three pairs of thoracic legs, five pairs of prolegs, and a distinct dark brown or light black head. Larvae in their second instar were similar to those in their first, except they were bigger and had a somewhat darker color. Compared to the previous instars, the third instar larvae were substantially longer and darker, with a unique pattern on the prothoracic shield and dark brown thoracic legs [31]. The fourth instar had a color that was somewhat pink. The fifth instar had three

distinct thoracic segments, five pairs of well-developed prolegs, and a cylindrical, pinkish-brown color. Still, six larval instars of the shoot and fruit borer were documented by Alam [2] and [92]. The typical larval phase was found to extend between 12.3 and 14.0 days, according to reports from [22] and [41]. The pupal time was observed by Pupa [16] and [57] to vary between 7 and 10 days. They saw that the pupae had eight hook- shaped, fine spines at the posterior end of the abdomen, a small anal end, and a broader cephalic lobe. The pupae were dark brown in color. According to Alam [3], [40], and [53], pupation occurred on glass jars, earth, muslin fabric, fruits, and occasionally on plant leaves. Saeed [85] report that the pupal duration was found to vary from 6 to 8 days. They didn't notice the adult emerging until the fifth day following pupation.

According to Kavitha [41], [35], [70], [86], and [102], larvae progressed through five instars before reaching the pupal stage. The typical lengths of the first, second, third, fourth, and fifth larval instars were determined to be 1-2, 2-3, 2-3, 2-4, and 2-4 days, sequential-ly. On the sixth day following pupation, the adult began to emerge, and it did so until the eighth day. Maximum adult emergence was noted on the seventh day following pupation, with an average of 14%, 30%, and 10% emerging on the sixth, seventh, and eighth-days following pupation. The average adult emergence rate was shown to be 54% [37-38].




Fig. 1. Fruit borer and brinjal sprout life cycle

2.3 Adult

Male moths were found to have a lifespan of one to three days, while female moths typically lived for two to five days, as indicated by studies conducted by Stommel [102], [40], and [2]. The moth had a blackish brown head and thorax and was white in color. The pinkish brown patterns on the white wings were larger on the forewings. As per Kavitha [41], the females had a rounded posterior end and a larger abdomen with greater wing spread than the males, who were smaller in size and had a narrower abdomen that tapered posteriorly. Singla [70] report that the partial sex ratio was determined to be 1.0:2.0 in favor of females and 1.0:1.3 in favor of males, respectively. According to Saeed [85], mature *L. orbonalis* insects typically mate at night or in the early morning. Pre-mating hours ranged from 6 to 9 (with an average of 7.1 hours). The adults spent between thirty and forty-nine minutes (avg. 41.2 minutes) in the mating posture. The post-mating time averaged 5.0 days, ranging from 4-6 days. It was also mentioned by Muthukumar [57] that mating often occurs in the early morning and lasts for 43 minutes. According to records kept by Alam [2], the life cycle of the brinjal shoot and fruit borer takes between 19.0 and 43.0 days to complete (Fig. 1).

3. POPULATION DYNAMICS BRINJAL SHOOT AND FRUIT BORER

In Kanpur, during the fourth week of August, 47 days after transplanting, Sreenivasa [99] discovered the first infestation of shoot and fruit borer. As the temperature increased, the incidence steadily decreased from its peak, which occurred 114 days after transplanting, during the second week of September. It was shown that there was little or no correlation between the shoot damage and the meteorological conditions. In an experiment conducted at Manipur University, Sreenivasa [99] discovered that during the second week of April in 2003 and 2004, shoots became infected with fruit borer, resulting in damage percentages of 11.6% and 9.7%, respectively. The second week of June 2003 and the third week of May 2004 recorded the highest infection levels on shoots, with 25.8% and 31.4% of the shoots affected. The percentage of *L. orbonalis* infestation was favorably linked with both R.H. (80.5- 87.2%) and temperature (22.93-25.45°C). In the first year of the experiment, maximum relative humidity, rainfall, and wind speed showed positive correlations with brinjal shoot and fruit borer

infection. Conversely, in the second year, maximum relative humidity and sunshine hours exhibited positive correlations with the infection rates. The population incidence of the fruit borer, *L. orbonalis*, and brinjal shoot on *S. melongena* L c.v. Pusa Purple long was investigated by Srinivasan [100] during the two cropping seasons (2003 and 2004) in Manipur. They observed that the incidence of shoot and fruit borer began in April and persisted through the end of June. During the first and second cropping seasons, the pest on shoot peaked in the first week of June (29.45%) and the fourth week of May (25.24%), respectively. Conversely, the second week of June 2003 (67.16%) and the third week of June 2004 (72.25%) exhibited the highest prevalence of this insect on fruit [46].

Correlation studies revealed that average sunlight had a significant negative correlation with pest infestation on brinjal, while average temperature and relative humidity showed a significant positive correlation. In rabi 2009 in Durgapur, Meena [52] investigated the impact of abiotic conditions on the seasonal occurrence of the shoot and fruit borer, *L. orbonalis*. The findings indicated that shoot damage had a negative correlation with mean relative humidity (ranging from 21.8% to 75.3%) and a positive correlation with both maximum temperatures (ranging from 18.1°C to 37.88°C) and lowest temperatures (ranging from 4.6°C to 20.84°C), rainfall (ranging from 0 mm to 2.6 mm), and wind speed (ranging from 2.5 km/hr to 7.3 km/hr) [55]. On the contrary, the percentage of fruit infestation showed a negative correlation with mean relative humidity and a non-significant correlation with maximum and lowest temperatures, rainfall, and wind speed. According to Shaikat [56], the highest percentage of shoot infestation was noted during the ninth standard week (5.4%), followed by the seventh standard week (4.6%) and the eighth standard week (4.5%). Fruit borer was first observed in the tenth standard week and persisted until the final harvest. Fruit borer infestation peaked in the 18th and 17th standard weeks (43.3 and 40.1%, respectively). According to Raina [49] from Kanpur, throughout the vegetative phase of the crop up till the third week of September, there was a higher seasonal prevalence of the fruit borer, *L. orbonalis*, on the shoot [58]. The infection on shoots steadily decreased as the fruit grew, and by the end of October, when the crop was ripening, it had vanished as the borer infestation had shifted to the fruits during the second week of October. As winter arrived, it progressively became worse

and by the end of November, it was totally gone. Temperature, precipitation, and RH (morning) all had a very positive impact on the amount and intensity of infection on the shoots and fruits; however, RH (evening) had the opposite effect. On forty brinjal germplasm samples from Kalyanpur, Krattiger [51] investigated the seasonal occurrence of the fruit and shoot borer, *L. orbonalis*. The shoot borer infection first surfaced during the 43rd standard week (18–24 October).

The brinjal shoot borer exhibited positive multiplication rates at higher temperatures, while a negative correlation was observed between minimum temperature and relative humidity. There was no discernible relationship between wind speed and rainfall, although evaporation rate had a beneficial influence on the infesting shoot's ability to multiply. In their study on the population dynamics of brinjal shoot and fruit borer in Hisar during the summer of 2009–10, Kumar [44] discovered that the 39th and 40th standard weeks of the year had the highest number of larvae (10 larvae per 90 plants), while the 48th standard week had the lowest mean population (0.0 larvae per 90 plants). Larval population was shown to be inversely connected with percent and positively correlated with temperature, according to correlation analysis. In 2014, R.H. Saeed [85] from Hisar experimented on brinjal (var. BR-112) from June to October. They discovered that whereas fruit infestation first appeared in July, *L. orbonalis* infestation first appeared in shoots in June. The third week of September saw the highest prevalence of fruit borer and shoot borer. The third week of September reported the highest shoot damage (48.75%), fruit damage (40.00%) based on the number of fruits, and the greatest larval population (12 larvae per 20 plants). The maximum temperature was 35.3°C, the minimum was 25.0°C, and the relative humidity was 87% in the morning and 45% in the evening. After that, both the incidence of *L. orbonalis* in fruits and shoots began to decline. Additionally, correlation analysis indicated that there was no significant association between abiotic parameters and the mean larval population, fruit damage, and the percentage of shoot damage [59]. Regression study, however, revealed that abiotic variables account for 68% of population variance.

4. NATURE OF DAMAGE

The main food source for the almost monophagous brinjal shoot and fruit borer is

eggplant. Although the pest is thought to be hosted by *Solanum melongena*, several plants in the solanaceae family are frequently implicated in this regard. Major hosts are *S. Melongena* (L.) and *S. tuberosum* (L.), while minor and alternate hosts are *S. indicum* L. and *S. myriacanthum Dunal* [74], Sweet potato (*Ipomoea batatas* L.), Green pod of Austrian winterpea (*Pisum sativum* var. arvense L.) [12], Dark nightshade (*S. nigrum* L.), Turkey berry (*S. torvum* Swartz) [33], and Giloi (*S. gilo* Raddi). The wild hosts of *L. orbonalis* include Black nightshade (*S. anomalum* Thonn) [101], African eggplant (*S. macrocarpon* L.) [48], Tropical Soda Apple (*S. viarum* Dunal), Indian nightshade or Kantakari (*S. xanthocarpum* Schrad), Cape ooseberry (*Physalis peruviana* L.), Pygmy groundcherry (*Physalis minima* L.), and Forest Bitter Berry (*Solanum anguivi* Lam.) [28]. For brinjal, the most virulent internal feeder pest is the Shoot and Fruit Borer (FSB) [63]. By creating holes in the fruits and shoots, it not only reduces the production (number and quality) but also the fruit's vitamin C content by up to 80% and its aesthetic value. When the larvae first hatch, they promptly bore into the nearest tender shoot, petioles, developing bud, and flower. Subsequently, as the fruits develop, they penetrate into the fruit and consume its mesocarp, leading to the destruction of the fruit tissue [66]. The larvae created a dead heart when they bored into fruits, and they frequently filled in the feeding tunnel's opening with their excrement, called frass. Though fading entry hole depressions are evident, the fruit's entry holes are hidden because they have either healed or been covered with frass. Only the injured fruits display the big circular exit holes, one or more of which are visible [83].

Fruits that are impacted become deformed and internally decay, rendering them unsuitable for selling or eating, [84]. One fruit can have up to 20 larvae, according to research from Ghana. According to Krishnamoorthy [43] a single larva may ruin four to seven good fruits. The primary cause of damage to the plant is fruit feeding by the larvae, which bores into the tender shoots. Consequently, the affected twigs, flowers, and fruits undergo drying, withering, and sometimes premature falling off. This ultimately results in the wilting of young shoots and dieback of the branch terminals, thereby reducing the plant's ability to bear fruit. As a result, there are fewer and smaller fruits on the plants. Although new shoots can emerge, this postpones crop maturation and exposes the newly developed shoots to harm from larvae. Damaged blooms

that are fed on by larvae do not develop into fruit. Damages caused by Brinjal Shoot and Fruit Borer illustrated in Fig. 2.

5. HOST PLANT RESISTANCE

Numerous researchers have studied the screening of brinjal genotypes against the fruit borer, *L. orbonalis*, and shoot borer, *L. esculentus*, using host plant resistance mechanisms such as tolerance, antixenosis, and antibiosis. Insect resistance in brinjal plants is known to be correlated with several morphological and biochemical characteristics. Table 1 details the methods of host plant resistance to brinjal. In contrast to resistant

cultivars, susceptible kinds displayed greater levels of shoot infestation. A thin stem, numerous branches, the length and width of the lower third of the leaf, more spines, a rough leaf surface area, a thick cuticle heavily lignified, a broad and thick hypodermis, a closely packed vascular bundle, and a small pith area are characteristics of antixenosis that may indicate a lower infestation or, conversely, a higher infestation. Numerous researchers have examined the antixenosis mechanism of various plant characteristics [75-76]. Their findings have shown that the biophysical characteristics of shoot and fruit borer insect populations are reduced, as shown in Table 2.



Fig. 2. Nature of damages caused by brinjal shoot and fruit borer

Table 1. Characters with different resistance mechanisms in brinjal

Mechanism (s)	Character (s)
Antixenosis (non-preference)	Fruit colour, shape and diameter, size, Calyx size, pericarp thickness, surface wax, glandular and non-glandular trichomes, leaf size
Antibiosis	Total phenol, sugar content, polyphenol oxidase and peroxidase enzyme, solasodine contents, flavonols and potassium
Avoidance (escape)	Earliness with cold tolerance

Table 2. Antixenosis characters which shows the resistance/ reduction to brinjal shoot and fruit borer

S. No	Biophysical Varietal Characters	Reference
1.	Leaf trichomes, stem thickness and stem hair density	[44]
2.	Leaf thickness and trichome density	[65]
3.	Number of shoots per plant, spines of leaves, branches, petioles, calyx of fruits, fruit skin thickness, shoot thickness and longfruited varieties	[95]

Table 3. Antibiosis characters which shows the resistance/ reduction in brinjal to brinjal shoot and fruit borer population

S. No	Biochemical Characters	Reference
1.	Solanine content and total phenols	[9]; [81]; [42];[111]
2.	Polyphenol oxidase activity, total phenol content and solasodine content	[78]
3.	Phenolics content	[29]; [80]
4.	Polyphenol oxidase (PPO), Phenylalanine ammonium lyase (PAL) and Lignin	[48]

Pubescent types characterized by dense and lengthy upright hairs on their surface obstruct adult insects from laying eggs and hatching them. Varieties, including the wild type and other resistant types, possess high levels of silica and crude fiber, along with lower levels of ash and crude fat protein in the stem, which impede larval feeding and digestion. It is evident that biochemical factors play a more crucial role than morphological and physiological factors in deterring insects through non-preference and antibiosis. Numerous biochemical elements are recognized for their association with insect resistance in agricultural plants. Some of these constituents may serve as feeding cues for insects. The occurrence at lower concentrations or the total absence of such biochemical constituents leads to insect resistance. Biochemical constituents such as glycoalkaloid (solasodine), phenols, and phenolic oxidase enzymes, namely polyphenol oxidase and peroxidase, are present in brinjal [88]. These biochemical constituents possess insect-resistant properties, as outlined in Table 3. Achieving complete borer resistance would be challenging, and therefore, the development of tolerant genotypes is considered. When selecting genotypes for shoot and fruit borer resistance, apart from their performance, consideration may also be given to the quantity of biochemical constituents and the isozyme banding pattern [88].

6. MANAGEMENT OF BRINJAL SHOOT AND FRUIT BORER

6.1 Organic Manure

The effects of applying neem and *pongamia* to various plants on vegetables were evaluated by [47]. In insecticide-treated plots, the incidence of fruit borer and shoot borer in brinjal was initially between 30 and 50%, but it reduced to 6–10% following treatment. Research by Taher [103] demonstrated that the application of neem cake at a rate of 250 kg/ha increased the yield by approximately 68% and decreased the incidence of borer to 8%. According to [77, 112-113], okra treated with FYM and vermicompost had lower

percentages of fruit borer infestation. Islam [34] observed the impact of organic manures and fertilizers on the incidence of the fruit borer *L. orbonalis* and the brinjal shoot borer. Neem cake had the lowest incidence of fruit borer, 1.700 kg per hectare (6.08%). Nonetheless, it was discovered to be on par with vermicompost at 4000 kg per hectare, double the K₂O dose, and half the FYM + half the fertilizer dose. According to *L. orbonalis* incidence in potatoes may be effectively decreased with a single application of neem cake at 240 kg/ha. The effects of applying five different organic manures on *L. orbonalis* in brinjal: neem cake, pongamia cake, castor cake (all at 1.0 t/ha), farmyard manure, and vermicompost (10.0 t/ha). Neem was found to be the best cake of all [114].

6.2 Pheromone Traps

Ali [5] discovered that the output of marketable fruit was higher in the pheromone-treated plots compared to the control plots. They also found that, compared to the 1.5 m height, the 0.5 m height had a significantly higher number of insects caught. According to Das [19], delta and wing traps baited with synthetic *L. orbonalis* female sex pheromone were observed to capture and retain ten times more moths compared to either *Spodoptera* or uni-trap designs. Additionally, "windows" were incorporated into the side panels of delta traps, and the performance of locally constructed water and funnel traps was found to be comparable to that of delta traps [115]. However, the trap catches significantly increased from 0.4 to 2.3 moths per trap each night. When wing traps were positioned at crop height, they captured significantly more moths than when positioned 0.5 m above or below the canopy. However, according to Taher [18], deploying pheromone traps at a rate of 75 traps per hectare provided considerable protection against *L. orbonalis* in terms of production (28.67%), fruit damage (33.73%), and shoot damage (58.35%) in brinjal crops. Using sex pheromone traps, Rani (2013) studied fruit borer, *L. orbonalis*, and brinjal shoot in nine

villages in and around the Bangalore rural area between 2012 and 2013. The best trap heights for catching BSFB moths were assessed for each of the four variations. The findings showed that the greatest number of moth captures (499 moths) were found in traps at the highest elevation of 0.6 m above the crop canopy. Likewise, five other trap densities (i.e., 8, 16, 24, 32, and 40 traps/acre) were evaluated as well; the findings indicated that, at 16 traps/acre, the greatest number of moth captures (1097 moths) and the least degree of fruit damage (6.48%) were recorded.

6.3 Biopesticides and Botanicals

In order to combat brinjal shoot and fruit borer, Raina [80] compared several *B. thuringiensis* (Bt) formulations with neem and other pesticides. Among the various treatments, the greatest yield of marketable fruits (196.96 q/ha) and the smallest shoot (9.56%) and fruit (11.78%) infection were obtained with five sprays of Dipel 8L @ 0.2 percent spaced ten days apart. In their evaluation of novel insecticides against *L. orbonalis*, the aubergineshoots and fruit borer, Deshmukh and Bhamare (2006) contrasted them with traditional pesticides. To achieve a decrease in shoot infestation to 4.20%, reduce fruit infestation to 23.72% on a numerical basis and 25.30% on a weight basis, and increase fruit output to 78.73 q/ha, researchers found that cartap hydrochloride at a concentration of 0.1% was the most effective. Spinosad at a concentration of 0.01% was the next most effective option.

According to Prabhu [72], plots treated with Spinosad 2.5 SC (50g a.i./ha) had the lowest levels of fruit and shoot infestation (7.47 and 9.88%) throughout the West Bengal kharif season. The Spinosad treatment resulted in the highest marketable fruit production at 143.50 q/ha, followed by indoxacarb at 126.90 q/ha and emamectin benzoate at 121.30 q/ha. Anil and Sharma (2010) investigated the bio efficacy of several pesticides on brinjal c. v Arka Nidhi in 2007 and 2008 in Palampur against the shoot and fruit borer, *L. orbonalis*. They discovered that in the case of emamectin benzoate, there were comparatively few drooping shoots and fruit infection. However, agrospray oil T (0.2%) was determined to have the highest cost-benefit ratio.

In the kharif trials of 2007 and 2008, the most effective treatment was emamectin benzoate, which resulted in 5.0% and 4.8% shoot damage,

respectively. During the course of the two cropping seasons, emamectin benzoate was also shown to have the lowest fruit infestation (11.51, 11.44, and 12.39, 12.44) and the best output of healthy fruits (24.06, 23.14 t ha⁻¹). In Meerut, Uttar Pradesh, Harit [32] conducted an experiment. Spinosad 45 SC demonstrated the highest success in reducing shoot and fruit damage, resulting in the highest yield of 253.30 q/ha, followed by novaluron 10EC with a yield of 242.30 q/ha. However, they found that the highest cost-benefit ratio for no-valuron was 1:8.50, whereas the highest ratio for carbosulfan was 1:7.34.

6.4 Chemical Control

According to Singh [97], using deltamethrin at a rate of 25 g a.i./ha was found to be more effective than chlorpyrifos at 500 g a.i./ha in reducing fruit damage in brinjal on both a number and weight basis while also increasing the production of healthy fruits. Misra (2008) conducted field evaluations in Bhubaneswar during the winter of 2007 and the summer of 2008 to assess the effectiveness of two recently developed insecticides, namely rynaxypyr 20SC and flubendiamide 480 SC, on brinjal cultivar "Utkal Anushree" against the shoot and fruit borer, *L. orbonalis* [61]. Ten days after the fourth spray, Rynaxypyr 20SC @40 and 50g a.i./ha reduced shoot damage by 95–97%, fruit damage by 87–90% on a numerical basis, and weight damage by 88–90% when compared to the untreated control. During both seasons, the plots treated with rynaxypyr20SC @ 40 and 50g a.i. ha⁻¹ had the maximum healthy fruit output. Insecticides were tried by inthe Bapatla district of Andhra Pradesh against the shoot and fruit borer. In comparison to the untreated control (6666.66 kg/ha), they discovered that Profenofos (0.1%) boosted fruit yields (14312.05 kg/ha) and provided the largest decrease (42.7%) of *L. orbonalis* shoot damage. In their 2003 and 2004 study at Palampur, Plazas [69] evaluated the effectiveness of ten insecticidal treatments against the fruit borer and brinjal shoot. They found that acetamprid had the lowest levels of fruit and shoot infestation along with the highest profit and cost-benefit ratios (Rs 24,146/ha and 1:13.24). The most successful insecticide in lowering the weight-based number of *L. orbonalis* shoots (39.91%) and fruit infestations (18.21 and 17.48%) as well as increasing fruit yield (310.50 q/ha) was found to be Profenofos @ 0.1% in a chemical control trial carried out at Kanpur by Sreenivasa [99]. In brinjal, Raina [49] found that

cypermethrin 0.0075% was more effective than endosulfan 0.05% at controlling *L. orbonalis*. This finding is mostly consistent with the current research. In Pusa, Bihar, Srinivasan [100] conducted bio efficacy studies against *L. orbonalis* on brinjal. They found that the most effective treatments were imidacloprid at 0.025 kg a.i. ha⁻¹ and fenvalerate at 0.150 kg a.i. ha⁻¹. The maximum fruit yield was recorded at 290.25 q ha⁻¹ and 268.5 q ha⁻¹, respectively. However, the highest ICBR (1:14.41) was noted for fenvalerate @ 0.150 kg a.i. ha⁻¹, with imidacloprid (1:12.99) and cypermethrin (1:13.85) coming next. In Jalna during the kharif seasons of 2009 and 2010, Singh [94] evaluated that flubendamide 39.35SC and chlorantraniliprole 18.50SC outperformed other insecticides in decreasing *L. orbonalis* infestation and produced higher yield efficacy on Mahyco brinjal hybrid MHB 39.

Insecticides against shoot and fruit borer were assessed by Shirale [90] from Sabour (Bihar) in the kharifs 2010–11 and 2012–13. In areas where rynaxypyr 20 SC was applied, minimal levels of shoot infestation (5.67%), fruit infestation (12.59%), larvae per plot (2.36), and holes per fruit (0.40) were reported. Additionally, they observed that rynaxypyr had the greatest mean yield (346.69 q/ha). After conducting controlled field trials in two cropping seasons at Coimbatore, Latha [47] discovered that flubendamide 20 WG @ 75 g a.i./ha was the most efficient pesticide in reducing fruit and shoot damage. In both the winter and summer seasons, flubendamide exhibited the most significant reduction in shoot damage (96.8% and 97.2%), fruit damage (98.2% and 98.1%), and resulted in the highest yield (21.7 and 26.3 tons/ha). According to Jayakrishnan [87], deltamethrin was the most successful in lowering fruit damage (88.89%) and shoot damage (60.40%) when compared to the control on both a number and weight basis. With 132.27 q/ha, deltamethrin had the largest marketable fruit yield, whereas nimbecidine had the lowest (33.53 q/ha). The cost-benefit ratios for the insecticides were as follows: deltamethrin had the highest ratio at 1:8.7, followed by cypermethrin at 1:6.5, fenvalerate at 1:8.5, chlorpyrifos at 1:4.5, Preempt at 1:1.9, malathion at 1:0.6, and nimbecidine at 1:0.3 [89].

7. INTEGRATED PEST MANAGEMENT

A combination of plant products and herbicides can effectively inhibit fruit borer and brinjal

shoots, according to Singh [98]. Among the various treatments evaluated, the study identified that the basal application of neem cake at 20 q/ha combined with a foliar spray of quinalphos at 0.05% was effective in reducing the incidence of fruit borer to 20.63%. Additionally, according to Atwal [9], the combination of spinosad at 0.01%, *Metarhizium anisole*, chelating agent Fe-EDTA, and cartaphydrochloride at 0.1% proved to be the most effective Integrated Pest Management (IPM) approach against the fruit borer, *L. orbonalis*, resulting in the lowest shoot infection (7.47%) and the highest yield (81.82 q/ha). According to Aremu-Dele [26], mechanical removal of contaminated fruits and shoots combined with a pheromone trap and neem was determined to be the most efficient IPM module in decreasing shoot damage (86.69%).

Following a sequence, a reduction in shoot damage of 79.24%, 78.75%, and 78.55% was observed immediately after the implementation of pheromone traps mixed with neem, mechanical removal of contaminated fruits and shoots combined with pheromone traps, and traditional farmer's practices, respectively. Conversely, neem had the lowest effectiveness, with 54.46% of shoot infection [93]. The greatest protection against fruit infestation was found when infested fruits and shoots were mechanically removed along with a pheromone trap and neem (59.36% reduction). Subsequently, the methods employed by farmers resulted in a reduction of 54.13%, while the mechanical removal of infected fruits and shoots using a pheromone trap led to a reduction of 52.77%. These were then succeeded by the following techniques: pheromone trap combined with neem, mechanical removal of infested fruits and shoots with neem, and mechanical removal of infested fruits and shoots with protection, resulting in reductions of 47.70%, 43.69%, and 42.93%, respectively [96]. On the other hand, installing merely traps reduced fruit damage by at least 38.17%.

8. CONCLUSION

Brinjal Fruit and Shoot borer (*L. orbonalis* Guinee), is a monophagous insect that mainly feeds on Brinjal and other vegetables of Solanaceae family. Due to its short life cycle and boring nature, it heavily infests on the Brinjal plants and it has resulted huge losses in several nations of the world including Nepal. The management of this insect is of utmost importance to increase the yield of Brinjal and

other Solanaceae vegetables. The successful management of this pest can be brought about only by effective IPM practices. Apart from chemical pesticides bio-pesticides like Neem oil and Neem Leaf extract have great effectiveness against *L. orbonalis*. Botanical oils and extract of different plants such as Neem, Pungam, etc. are found to be very effective against the pests and insects. Biocontrol agents such as *Bacillus thuringiensis*, *Trichogramma chilonis* which is an egg parasitoid and larval parasitoid-*Trathala flavo-orbitalis* (Cameron) can also be utilized as means of potential parasitoids of this pest since they also show significant result in the reduction of shoot and fruit damage of Brinjal. Integrated pest control measure tactics such as breeding resistant cultivars, adopting good agronomic practices, mechanical, physical and biological control, and biorational control can do Eco-friendly management of brinjal shoot and fruit borer. Some cultural practices such as proper spacing, followed by clipping and burning of infested twig/ fruits/ stem, removal of alternate host, inter/trap crops (viz., coriander, cluster bean, fennel, chilly, radish, marigold, mint, onion, clover, fenugreek and cereal including maize) uses of organic amendment, and installing animated bird perches of T-shaped are optimal for getting high yields beside with eco-friendly management of the pest. Given this, the present review concluded that the use of IPM options, along with growing resistant varieties, good agronomic practices, biological control and chemical control (only if necessary) etc., reduce the unenthusiastic force of insecticides on the natural enemies, beneficial insect, pollinators, animal and human being that are present in the appropriate ecological niche and will defend the flora and fauna and the atmosphere from toxicological hazards contents.

Research gap: The bio-ecology and management of brinjal shoot and fruit borer pertains to the development and implementation of integrated pest management (IPM) strategies tailored to the specific needs and ecological context of brinjal cultivation. While conventional chemical pesticides are commonly used, there is a need for sustainable alternatives that minimize environmental impact and preserve natural ecosystems. Additionally, further investigation into the biology and behavior of the pest, as well as its interaction with the brinjal plant and surrounding environment, is warranted to inform the design of effective and holistic

management approaches for sustainable brinjal production.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Alam MZ, Sana DL. Biology of the brinjal shoot and fruit borer, *Leucinodes orbonalis* G.(*Pyralidae: Lepidoptera*) in East Pakistan. The scientist. 1962;5(1-4):13-24.
2. Alam MA, Rao PK, Rao BH. Biology of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee. Indian J. Agril. Sci. 1982;52(6):391-5.
3. Alam SN, Hossain MI, Rouf FM, Jhala RC, Patel MG, Rath LK, Sengupta A, Baral K, Shylesha AN, Satpathy S, Shivalingaswamy TM. Implementation and promotion of an IPM strategy for control of eggplant fruit and shoot borer in South Asia. AVRDC; 2006 Oct 4.
4. Alam SN, Rashid MA, Rouf FM, Jhala RC, Patel JR, Satpathy S, Shivalingaswamy TM, Rai S, Wahundeniya I, Cork A, Ammaranan C. Development of an integrated pest management strategy for eggplant fruit and shoot borer in South Asia. AVRDC-World Vegetable Center; 2003Dec 1.
5. Ali MH, Sanghi PM. Observations on oviposition, longevity and sex ratio of brinjal shoot and fruit borer *Leucinodes orbonalis*. Madras Agricultural Journal. 1962;49(8):267-8.
6. Alpuerto AB. Ecological studies and management of brinjal fruit and shoot borer, *Leucinodes orbonalis* Guenee. Indian Journal of Agricultural Sciences. 1994;52(6):391-5.
7. Asati BS, Sarnaik DA, Thakur BS, Guhey A. Shoot and fruit borer incidence as influenced by total phenol and total chlorophyll content in round fruited brinjal varieties. Orissa Journal of Horticulture. 2002;30(2):100-104.

8. Asmita S, Gotarkar SB, Jaya T, Dhumale UM. Integrated management of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guen. Journal of Applied Zoological Researches. 2006;17(2):189-91.
9. Atwal AS. Agricultural pests of India and South-East Asia. Agricultural Pests of India and South-East Asia; 1976.
10. Atwal AS, Dhaliwal GS. Agricultural pests of South Asia and their management. Kalyani Publishers; 2015.
11. Baral K, Roy BC, Rahim KM, Chatterjee H, Mondal P, Mondal D, Ghosh D, Talekar NS. Socio Economic Parameters of Pesticide Use and Assessment of Impact of an IPM Strategy for the Control of Eggplant Fruit and Shoot Borer in West Bengal, India. Tashkent, Uzbekistan: AVRDC; 2006 Oct 4.
12. Bhushan S, Chaurasia HK, Shanker RA. Efficacy and economics of pest management modules against brinjal shoot and fruit borer (*Leucinodes orbonalis*). The Bioscan. 2011;6(4):639-42.
13. Butani DK, Jotwani MG. Insect in vegetables periodical. Expert Book Agency D-42, Vivek Vihar, Dehli (India). 1984;6:220-36.
14. Butani DK, Varma SH. Pests of vegetables and their control: Brinjal. Pesticides. 1976;10(2):32-35.
15. Chakraborti S, Sarkar P. Management of *Leucinodes orbonalis* Guenee on eggplants during the rainy season in India. Journal of Plant Protection Research; 2011.
16. Hira C. Pheromones for the management of brinjal fruit and shoot borer, *Leucinodes orbonalis* Guenee. Karnataka Journal of Agricultural Sciences. 2009;22(3):594-6.
17. Nusra MS, Paranagama PA, Amarasinghe LD, Udukala DN. Pheromone baited biopesticide for control of *Leucinodes orbonalis* Guenee in brinjal plant. Frontiers in Bioscience. 2020;12(1):35-47.
18. Taher D, Solberg SØ, Prohens J, Chou YY, Rakha M, Wu TH. World vegetable center eggplant collection: origin, composition, seed dissemination and utilization in breeding. Frontiers in Plant Science. 2017 Aug 25;8:1484.
19. Das MS, Patnaik BH. A new host of the brinjal shoot and fruit borer *Leucinodes orbonalis* Guen, and its biology. Journal of the Bombay Natural History Society. 1971;67(3):601-603.
20. Arora R, Shera PS. bio-intensive ipm for ecofriendly management of insect pests.
21. Dhandapani N, Shelkar UR, Murugan M. Bio-intensive pest management in major vegetable crops: An Indian perspective. J Food Agric Environ. 2003;1(2):330-339.
22. Dhankar BS. Progress in resistance studies in the eggplant (*Solanum melongena* L.) against shoot and fruit borer (*Leucinodes orbonalis* Guen.) infestation.
23. Dutta P, Singha AK, Das P, Kalita S. Management of brinjal fruit and shoot borer, *Leucinodes orbonalis* Guenee in agro-ecological condition of West Tripura. Scholarly Journal of Agricultural Science. 2011;1(2):16-9.
24. Elanchezhyan K, Baskaran RK, Rajavel DS. Bio-chemical basis of resistance in brinjal genotypes to shoot and fruit borer, *Leucinodes orbonalis* Guen. Journal of Entomological Research. 2009;33(2):101-4.
25. Elekofehinti OO, Kamdem JP, Bolingon AA, Athayde ML, Lopes SR, Waczuk EP, Kade IJ, Adanlawo IG, Rocha JB. African eggplant (*Solanum anguivi* Lam.) fruit with bioactive polyphenolic compounds exerts *In vitro* antioxidant properties and inhibits Ca²⁺-induced mitochondrial swelling. Asian Pacific Journal of Tropical Biomedicine. 2013 Oct 1;3(10):757-66.
26. Aremu-Dele O, Sobowale IO, Nduka BA, Adesanya KA, Solomon O. Cocoa production improvement in some major producing countries of the world. In presentation at the conference natural ecosystem sustainability in the 21st Century, Ibadan, Nigeria; July 2022.
27. Fray A, Doganlar S, Daunay MC. Eggplant. In vegetables. Berlin, Heidelberg: Springer Berlin Heidelberg. 2007;287-313)
28. Frempong E. The nature of damage to egg plant (*Solanum melongena* L.) in Ghana by two important pests, *Leucinodes orbonalis* Gn. and *Euzophera villora* (Fldr.) (*Lepidoptera Pyralidae*). Bulletin de l'Institut Fondamental d'Afrique. 1979; 41(2):408-16.
29. Gangwar RK, Singh DV, Bhatnagar A, Sachan SK. Brinjal shoot and fruit borer management with insecticides. Indian Journal of Entomology. 2014;76(2):97-100.
30. Gautam M, Kafle S, Regmi B, Thapa G, Paudel S. Management of brinjal fruit and shoot borer (*Leucinodes orbonalis*

- Guenee) in Nepal. Acta Sci. Agric. 2019; 3(9):188-95.
31. Godase SK, Patel CB. Effect of organic manures and fertilizer doses on the incidence of brinjal shoot and fruit borer (*Leucinodes orbonalis* Guen). Pestology. 2003;27(1):5-6.
 32. Harit DN, Shukla GR. Laboratory biology of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guen.(Lepidoptera). Journal of Experimental Zoology, India. 2005;8(2): 307-311.
 33. Herbst ST. The new food lover's companion: Comprehensive definitions of nearly 6000 food, drink, and culinary terms. (No Title); 2001.
 34. Islam MN, Karim MA, Mannan MA, Chowdhury JC, Islam M. Integrated management of the brinjal shoot and fruit borer, *Leucinodes orbonalis* Guen.(Lepidoptera: Pyralidae), based on insecticide. Bangladesh Journal of Entomology. 1999;9(1/2):75-85.
 35. Islam MS, Choudhury MA, Maleque MA, Mondal MF, Hassan K, Khan AU. Management of brinjal shoot and fruit borer (*Leucinodes orbonalis* Guen.) using selected bio-rational insecticides. Fundamental and Applied Agriculture. 2019 Nov 17;4(4):1025-31.
 36. Jagginavar SB, Sunitha ND, Biradar AP. Bioefficacy of flubendiamide 480 SC against brinjal fruit and shoot borer, *Leucinodes orbonalis* Guen. Karnataka Journal of Agricultural Sciences. 2009; 22(3):712-3.
 37. Jat KL, Pareek BL. Biophysical and biochemical factors of resistance in brinjal against *Leucinodes orbonalis*. Indian Journal of Entomology. 2003;65(2):252-258.
 38. Javed H, Mohsin A, Aslam M, Naeem M, Amjad M, Mahmood T. Relationship between morphological characters of different aubergine cultivars and fruit infestation by *Leucinodes orbonalis* Guenee. Pak. J. Bot. 2011 Aug 1;43(4): 2023-8.
 39. Jayaraj J, Manisegaran S. Management of fruit and shoot borer in brinjal. The Hindu Sci-Tech. Agri. College and Res. Inst. Madurai. 2010.
 40. Kaur P, Yadav GS, Wargantiwar RK, Burange PS. Population dynamics of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee (Lepidoptera: Crambidae) under agroclimatic conditions of Hisar, Haryana, India. The Ecoscan. 2014;8(1&2):1-5.
 41. Kavitha VS, Revathi N, Kingsley S. Biology of brinjal pest, *Leucinodes orbonalis* Guen of Erode region in Tamil Nadu. Journal of Entomological Research. 2008;32(3):255-7.
 42. Khorsheduzzaman AK, Alam MZ, Rahman MM, Mian MK, Mian MI. Biochemical basis of resistance in eggplant (*Solanum melongena* L.) to *Leucinodes orbonalis* Guenee and their correlation with shoot and fruit infestation. Bangladesh Journal of Agricultural Research. 2010;35(1):149-155.
 43. Krishnamoorthy SV, Banuchitra R, Devi SJ, Karuppuchamy P. Management of brinjal shoot and fruit borer *Leucinodes orbonalis* Guenee with novel insecticides. Journal of Insect Science (Ludhiana). 2014;27(1):17-22.
 44. Kumar NK, Sadashiva AT. *Solanum macrocarpon*: a wild species of brinjal resistant to brinjal shoot and fruit borer, *Leucinodes orbonalis* (Guen.). Insect Environment. 1996;2(2): 41-2.
 45. Kumar S, Singh D. Seasonal incidence and economic losses of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee. Agricultural Science Digest-A Research Journal. 2013;33(2):98-103.
 46. Latif MA, Rahman MM, Alam MZ. Efficacy of nine insecticides against shoot and fruit borer, *Leucinodes orbonalis* Guenee (Lepidoptera: Pyralidae) in eggplant. Journal of Pest Science. 2010 Dec;83:391-7.
 47. Latha VP, Nagar S, Chaitanya G. Seasonal incidence of brinjal shoot and fruit borer, *Leucinodes orbonalis* guenee,(Lepidoptera: Crambidae) during Kharif Season in 2017. Journal of Pharmacognosy and Phytochemistry. 2018;7(5):3261-2.
 48. Anjali Mathur AM, Singh NP, Mahesh Meena MM, Swaroop Singh SS. Seasonal incidence and effect of abiotic factors on population dynamics of major insect pests on Brinjal crop.
 49. Raina J, Yadav GS. Brinjal shoot and fruit borer: Bio-ecology and management. Journal of Pharmacognosy and Phytochemistry. 2018;7(4):444-9.
 50. Matsubara K, Kaneyuki T, Miyake T, Mori M. Antiangiogenic activity of nasunin, an antioxidant anthocyanin, in eggplant peels.

- Journal of Agricultural and Food Chemistry. 2005 Aug 10;53(16):6272-5.
51. Krattiger A, Mahoney RT, Nelsen L, Thomson JA, Bennett AB, Satyanarayana K, Graff GD, Fernandez C, Kowalski S. Intellectual Property Management in Health and Agricultural Innovation: A Handbook of Best Practices; 2.
 52. Meena GS, Pachori RK, Panse R. Extent of damage and seasonal incidence of *Leucinodes orbonalis* (Guen.) on Brinjal. Annals of Plant Protection Sciences. 2012; 20(1):114-6.
 53. Mehto DN, Singh KM, Singh RN, Prasad D. Biology of brinjal fruit and shoot borer, *Leucinodes orbonalis* Guen. Bulletin of Entomology. 1983;24(2):112-5.
 54. Mennella G, Lo Scalzo R, Fibiani M, D'Alessandro A, Francese G, Toppino L, Acciarri N, de Almeida AE, Rotino GL. Chemical and bioactive quality traits during fruit ripening in eggplant (*S. melongena* L.) and allied species. Journal of Agricultural and Food Chemistry. 2012 Nov 28;60(47): 11821-31.
 55. Misra HP. New promising insecticides for the management of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee. Pest Management in Horticultural Ecosystems. 2008;14(2):140-7.
 56. Shaukat MA, Mustafa MA, Ammad A, Maqsood S, Umer H, Mustafa F, Malik G. Life aspects and mode of damage of brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee) on eggplant (*Solanum melongena* Linnaeus): A review. International J Entom Res. 2018 Mar; 3(2).
 57. Muthukumaran N, Kathirvelu C. Preliminary screening of different accessions of eggplants on the biology of shoot and fruit borer (*Leucinodes orbonalis*). Insect Environment. 2007; 12(4):173-174.
 58. Nandi C, Narabanchi G, Jakatimath S, Prafulkumar MV. Seasonal incidence of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee, (*Lepidoptera: Crambidae*) during rabi season. Journal of Experimental Zoology India. 2017 Jul 1;20(2).
 59. Naqvi AR, Pareek BL, Nanda US, Mitharwal BS. Leaf morphology and biochemical studies on different varieties of brinjal in relation to major sucking insect pests.
 60. Netam M, Lakra R, Koshta VK, Sharma D, Deole S. Screening of Shoot and Fruit Borer (*Leucinodes orbonalis* Guenee.), for Resistance in Brinjal (*Solanum melongena* L.) Germplasm Lines. Int J Curr Microbiol App Sci. 2018;7(02):3700-6.
 61. MALLIK H. Comparative study for the management of brinjal shoot and fruit borer and Epilachna beetle with botanicals and some selected chemical insecticides (doctoral dissertation, department of entomology, Sher-e-Bangla Agricultural University, Dhaka-1207).
 62. Ahmad H, Hasan MR, Rahul S, Mahbuba S, Uddin AJ. Growth and yield analysis of some exotic brinjal lines. Bangladesh Research Publication Journal. 2016;12(2): 112-6.
 63. Patel LC, Konar AM, Sahoo CS. Biophysical screening of brinjal genotype s against fruit and shoot borer, *Leucinodes orbonalis* (GUEN). The Bioscan. 2015 May 24;10:905-9.
 64. Anjana P, Mehta PK. Pest complex of brinjal and their succession under mid hills of Himachal Pradesh. Journal of Insect Science (Ludhiana). 2008;21(1):67-71.
 65. Anjana P, Mehta PK, Sharma PC. Field efficacy of some insecticides and biopesticides against *Leucinodes orbonalis* Guenee on brinjal. Environment and Ecology. 2009;27(1A):320-5.
 66. Anjana P, Mehta PK, Sood AK. Developmental biology of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee in mid-hills of Himachal Pradesh. Entomon. 2007;32(2):137-41.
 67. Patil PD. Technique for mass rearing of the brinjal shoot and fruit borer, *Leucinodes orbonalis* Guen. Journal of Entomological Research. 1990;14(2):164-72.
 68. Patra S, Chatterjee ML, Mondal S, Samanta A. Field evaluation of some new insecticides against brinjal shoot and fruit borer, *Leucinodes orbonalis* (Guen.). Pesticide Research Journal. 2009;21(1): 58-60.
 69. Plazas M, Andujar I, Vilanova S, Hurtado M, Gramazio P, Herraiz FJ, Prohens J. Breeding for chlorogenic acid content in eggplant: interest and prospects. Notulae Botanicae Horti Agrobotanici Cluj-Napoca. 2013 May 28;41(1):26-35.
 70. Singla P, Bhullar MB, Kaur P. Biological studies on brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee. Journal of

- Entomology and Zoology Studies. 2018; 6(1):161-5.
71. Prabhu M, Natarajan S, Pugalendhi L. Biochemical basis of shoot and fruit borer (*Leucinodes orbonalis* G.) resistance in brinjal (*Solanum melongena* L.). In Recent trends in horticultural biotechnology, ICAR national symposium on biotechnological interventions for improvement of horticultural crops: issues and strategies, Vellanikkara, Kerala, India, 10-12 January, 2005. New India Publishing Agency. 2007;11:829-836.
 72. Prabhu M, Natarajan S, Veeraragavathatham D, Pugalendhi L. The biochemical basis of shoot and fruit borer resistance in interspecific progenies of brinjal (*Solanum melongena*). Eur Asian Journal of Bio Sciences. 2009 Dec 1;3(1):50-7.
 73. Prakash YS, Bhadoria PB, Amitava Rakshit AR. Relative efficacy of organic manures in improving resistance and pest tolerance of okra (cv. Parbhani Kranti).
 74. Prasad TV, Bhardwaj R, Gangopadhyay KK, Arivalagan M, Bag MK, Meena BL, Dutta M. Biophysical and biochemical basis of resistance to fruit and shoot borer (*Leucinodes orbonalis* Guenee) in eggplant. Indian Journal of Horticulture. 2014;71(1):67-71.
 75. Preneetha S. Breeding for shoot and fruit borer (*Leucinodes orbonalis* Guenee) resistance in brinjal (*Solanum melongena* L.) (Doctoral dissertation, Tamil Nadu Agricultural University).
 76. Puranik TR, Hadapad AB, Salunkhe GN, Pokharakar DS. Management of shoot and fruit borer, *Leucinodes orbonalis* Guenee through *Bacillus thuringiensis* formulations on brinjal. Journal of Entomological Research. 2002;26(3):229-32.
 77. Rahman AK. Screening of 28 brinjal line for resistance/tolerance against the brinjal shoot and fruit borer. Annual Report. Entomology Division, Bangladesh Agricultural Research Institute, Joydebpur, Gazipur, Bangladesh. 1997:32-5.
 78. Dash L, Lakshmi VR, Padhy D. Host range, life aspects and mode of damage of brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee) on brinjal: A review. International Journal of Agriculture, Environment and Biotechnology. 2020; 13(1):81-5.
 79. Rai AB, Vora VT, Patel CV. Investigations on the integrated control of *Tetranychus urticae* (Acari: Tetranychidae) infesting brinjal in South Gujarat. In Abstract presented in 5th national symposium on acarology, Bangalore. 1995 Sep 20;20:22.
 80. Raina J, Yadav GS. Brinjal shoot and fruit borer: Bio-ecology and management. Journal of Pharmacognosy and Phytochemistry. 2018;7(4):444-9.
 81. Raina J, Yadav GS. Influence of abiotic factors on population dynamics of *Leucinodes orbonalis* Guenee on brinjal in Hisar agro-climatic conditions. Annals of Plant Protection Sciences. 2017;25(2):277-80.
 82. Raina J, Yadav GS. Studies on biological parameters of *leucinodes orbonalis* guenee on brinjal under laboratory conditions. Journal of Experimental Zoology India. 2017 Jul 1;20(2).
 83. Raina J, Yadav GS, Sharma SS. Bioefficacy of some insecticides against shoot and fruit borer, *Leucinodes orbonalis* Guenee on brinjal under Hisar agro-climatic conditions during kharif season. Journal of Applied and Natural Science. 2016 Dec 1;8(4):1801-5.
 84. Rotino GL, Perri E, Acciarri N, Sunseri F, Arpaia S. Development of eggplant varietal resistance to insects and diseases via plant breeding. Advances in Horticultural Science. 1997 Jan 1:193-201.
 85. Saeed MQ, Khan IA. Population abundance and chemical control of brinjal fruit borer *Leucinodes orbonalis* Guen. (*Pyrilidae*; *Lepidoptera*).
 86. Saha T, Chandran N, Kumar R, Ray SN. Field efficacy of newer insecticides against brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee (*Lepidoptera*: *Pyrilidae*) in Bihar. Pesticide Research Journal. 2014;26(1):63-7.
 87. Jayakrishnan S, Madhuban G. Evaluation of synthetic and natural insecticides for the management of insect pest control of eggplant (*Solanum melongena* L.) and pesticide residue dissipation pattern. American Journal of Plant Sciences. 2012 Feb 27;2012.
 88. Raina J, Yadav GS. Brinjal shoot and fruit borer: Bio-ecology and management. Journal of Pharmacognosy and Phytochemistry. 2018;7(4):444-9.
 89. Shaukat MA, Ahmad A, Mustafa F. Evaluation of resistance in brinjal (*Solanum melongena* L.) against brinjal shoot and fruit borer (*Leucinodes orbonalis* Guen.) infestation: A review. International

- Journal of Applied Sciences and Biotechnology. 2018 Oct 1;6(3):199-206.
90. Shirale D, Patil M, Zehr U, Parimi S. Evaluation of newer insecticides for the management of brinjal fruit and shoot borer *Leucinodes orbonalis* (Guenee). Indian Journal of Plant Protection. 2012;40(4):273-5.
 91. Shukla A, Khatri SN. Incidence and abundance of brinjal shoot and fruit borer *Leucinodes orbonalis* Guenee. Bioscan. 2010 Apr 17;5(2):305-8.
 92. Shukla V, Naik LB. Agro-techniques for solanaceous vegetables. Advances in Horticulture; 1993.
 93. Singh DK, Singh R, Datta SD, Singh SK. Seasonal Incidence and Insecticidal Management of Shoot and Fruit Borer (*Leucinodes orbonalis* Guenee) in Brinjal. Annals of Horticulture. 2009;2(2):187-90.
 94. Singh PK. Control of brinjal shoot and fruit borer, *Leucinodes orbonalis* with combination of insecticides and plant extracts.
 95. Singh RK, Devjani P, Singh TK. Population dynamics of *Leucinodes orbonalis*. Annals of Plant Protection Sciences. 2009;17(2):486-7.
 96. Singh SP, Kumar N. Insecticidal evaluation against *Leucinodes orbonalis* Guen on brinjal. Pesticide Research Journal. 2011; 23(2):227-9.
 97. Singh TH, Kalda TS. Source of resistance to shoot and fruit borer in eggplant (*Solanum melongena* L.). PKV Research Journal. 1997;21(2):126-8.
 98. Singh YP, Singh PP. Biology of shoot and fruit borer (*Leucinodes orbonalis* of egg plant at medium high altitude hills of Meghalaya. Indian Journal of Entomology. 2001;63(4):373-6.
 99. Sreenivasa Murthy D, ward Raja ED, Shivanna M. Economic assessment of neem cake to management fruit borer in Brinjal. In Proc. Second Nation. Symp. Int. Pest Mgmt. Hort. Crops: New Molecules, Biopest. Environ, Bangalore. 2001 Oct 17;13.
 100. Srinivasan R. Integrated Pest Management for eggplant fruit and shoot borer (*Leucinodes orbonalis*) in south and Southeast Asia: Past, Present and Future. Journal of Biopesticides. 2008;1(2):105-12.
 101. Srinivasan R. Insect and mite pests on eggplant. AVRDC-World Vegetable Center; 2009.
 102. Stommel JR, Whitaker BD, Haynes KG, Prohens J. Genotypex environment interactions in eggplant for fruit phenolic acid content. Euphytica. 2015 Oct;205:823-36.
 103. Taher D, Solberg SØ, Prohens J, Chou YY, Rakha M, Wu TH. World vegetable center eggplant collection: origin, composition, seed dissemination and utilization in breeding. Frontiers in Plant Science. 2017 Aug 25;8:1484.
 104. Thapa RB. Integrated management of brinjal fruit and shoot borer, *Leucinodes orbonalis* Guen: An overview. J. Inst. Agric. Anim. Sci. 2010;30(32):1-6.
 105. Adeyeye EI. The contribution of flesh, placenta and seeds to the nutritional attributes of a variety of *Capsicum annum* (Bell pepper). Elixir Food Sci. 2014 Feb 28;68:22587-94.
 106. Tiwari G, Prasad CS, Kumar A, Nath L. Influence of weather factors on population fluctuation of pest complex on brinjal. Annals of Plant Protection Sciences. 2012;20(1):68-71.
 107. Varma S, Anandhi P, Singh RK. Seasonal incidence and management of brinjal shoot and fruit borer, *Leucinodes orbonalis*. Journal of Entomological Research. 2009;33(4):323-9.
 108. Venkateswarlu P, Sitaramaiah S, Sreedhar U, Rao SG, Sawant SK, Rao SN. Role of organic and inorganic manures on the incidence of insect pests and their natural enemies in rabi groundnut. In Biological control of lepidopteran pests. Proceedings of the Symposium of Biological Control of Lepidopteran Pests, July 17-18, 2002, Bangalore, India. Society for Biocontrol Advancement. 2003;295-299.
 109. Wankhede SM, Kale VD. Effect of insecticides on brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee. Annals of Plant Protection Sciences. 2010;18(2):336-9.
 110. Paudel P, Pandey MK, Subedi M, Paudel P, Kumar R. Genomic approaches for improving drought tolerance in wheat (*Triticum aestivum* L.): A comprehensive review. Plant Archives. 2024;24(1):1289-300.
 111. Tutlani A, Kumar R, Kumari S, Chouhan S. Correlation and path analysis for yield and its phenological, physiological, morphological and biochemical traits under salinity stress in chickpea (*Cicer arietinum* L.). International Journal of Bio-resource

- and Stress Management. 2023;14(Jun, 6):878-90.
112. Chaudhary PL, Kumar B, Kumar R. Analysis of heterosis and heterobeltiosis for earliness, yield and its contributing traits in okra (*Abelmoschus esculentus* L. Moench). International Journal of Plant and Soil Science. 2023;35(11): 84-98.
113. Kumar R, Pandey MK, Pitha CC, Mehandi S, Chaudhary PL. Analysis of heterotic potential for yield and its attributing traits in okra (*Abelmoschus esculentus* L. Moench). Electronic Journal of Plant Breeding. 2023;14(4):1515-26.
114. Kumari S, Kumar R, Chouhan S, Chaudhary PL. Influence of various organic amendments on growth and yield attributes of mung bean (*Vigna radiata* L.). International Journal of Plant & Soil Science. 2023;35(12):124-30.
115. Chouhan S, Kumari S, Kumar R, Chaudhary PL. Climate resilient water management for sustainable agriculture. Int. J. Environ. Clim. Change. 2023; 13(7):411-26.
116. Gesraha, Mohamed Ahmed, and Amany Ramadan Ebeid. Evaluate the effects of potential botanical and conventional insecticides on the reproductive and developmental aspects of the pest agrotis ipsilon (*Lepidoptera: Noctuidae*). Asian Journal of Biology. 2021;10(4):83-91. <https://doi.org/10.9734/ajob/2020/v10i430128>.
117. Tiwari, Awanindra Kumar. IPM essentials: Combining biology, ecology, and agriculture for sustainable pest control. Journal of Advances in Biology & Biotechnology. 2024;27(2):39-47. Available:<https://doi.org/10.9734/jabb/2024/v27i2697>.
118. Rahman MM, Rahman MM, Ali MR. Evaluation of some selected options for managing brinjal shoot and fruit borer in two intensive brinjal growing areas. World Journal of Zoology. 2009;4(3):169-75.
119. Abrol DP, Singh JB. Relative efficacy of some insecticides against brinjal fruit and shoot borer, *Leucinodes orbonalis* Guen and their impact on fruit yield. Journal of Asia-Pacific Entomology. 2003 May 1;6(1):83-90.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/119027>