



# **Emerging Role of Probiotics in Advancement of Combating Physical Abnormalities and Diseases: A Systematic Perspective Analysis**

**Mst. Tania Khatun <sup>a</sup>, Forkanul Hoque <sup>a</sup>,  
Naeema Salatia Hoque <sup>b</sup>, Md. Sakhawat Hossain <sup>b</sup>,  
Md. Ashrafal Alam <sup>c</sup>, Sabrina Afrin <sup>c</sup>,  
Tanjina Nasrin Eva <sup>c</sup> and Rahima Tanbin Tama <sup>b\*</sup>**

<sup>a</sup> Department of Pharmacy, Islamic University, Kushtia-7003, Bangladesh.

<sup>b</sup> Department of Biotechnology and Genetic Engineering, Islamic University, Kushtia-7003, Bangladesh.

<sup>c</sup> Department of Applied Chemistry and Chemical Engineering, Islamic University, Kushtia-7003, Bangladesh.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. Author MTK conceptualized the study, performed methodology, did formal analysis, Data curation, wrote original draft and did Funding Acquisition. Author FH did data curation and funding acquisition. Author NSH wrote original draft, did Data curation and funding acquisition. Authors MSH, SA and TNE did data visualization and Funding acquisition. Author MAA did software analyses and funding acquisition. Author RTT supervised and investigated the study, performed methodology, did Software analysis, Data curation, wrote, reviewed and edited the manuscript and did funding acquisition. All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/ajbgmb/2024/v16i8397>

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119570>


\*Corresponding author: E-mail: [tanbintama.btge@std.iu.ac.bd](mailto:tanbintama.btge@std.iu.ac.bd);

**Cite as:** Khatun, Mst. Tania, Forkanul Hoque, Naeema Salatia Hoque, Md. Sakhawat Hossain, Md. Ashrafal Alam, Sabrina Afrin, Tanjina Nasrin Eva, and Rahima Tanbin Tama. 2024. "Emerging Role of Probiotics in Advancement of Combating Physical Abnormalities and Diseases: A Systematic Perspective Analysis". *Asian Journal of Biochemistry, Genetics and Molecular Biology* 16 (8):1-23. <https://doi.org/10.9734/ajbgmb/2024/v16i8397>.

## ABSTRACT

Probiotics are live microorganisms, primarily bacteria that give the host health benefits when given in sufficient amounts. It works in various defensive mechanisms, including releasing antimicrobial compounds, competitive adhesion to the mucosa and epithelium, fortifying the gut epithelial barrier and regulating the immune system. Probiotics have a considerable effect in reducing allergies and gastrointestinal problems like diarrhoea and ulcers. Probiotics are quite beneficial for managing and preventing complex diseases including diabetes, cardiovascular diseases, anaemia, urogenital diseases, dental caries, URTI, cancers and various abnormalities like obesity, depression, etc. Nowadays, Antibiotic resistance is a global challenge associated with high morbidity and mortality. The current lack of efficacious treatments necessitates the creation of innovative therapeutic approaches and substitute antimicrobial therapies. So, natural products or probiotics would be a better option for treating human health. Therefore, we systematically reviewed the sources, including Google Scholar, PubMed, Science Direct, Jstor and Base; studies only about human health were included here. All types of animal diseases and articles before 1990 are excluded. A total of 6,410 articles were identified and 300 articles were taken for further evaluation based on context. However, Probiotics have already become a successful therapeutic intervention strategy among physicians. This investigation covers a comprehensive overview of the current state of probiotic research which comprises a wide range of topics including sourcing, mode of action, probiotic strains, isolating procedures and the role of probiotics.

### Graphical abstract



**Keywords:** Abnormality; bacteria; diseases; defensive mechanism; probiotics.

## 1. INTRODUCTION

Antibiotics have been widely employed in managing bacterial infections because of their capacity to either stop the growth or eradicate living microbes [1]. However, the spread of antibiotic-resistance genes into harmful

bacteria has sparked worries about the short-term efficacy of the current antibiotic inventory [2]. Due to the global expansion of antibiotic and antimicrobial resistance, the World Health Organization has classified these concerns as an unanticipated global health danger with wide-ranging effects on the safety of humans, animals,

food and the environment across numerous sectors [2]. Thus, Probiotics are considered one of the best substitute treatments [3]. German physicist Werner Kollath first used the term "probiotic" in 1953 [4]. It is a combination of the Latin word "pro" and the Greek word "βίος," which means "for life" [4]. Kollath uses this word to describe the "active substances" that are essential to the healthy growth and well-being of living things [4]. According to the definition given to probiotics in 1965, they are chemicals secreted by microbes that encourage the growth of other microorganisms [5]. Probiotics are defined by the World Health Organization (WHO) as "living microorganisms that, when administered in adequate amounts, confer benefits to the host's health" [6].

Probiotics which can exist in food, supplements and the intestine are a significant class of advantageous bacteria that are taken or added to diets [7]. These bacteria are capable of colonizing the intestine and competing with dangerous microorganisms. Additionally, these bacteria produce certain enzymes that aid in feed breakdown. Probiotics can enhance immune activities by interacting with various immune cells and altering the composition of gut bacteria when ingested [8-10]. Fundamental selection criteria

should generally be used in the selection and characterization of novel probiotic strains [11]. The Food and Agriculture Organization (FAO) and the World Health Organization (WHO) of the United Nations released guidelines in 2002 outlining the various factors that need to be considered when choosing probiotic microorganisms [11]. These factors include antimicrobial activity, the ability to adhere to epithelial tissues, resistance to harsh conditions inside the human body and safety for use [11]. Furthermore, a probiotic must be highly able to endure under intestinal circumstances, such as acidic pH, enzymes, bile salts, etc. [12]. They have no harmful side effects and are not poisonous or pathogenic [13]. To preserve the intended qualities, they are also appropriately labelled and compatible with the product matrix, processing, and storage circumstances [13]. It has been demonstrated that consuming  $10^9 - 10^{10}$  viable colony-forming units (CFU) daily improves host health [14]. So, probiotics are frequently prescribed to enhance health in nations like Canada, Switzerland and Japan [15]. This manuscript reviews available information and summarizes the current research knowledge about probiotics, their defensive mechanism, sources, isolation techniques, and the effects of various diseases that impact human health.

**Table 1. List of some probiotic bacteria (gram-positive and gram-negative).**

Previous Generation Probiotics		
Bacteria Type	Bacteria Name	References
Gram-positive	<i>Lactobacillus acidophilus</i> , <i>Lactobacillus casei</i> , <i>Lactobacillus fermentum</i> , <i>Lactobacillus gasseri</i> , <i>Lactobacillus johnsonii</i> , <i>Lactobacillus paracasei</i> , <i>Lactobacillus plantarum</i> , <i>Lactobacillus salivarius</i> , <i>Bifidobacterium adolescentis</i> , <i>Bifidobacterium animalis</i> , <i>Bifidobacterium bifidum</i> , <i>bifidobacterium genera</i> , <i>Leuconostoc</i> , <i>Enterococcus</i> , <i>Streptococcus salivarius</i> , <i>Bifidobacterium longum</i> , <i>Lactobacillus crispatus</i> , <i>Lacticaseibacillus rhamnosus</i> , <i>Lacticaseibacillus casei</i> , <i>Lactiplantibacillus plantarum</i> , <i>Limosilactobacillus reuteri</i> , <i>Levilactobacillus brevis</i> , <i>Ligilactobacillus salivarius</i> , <i>Lactococcus lactis</i> , <i>Pediococcus mesenteroides</i> , <i>Streptococcus thermophiles</i> , <i>Bacillus subtilis</i> , <i>Bacillus coagulans</i> .	[16] [17] [18] [19] [20] [21] [22]
Gram-negative	<i>Escherichia coli</i> , <i>Veillonella sp.</i>	[23] [24] [25]
New Generation Probiotics		
Bacteria type	Bacteria name	References
Gram-positive	<i>Clostridium butyricum</i> , <i>Anaerobutyricum hallii</i> , <i>Roseburia spp.</i>	[26]
Gram-negative	<i>Akkermansia muciniphila</i> , <i>Faecalibacterium prausnitzii</i> , <i>Bacteroides spp.</i> , <i>Bacteroides uniformis</i> , <i>Bacteroides acidifaciens</i> , <i>Bacteroides thetaiotaomicron</i> , <i>Prevotella copri</i> , <i>Christensenella minuta</i> , <i>Parabacteroides goldsteinii</i> .	[27] [28] [29]

## 1.1 Mode of Action of Probiotics

Though their exact processes are unclear, probiotics are thought to have several positive impacts [13]. Probiotics work against various pathogens in several ways, including releasing antimicrobial compounds, competitive adhesion to the mucosa and epithelium, fortifying the gut epithelial barrier and regulating the immune system in Fig. 1 [13]. Probiotics can increase the intestinal defence system through various strategies, including acting as an antagonist against pathogens, enhancing the intestinal epithelial layer, producing neurotransmitters, increasing innate immunity, and supporting adaptive immunity in Fig. 1 [30,31]. The intestinal barrier is one important defensive mechanism that keeps the organism safe from the environment and preserves epithelial integrity [32]. The mucous layer, antimicrobial peptides, secretory IgA and the epithelial junction adhesion complex are the intestinal barrier's defences [32]. When this barrier function is damaged, food antigens and bacteria can enter the submucosa and cause inflammatory reactions that could lead to digestive problems. Probiotic microorganisms have been thoroughly investigated in terms of their role in maintaining this barrier. Mucin glycoproteins, often called mucins, are important macromolecules that make up epithelial mucus and have been linked to health and illness. Probiotics may increase mucus secretion to enhance barrier performance and pathogen exclusion in Fig. 1 [18].

It is thought that adhesion to intestinal mucosa is necessary for colonization and plays a crucial role in the relationship between probiotic strains and their host. The particular interaction has suggested a potential link between probiotic bacterial surface proteins and pathogens' competitive removal from mucus [33,34]. Probiotics also control innate and adaptive immune responses by influencing B and T lymphocytes, dendritic cells (DC) and macrophages [35]. Probiotics interact with intestinal epithelial cells and draw macrophages and mononuclear cells, increasing the generation of anti-inflammatory cytokines [35]. Additionally, probiotics can use the gut-brain axis to create neurotransmitters in the gut. The amounts of serotonin, gamma-aminobutyric acid (GABA) and dopamine can be modulated by some probiotic strains which can impact mood, behaviour, gut motility and stress-related pathways [36-38].

## 1.2 Sources of Probiotics

Probiotic foods are the subject of intense interest due to their positive effects on human health [39]. Thus, there is a corresponding increase in the study of probiotic microbes [39]. Consequently, probiotics may be presented in many ecological niches besides humans [40]. Probiotics are becoming increasingly common in the food, pharmaceutical and fermentation industries, and customers generally accept them favourably [40]. Besides, probiotic-rich foods include dairy and dairy-related products [41]. The primary supplies could come from human origins such as breast milk or the large or small intestine of humans [41]. Milk is an excellent medium for both good and bad microbes due to its high nutritional content [42]. Moreover, the best source of probiotics for human consumption is the faeces of newborns, kids and adults [43]. Their capacity to withstand gastrointestinal transit and colonize the intestines for beneficial purposes is the reason behind this [43]. Additionally, foods of animal origin and foods of plant origin may provide additional sources for isolating probiotic strains [44]. Probiotics found in animal origin can be divided into three categories such as meat products (emulsified, fermented, and restructured products), milk and dairy products (yogurt, cheese, and ice cream) and fish and seafood (keureling fish, *labeo rohita* and juvenile rainbow trout) [45-47].

Probiotics can also be found in plants of origin. They can be divided into four categories such as pickled vegetables (kimchi, sauerkraut and humble pickle), grain (bran, barley and oats), non-fermented fruits (banana, almonds and watermelon) and non-fermented vegetables (cucumbers, carrots and radishes) [48-50]. However, Supplements containing probiotics are also offered [51]. The bacterial strains found in probiotic foods and supplements should be essential for the following processes: lowering and metabolism of cholesterol; colonization of the intestinal, respiratory and urogenital tracts; inhibition of carcinogenesis, either directly or indirectly, through immune system stimulation; lactose metabolism; calcium absorption and vitamin synthesis potential; the reductive potential of yeast and vaginal infection; reduction of constipation and diarrheal disorders; mitigation of gastritis and ulcers; reduction of acne, rash face and skin issues; and the synthesis of natural antimicrobials [51-53].




Fig. 1. Major mechanisms of action of probiotics [16-18]





Fig. 2. Features of probiotics [21-24]

### 1.3 Isolation of Probiotics

According to FAO/WHO guidelines, the first step is to isolate possible probiotic strains [55]. Researchers have determined that new probiotic bacterial strains are still needed for certain therapeutic applications [55]. The probiotics derived from dairy products differ in several ways from the probiotics isolated from the intestines of people and animals [56]. A prominent distinction between these probiotics is their adherence. The adhesion activity of intestinal isolates is often higher than that of dairy isolates [56]. To extract probiotic strains from dairy products, samples were gathered from various sources, including camel milk, sheep yogurt, cow yogurt, goat milk

and native buttermilk [57]. Also, the sample was taken from clinical samples including blood, urine, respiratory, pericardial, abdominal and cerebral fluid (CSF) [58]. These samples were cultured on agar and broth media and incubated for 24.0 to 48.0 hours at 37.0 °C [57]. The strains were identified using a variety of tests, such as Gram staining, catalase testing, growth at temperatures between 15.0 °C and 45.0 °C, production of acid and gas from glucose, production of ammonia from arginine, and fermentation of sugars (arabinose, cellobiose, mannitol, mannose, melibiose, raffinose, ribose, salicin, rhamnose and xylose) [57-59].





**Fig. 3. Probiotic sources: animal and plant-origin foods [54].**

## 2. METHODOLOGY

Different sources were generated for article searches including Google Scholar, PubMed, Science Direct, Jstor, and Base. The information mainly searches in several articles from 2000 to July 2024. The keywords that were used for searches included probiotics, advancements and disease management. The obtained articles were either full text or just the title and abstract. Some criteria followed to search the articles included Probiotics, Sources of probiotics, Mechanism of action of probiotics, Probiotics in human health,

Role of probiotics in different diseases (probiotics and allergy, probiotics and diabetes, probiotics and cancer, probiotics and probiotics and cardiovascular diseases, probiotics and diarrhoea, probiotics and anaemia, probiotics and ulcers, probiotics and obesity). The articles that met the requirements for inclusion in this review were either published in peer-reviewed journals, reviewed or authored in English. Only studies about human health were included in the review. The exclusion criteria were as follows: no conference publications or book chapters, no mention of animal diseases that are host specific

like foot and mouth disease, no article searches in other languages and yeast or fungus strains not acting as probiotics.

### 3. ROLE OF PROBIOTICS IN VARIOUS DISEASES AND ABNORMALITIES OF THE HUMAN BODY

#### 3.1 Prevention of Allergies

Nowadays, Food allergies are becoming very common globally which lowers the quality of life for people who are impacted. People of all ages have been especially affected by food allergies. Nearly 6.0 % of children and 3.0 to 4.0 % of adults suffer from food allergies [60]. Mainly, food allergy is a consequence of several genetic and environmental variables, leading to an absence or reduction in tolerance of particular foods. Sometimes, Food-induced immune responses can result in various symptoms and conditions, such as anaphylaxis and severe allergic reactions [61]. So, avoiding allergens and taking emergency medicine is the only way to manage food allergies without a successful course of treatment [62]. Probiotics help allergies by healing the digestive tract [63]. As a result of this intervention, the immune system stabilizes, the intestinal epithelial barrier is strengthened and inflammation is reduced. Immunological processes cause hypersensitivity reactions which are the outward manifestation of allergies [64]. Probiotics can rearrange antigens, lessening their capacity to elicit an immunological response [64]. This ultimately results in a reduction of intestinal permeability and the class of pro-inflammatory compounds. People who suffer from various allergies often experience these symptoms. Research indicates that *L. rhamnosus* GG, a probiotic bacteria, may lessen the intensity of allergic reactions and ease the symptoms associated with food allergies [63,64]. According to a more contemporary hygiene hypothesis, certain organisms such as *helminths* and microorganisms such as *bifidobacteria*, *lactobacilli* and *saprophytic mycobacteria* are necessary for the healthy development of the human immune system. To support immunological tolerance, these microorganisms are believed to have coevolved with the human immune system as part of our microbiota during human evolution. This involves the production of regulatory T cells (Treg) and the regulation of the balance between Th2 and Th1 which helps avoid the development of autoimmune and allergic diseases [65,66]. A study was conducted that discovered that supplementing with *L. casei*

CRL431 and *B. lactis* Bb-12 for twelve (12.0) months did not influence the development of tolerance in 119 infants who were allergic to cow's milk [60].

#### 3.2 Antimicrobial Effect

One possible strategy to stop the spread of antibiotic-resistant bacteria is using probiotics [67]. Probiotics use several important antimicrobial pathways to accomplish this, including immune system modulation, mucin and tight junction protein production, competitive exclusion, and improved intestinal barrier function [68]. Probiotics produce antimicrobial peptides essential for competitively excluding microorganisms and are a major factor in their antimicrobial effectiveness. The antibacterial activity of *Lactobacillus* strains such as *S. aureus*, *P. aeruginosa*, *Shigella sp.*, *Clostridium difficile*, *Klebsiella sp.*, and *E. coli* has been particularly effective against a range of infections [69]. They produce lactic acid, bacteriocin, and hydrogen peroxide which prevent the attachment of harmful bacteria to the mucosa and enhance the immune response, among other efforts to outcompete and suppress these pathogenic bacteria [70]. The release of chemicals known as bacteriocins is the main way that *Lactobacillus* strains demonstrate antibacterial activity [71]. Antimicrobial peptides known as bacteriocins fight both Gram-positive and Gram-negative bacteria. Interestingly, the bacteria that make these bacteriocins have evolved defence mechanisms against the effects of their antimicrobial peptides [72]. It is also known that certain strains of *Lactobacillus* and *Bifidobacterium* produce bacteriocins. The antimicrobial peptides exhibit multiple modes of action such as blocking the creation of peptidoglycan and pore formation and inhibiting lipid II, an essential constituent of bacterial cell membranes [73]. A study showed that by restoring cell viability and regulating the production of inflammatory cytokines, *B. longum* and *L. reuteri* provide protective benefits in corneal infection induced by *P. aeruginosa* [74].

#### 3.3 Prevention of Diarrhoea

Numerous clinical investigations have evaluated the probiotics' effectiveness in preventing acute diarrhoeal illness [75]. The prevalent side effect of repeated short- and long-term antibiotic usage is diarrhoea [75]. To stop antibiotic-associated diarrhea (AAD), two meta-analyses utilizing the strains *Lactobacillus acidophilus*, *Lactobacillus*

*rhamnosus* strain GG, *Lactobacillus bulgaricus* and yeast *Saccharomyces boulardii* could be employed [75]. A study was conducted in patients with spinal cord injury (SCI) who frequently took proton pump inhibitors (PPI) in 7.0 days. This study aimed to evaluate the effectiveness of a probiotic containing at least  $6.5 \times 10^9$  live *Lactobacillus casei* Shirota (LcS) in reducing antibiotic-associated diarrhoea (AAD). When LcS is administered regularly to SCI patients who fall into a specific vulnerable group, it may be able to prevent AAD [76]. Another study was conducted to assess *Lactobacillus rhamnosus* GG (LGG) effectiveness in preventing antibiotic-associated diarrhoea (AAD) in adults and children (1499 participants). This finding indicates that in children and adults receiving antibiotic treatment for any cause, *Lactobacillus rhamnosus* GG is useful in avoiding antibiotic-associated diarrhoea [77]. Thus, nosocomial diarrhoea is a major issue in pediatric hospitals and is a global problem. Probiotics (*Streptococcus thermophiles*, *Bifidobacterium bifidum*, and *L. rhamnosus* starin GG) can be used prophylactically to prevent acute diarrhoea, especially in babies [75]. A study was also conducted in children aged six (6.0) months to five (5.0) years with watery diarrhoea in five (5.0) days. The result shows that probiotics are more effective in treating acute watery diarrhoea in hospitalized children. Probiotics may help lessen the intensity and length of diarrhoea when taken in addition to conventional treatment [78].

### 3.4 Prevention of Obesity

Human studies that examined the benefits of probiotic use included changes in lipid profiles, namely lower levels of total cholesterol, LDL cholesterol and plasma TG and higher levels of HDL cholesterol [79]. During 4.0 to 6.0 weeks of intake, the majority of research indicates the anti-obesity advantages of some types of probiotics such as *Lactobacillus* (primarily *Lactobacillus casei* strain Shirota (LAB13), *Lactobacillus gasseri*, *Lactobacillus rhamnosus* and *Lactobacillus plantarum*) and *Bifidobacterium* (particularly *Bifidobacterium infantis*, *Bifidobacterium longum*, and *Bifidobacterium breve* B3) [80,81]. Additionally, probiotics with anti-obesogenic properties, including *L. rhamnosus* CGMCC1.3724, help prevent obesity-related issues [82]. A study was

conducted to examine how the probiotic *Lactobacillus gasseri* SBT2055 (LG2055) affected body weight, other body measurements and abdominal adiposity in persons who tended toward obesity while consuming 200.0 g/day of Fermented milk for 12.0 weeks. The result demonstrated that the probiotic LG2055 decreased the effects on body weight, abdominal obesity and other metrics [83]. Another study was conducted with 45.0 patients with obesity, separated into three groups: probiotics (30.0 g/day), prebiotics (30.0 g/day) and single diets (low-carb and energy-reduced diets with *Bifidobacterium longum*, *Lactobacillus helveticus*, *Lactococcus lactis* and *Streptococcus thermophilus* per day). There is clinical evidence that the prebiotic and probiotic groups showed a significant decrease in fat mass, increased muscle strength, and reduced fasting blood glucose [84]. The other study was carried out with 66 .0 obese patients for three (3.0) months with a BMI in the range of 30.0 to 40.0 kg/m<sup>2</sup> in obese adults. The result shows that total cholesterol, triglycerides and low-density lipoprotein levels were decreased [85].

### 3.5 Prevention of Diabetes

Patients with type 2 diabetes may have changed gut microbial composition. This implies that altering the gut microbiota by probiotic medication could be a strategy for controlling glucose metabolism. Probiotics may have a hypoglycemic effect, according to numerous studies but the exact processes behind this are still up for debate because the aetiology of type 2 diabetes is so complicated [86]. Probiotics may help to mitigate several metabolic syndrome components, many of which are connected to type 2 diabetes [87]. The preservation of a range of mucosal immune cells may inhibit potential pathogens; pro-inflammatory cytokines (TNF- $\alpha$  and interleukin-6) may be reduced; delayed glucose absorption in the gut may contribute to these improvements; and rate-controlling enzymes of gluconeogenesis, such as glucose-6-phosphatase (G-6-Pase) and phosphoenolpyruvate carboxykinase (PEPCK), may be suppressed. In patients with type 2 diabetes, unchecked gluconeogenesis results in hyperglycemia [156]. Thus, probiotic therapy that inhibits G6Pase and PEPCK lowers glycemia by reducing the amount of glucose released into the bloodstream from the liver [88].



**Table 2. Overview of important studies demonstrating the effect of probiotics on type 2 diabetes mellitus.**

Probiotics	Model Types	Time of Experiment	Finding	References
<i>Lactobacillus brevis</i> 15 and <i>Lactobacillus plantarum</i> 13	24, 2.0-month-old Wistar rat	8.0 weeks	According to this study, the recently identified Bulgarian <i>Lactobacillus</i> strains <i>L. brevis</i> 15 and <i>L. plantarum</i> 13 may be potential probiotics that might help avoid certain metabolic disorders and diabetes-related difficulties.	[89]
Symbiter	53, patients	8.0 weeks	A little improvement in insulin resistance.	[90]
<i>L. rhamnosus</i> BSL and <i>L. rhamnosus</i> R23	Male Sprague-Dawley rats (n = 24)	1.0 months	In a streptozotocin-induced diabetic rat model, the liver showed better gene expressions and reduced fasting blood glucose after receiving 30 days' worth of <i>L. rhamnosus</i> BSL and <i>L. rhamnosus</i> R23 dose, reduction of glucose-6-phosphatase to increase insulin sensitivity.	[91]
<i>Bifidobacterium animalis</i> A6	28, type II diabetic patients	4.0 weeks	Patients with type 2 diabetes can greatly lower their fasting blood glucose by taking probiotic camel milk powder twice a day for four (4) weeks in a row.	[92]
Ecologic Barrier®	patients with type 2 diabetes mellitus	6.0 months	Multistrain probiotics show promise as an adjunct treatment for diabetes.	[93]

**Table 3. Different studies on the treatment of anaemia using probiotics.**

Subject	Number of Patients	Study Duration	Observation	References
Healthy woman	Study 1- 14; Study 2- 28	4.0 weeks	In the probiotics group: significant increase of iron in serum; In the placebo group: Changes are insignificant.	[97]
Healthy woman	20	1.0 week	Probiotics Group: Significant increase in iron.	[98]
Children with anaemia	52	3.0 years	No significant differences in serum ferritin levels between the probiotics and control group.	[99]

### 3.6 Prevention of Anemia

Numerous researches have demonstrated a strong correlation between iron deficiency and gut microbiota. Moreover, studies have demonstrated the positive effect of probiotic bacteria on improving iron absorption and the reciprocal association between gut microbiota composition and iron insufficiency [94,95], Probiotic bacteria such as *Lactobacillus*

*cerevisiae*, *Lactobacillus cremoris*, *B. pseudocatenulatum*, *Candida famata*, *B. adolescentis*, *Candida glabrata*, *Candida guilliermondii*, *S. cerevisiae*, *Yarrowia lipolytica* and *Pichia glucozyma* are the source of folic acid, a water-soluble B vitamin that is essential for treating and managing anemia. These microbes are used to improve folic acid absorption through the intestines. *P. denitrificans* and *P. shermanii* have also been discovered to

be effective in treating vitamin B12 insufficiency. Foods fermented with lactic acid have a crucial role in boosting the absorption of iron and are utilized in the treatment of anaemia in patients. They aid in regulating the digestive tract's pH and stimulate the phytase enzyme which aids in the absorption of nutrients. Moreover, megaloblastic anaemia has been treated by adding a mix of probiotic bacteria to the diet. These probiotics mitigate the negative effects of antibiotics and encourage intestinal fermentation [96].

### 3.7 Prevention of Urogenital Disease

According to recent research, maintaining a lactobacilli-dominated microbiome is crucial for preserving women's quality of life and preventing STDs and premature labour. It was questioned if artificial lactobacilli supplementation may reduce infection rates because women who are prone to vaginal and urinary infections lose lactobacilli organisms. However, uro-pathogens are becoming more resistant to common antibiotics such as trimethoprim/sulfamethoxazole and patients are turning increasingly to natural alternatives like cranberry juice which seems to contain antiadhesive compounds that are effective against uro-pathogens and can help prevent UTIs. Probiotic therapy using lactobacilli has been explored as a non-chemotherapeutic way to maintain and repair a healthy urogenital tract. There is evidence that some strains of lactobacilli can be beneficial when inserted directly into the vagina or when ascending from the rectum after oral ingestion. Probiotic consumption regularly may be beneficial for all healthy women as well as those who experience recurrent urogenital infections to prevent serious infections and superinfections of the vaginal mucosa. Probiotics have been shown to help keep the vaginal flora in balance which protects against UTIs [100]. A study was conducted with 174 premenopausal women with a history of recurrent UTIs at four (4.0) months and examined the efficacy of oral and intravaginal probiotic supplementation as a preventative measure against recurrent urinary tract infections. The result demonstrated the effectiveness in preventing recurrent symptomatic UTI episodes [101]. Another study was conducted with a population of 132 individuals with probiotics, antibiotics and probiotics plus antibiotics being administered to 42, 48 and 42 of the participants in each group at 30.0 days. This study found that probiotics are a safe and efficient way to treat UTIs in Bangladeshi women [102].

### 3.8 Prevention of Dental Caries

Dental caries is one of the most prevalent illnesses in the world, causing exorbitant expenses and being regarded as a public health issue in youngsters. Various therapies have been suggested to lower the prevalence of this illness, with probiotic supplementation being one of the main ones. According to recent studies, probiotic supplementation appears to lower the risk of dental caries [103-105]. *Streptococcus mutans* (SM), one of the several bacterial species found in saliva, is widely regarded as the pathogen primarily responsible for the early stages of tooth decay, whereas *Lactobacillus* plays a significant part in the disease's progression. The primary traits of *Streptococcus mutans*' pathogenicity include its acidogenicity, capacity to endure in acidic conditions, capacity to form biofilms, and capacity to cling to teeth. The probiotic *Lactobacillus paracasei* SD1 was introduced and used in the oral cavity due to its well-known ability to prevent *Streptococcus mutans* from growing, lessen its acid production, and adhere well to oral epithelial cells [106]. A study involved 260 participants of which 130 were in the control group and 130 in the intervention group at 4.0, 8.0 and 12.0 months. The group that took probiotic milk tablets demonstrated statistically significant decreases in carious lesions when compared to the control group [107].

### 3.9 Prevention of Depression

The gut-brain axis is the term for the reciprocal connections between the brain, intestinal microbiota and the digestive system via immunological, neurological and endocrine pathways. The gut microbiota responds to the physical and physiological strain its host endures. Stress drastically reduces the amount of Firmicutes phylum, particularly in *Bifidobacterium* and *Lactobacillaceae* family bacteria and is a major risk factor for depression. It also modifies the percentage and makeup of the gut microbiota. Stress weakens the brain's relationship with the gut microbiota which can result in metabolic depression. However, Changes in brain activity in patients with depression are known to be caused by higher levels of cortisol in plasma and corticotrophin-releasing factor in the cerebrospinal fluid. Furthermore, catecholamine levels (primarily those of norepinephrine and dopamine) rise when these factors are repeatedly activated. This raises the risk of various physiological problems

such as attention deficit disorder, balance and posture, cognitive function, emotions and sleep-wake cycles. *Bifidobacterium* baby restored the stress-induced drop in norepinephrine, indicating that probiotics influenced catecholamine metabolism [108]. A clinical trial consisted of 45 patients with *Bifidobacterium breve* CCFM1025 in 4.0 weeks to examine the psychotropic potential for the treatment of major depressive disorder (MDD). This trial shows that *B. breve* CCFM1025 is a good candidate that reduces depression and related gastrointestinal issues [109].

### 3.10 Prevention of Cardiovascular Diseases

Cardiovascular problems commonly affecting the heart include arrhythmia, thromboembolic disease, cardiomyopathy and atherosclerosis. Globally, the prevalence of cardiovascular disorders is rising. Cardiovascular diseases accounted for 18.0 million fatalities in 2015 or over one-third of all deaths with documented causes. This number indicates a noteworthy increase of 12.50 % when compared to the data from 2005. Predictions from the American Heart Association suggest that by 2030, around 43.90 % of the US population will be suffering from cardiovascular disease. In the United States, 92.10 million persons currently suffer from cardiovascular disease. Atherosclerosis is a chronic inflammatory disease characterized by the creation and progression of lipid-fueled atherosclerotic plaques in the artery wall [110]. From spontaneously fermented mustard, researchers obtained the *Lactobacillus plantarum* DMDL 9010 strain (CGMCC No. 5172). The potential benefits of this strain in lowering cholesterol were examined using an in vivo model. The study's findings demonstrated that rats given 109 cells of *L. plantarum* DMDL 9010 daily experienced significant reductions in their levels of total cholesterol (TC), low-density lipoprotein cholesterol content (LDL-C), and atherosclerosis index (AI) of 23.03 %, 28.00 %, and 34.03 %, respectively. After administering *L. plantarum* DMDL 9010, there was no statistically significant alteration in blood triglyceride levels. Rat hepatocyte protection against steatosis was demonstrated by morphological and pathological alterations in the liver following *L. plantarum* DMDL 9010 therapy. Significant reductions in hepatic cholesterol (33.20 %) and triglyceride levels (40.86 %) were also observed upon administration of substantial dosages of *L. plantarum* DMDL 9010. Significant increases

were also observed in bile acid (+ 70.18 %) and faecal cholesterol (+ 31.07 %). After receiving *L. plantarum* DMDL 9010 treatment, there was a reduction in blood and total hepatic cholesterol and triglyceride levels and increased bile acid excretion through stool. These outcomes seemed to depend on the dosage of *L. plantarum* DMDL 9010. Thus, Probiotic supplementation may help those who are susceptible to ischemic heart disease escape the devastation that comes with it [111].

### 3.11 Prevention of Upper Respiratory Tract Infections

The most prevalent illnesses are upper respiratory tract infections (URTIs), including tonsillitis, pharyngitis, laryngitis, sinusitis, acute respiratory distress known as coronavirus, and the common cold [112,113]. Adults experience colds two to three times a year, and 17.2 billion URTIs are thought to have occurred globally in 2015 [114,115]. Numerous viruses, including the influenza virus, adenovirus, rhinovirus and respiratory syncytial virus are the primary cause of URTIs. The symptoms can include chills, fever, headache, body aches, sore throat, runny or stuffy nose, cough and so on [116,117]. Retaining the immune system in a healthy, normal state reduces the likelihood of recurrent infections. Salivary immunoglobulin A (IgA) and NK cell activity are thought to be crucial in the prevention of URTIs. The immune system can be weakened by several environmental variables such as leading a stressful lifestyle which could raise the risk of URTIs. Evidence shows that certain probiotic strains increase salivary IgA levels and activate NK cells again. Therefore, regular use of a probiotic beverage may help to regulate URTIs and preserve normal immunological function. A probiotic known as *Lactobacillus casei* strain Shirota (LcS) can survive in the digestive system and restore a balanced gut flora. Numerous studies have been conducted on human and animal models to examine the strain's immunomodulatory properties. Research involving healthy individuals with low NK cell activity revealed that consumption of LcS-fermented milk (LcS-FM) increased the activity [118].

### 3.12 Prevention of Ulcers

The sores that form on the skin's surface that are painful and inflammatory are called ulcers. Clinical characteristics determine whether the illness is considered mild, substantial, or herpetiform. Many theories have been put out to

explain the illness, including immunologic abnormalities, viral infections, and environmental and psychological stress. However, the disorder's aetiology is still unknown and up for debate. Probiotics given in sufficient amounts have been shown to improve host health. One way probiotics help reduce sickness is by enhancing and modulating the immune system. Probiotics can produce a variety of substances, such as peptides, bacteriocins and organic acids. Consequently, there is a decreased chance of harmful germs colonizing [119]. Oral ulcers are a common oral ailment that is distinguished by a high degree of complexity, diversity and frequency. Several etiological factors such as allergic reactions, traumatic experiences and infectious agents, can result in oral ulcers. The hallmarks of mouth ulcers are the loss of connective tissue and the development of crater-like depressions brought on by persistent disruption or damage to the integrity of the mouth epithelium. Bacteria that are resistant to infections and benign are known as probiotics. In addition to strengthening immunity and preventing infection by fighting harmful bacteria, they may also improve digestion and make it easier for the host to absorb nutrients. They influence both immune system regulation and tissue healing. It may be possible to increase the effectiveness of probiotics in aiding the healing process of ulcers by using fucoidan (FD), a prebiotic obtained from marine sources. Probiotics can create a biofilm in the oral cavity that can successfully compete with periodontal and cancer-causing bacteria. They also help prevent diseases by enhancing immune system performance and modulating the host's immunological response. Dental health is significantly improved while using oral probiotics [120,121].

The development of peptic ulcers in the duodenum or stomach is known as peptic ulcer disease. The main risk factors for gastric and duodenal ulcers are the use of aspirin or nonsteroidal anti-inflammatory medicines (NSAIDs) and the existence of an *H. pylori* infection. The function of the barrier may be improved by probiotics. The primary defence against pathogenic bacteria is the acid and mucus barrier found in the stomach's gastric mucosa. Some probiotics can increase the upregulation of fitting linked proteins and promote the secretion of mucus and mucin which strengthens the mucus secretion process and improves the protective capacity of the stomach mucosal barrier. Furthermore, some probiotic

bacteria can create antimicrobial substances such as hydrogen peroxide, lactic acid, short-chain fatty acids (SCFAs) and bacteriocins. Short-chain fatty acids (SCFAs) and lactic acid both show signs of partial dissociation and when these organic acids are in their dissociated stages, *H. pylori* is negatively impacted [122,123].

### 3.13 Prevention of Breast Cancer

Breast cancer makes for 29.0 % of all cancer cases identified in women, making it the most frequent cancer among them. An estimated 2.50 million instances of breast cancer were reported in 2016 [124]. It was previously believed that disparities in hormonal components, reproductive patterns, and detection techniques at various stages were the causes of the wide variances in breast cancer incidence [125,126]. Breast cancer mortality has continuously dropped as a result of advancements in breast cancer detection and treatment. It was previously believed that probiotic supplements could lower the incidence of breast cancer in perimenopausal women. Bacteria believed to have anticancer properties include *Lactococcus* and *Streptococcus* which are more common in healthy breast tissue. To see whether they may cause DNA damage, bacteria from malignant females were cultivated on human breast cancer cells. Triple-stranded DNA breaks generated by *Enterobacteriaceae*, *Escherichia coli*, and *Staphylococcus* bacteria were alarmingly visible throughout the proliferation of these malignant cells. Carcinogens can be broken down by the probiotic bacterium *Lactobacillus*. The probiotic *S. thermophilus* strain produces antioxidants that reduce DNA damage [127]. Moreover, Research has indicated that *Lactobacillus crispatus* and *Lactobacillus acidophilus* exhibit antiproliferative properties against breast cancer cells. Furthermore, the transcriptional activity of four distinct cancer-testis antigens can be decreased by lactobacilli. It has been revealed that epigenetic controls have a role in the expression of cancer-testis antigens. Thus, its expression may be epigenetically downregulated by lactobacilli. A new era of clinical research applications could be enabled by lactobacilli-mediated expressional downregulation of cancer-testis antigens, as these antigens have been associated with high-grade malignancies and poor prognosis [128].

### 3.14 Prevention of Blood Cancer

Most hematopoietic malignancies, often known as blood cancers, start in the bone marrow,

which is where blood cells are made. Research indicates that probiotics may be useful in the treatment of cancer. A study found a correlation between chronic myeloid leukaemia and the therapeutic effects of heat-inactivated *Lactobacillus casei* and *Lactobacillus paracasei* on the K562 cell line. Bacteria were cultivated in an oxygen-free environment using a particular medium, and their cells were then exposed to a temperature of 100.0 °C to induce cell death. Lyophilization was used to extract moisture from the dead cells and autoclaving was used to sterilize them thereafter. The next step involved creating several heat-inactivated bacterial suspensions at different concentrations (125.0, 250.0, 500.0, 1000.0 and 2000.0 g/ml). Three distinct time points 24.0, 48.0 and 72.0 hours were used to evaluate the in vitro anticancer qualities of heat-killed bacteria using the MTT test. The results showed that *L. casei* and *L. paracasei* heat-killed cells have anticancer properties in K562 cells. Moreover, the anticancer activities of heat-killed cells are closely correlated with their concentration. In summary, heat-inactivated *Lactobacillus* has great potential and should be investigated further in the field of treatments for chronic myeloid leukaemia [64].

### 3.15 Prevention of Colorectal Cancer

One of the most prevalent cancers in the world and the primary cause of cancer-related mortality is colorectal cancer (CRC) [129]. The development of colorectal cancer and its incidence are intimately correlated with the gut microbiota. The altered gut microbiota can initiate carcinogenesis through modifications to the immune system, epithelial hemostasis, metabolic profile and activity, DNA damage and aberrant cellular and molecular activities in colonocytes. Numerous unfavourable treatment-associated side effects negatively impact the quality of life. Although multiple cutting-edge medical methods (such as immunotherapy, surgery, chemotherapy, and radiation therapy) are available for the treatment of colorectal cancer, the survival rates are poor [130,131]. So, one well-known bioactive option for treating several illnesses and disorders is probiotics [132,133]. A powerful probiotic strain can help prevent tumorigenesis, including colorectal cancer (CRC), by competing for adhesion sites, producing microbicidal agents like bacteriocin, improving intestinal permeability, releasing bioactive metabolites, regulating immune pathways and stimulating cell protective

responses [131]. Despite several attempts to elucidate the mechanism behind probiotics' anticarcinogenic properties, a definitive explanation of probiotics' anti-CRC action remains to be discovered [134,135].

The production of carcinogenic compounds such as cresols, ammonia, phenols, aglycones and N-nitroso compounds, is linked to certain bacterial enzymes, including nitrate reductase, azoreductase,  $\beta$ -glucosidase,  $\beta$ -glucuronidase and 7- $\alpha$ -dehydroxylase. These compounds also induce antiapoptotic pathways, which contribute to the development of colorectal cancer [136,137]. Research demonstrated that adding probiotics to the diet dramatically decreased the activity of bacterial enzymes [138-140]. The carcinogenic chemicals are released through faeces after binding with peptidoglycan, which is found in the probiotic microorganisms' cell walls. Certain probiotic strains can metabolize carcinogenic substances, particularly amines and N-nitroso substances. Probiotic supplementation has been shown to reduce the incidence of colorectal cancer (CRC) through a variety of mechanisms, including altered metabolic activity of intestinal microbiota (i.e., decreased endogenous generation of carcinogenic chemicals) and the binding and destruction of carcinogens [141].

### 3.16 Prevention of Liver Cancer

The second greatest death rate globally is from primary liver cancer. There are various types of malignant tumor in this condition which can include fibrolamellar HCC (FLC), pediatric neoplasm hepatoblastoma, mixed hepatocellular cholangiocarcinoma (HCC-CAA [142]), intrahepatic cholangiocarcinoma (iCAA) and HCC. Of all primary liver malignancies, HCC accounts for 75.0 to 85.0 % of cases [142,143]. The more aggressive proliferative variant of HCC is linked to elevated serum levels of  $\alpha$ -fetoprotein (AFP), progenitor cell phenotype expression, mutations in tumor protein 53 (TP53) and activation of the transforming growth factor  $\beta$  (TGF- $\beta$ ), hepatocyte growth factor receptor (MET), protein kinase B (AKT), and insulin-like growth factor (IGF) 2 pathways [144]. The prognosis for HCC patients is still not good generally, despite advances in treatment such as immune therapy, multi-kinase inhibitors, and intra-arterial therapy [145]. Pathogenesis of HCC is connected to adverse changes in the gut microbiota because the hepatic portal circulation connects the liver and the intestines directly

[146]. According to this anatomical link, probiotic bacteria may be able to reverse the dysbiosis linked with HCC by reestablishing the complexity and colonization resistance of the gut flora [147]. The expression of tight junction proteins is upregulated by probiotic metabolites and probiotic cell components, which improve gut epithelial integrity. The integrity of the gut epithelium can be preserved by the probiotic bacteria's cell surface proteins which can reduce gut epithelial cell inflammation and prevent epithelial cell apoptosis. Probiotics also shield the gut epithelium from harmful bacteria by increasing goblet cells' mucus production and antimicrobial peptides [148]. Probiotic bacteria, particularly *Bifidobacterium spp.* and *Lactobacillus spp.*, can lessen fatty liver, lowering the risk of HCC [149].

### 3.17 Prevention of Lung Cancer

Lung cancer is the second most frequent type of cancer and the primary cause of cancer-related mortality [150]. According to estimates, smoking is the primary cause of 90.0 % of lung cancer occurrences and other known risk factors include tobacco smoke, air pollution and other toxins. The precise mechanics, nevertheless, are not fully understood. Lungs offer a special chance to be exposed to germs and environmental hazards since they are the mucosa site with the biggest surface area in the body and an important interaction with the outside world. Despite being believed to be sterile, many microorganisms reside in the lungs. Introducing new bacteria, their removal by the body and immune system, and their ability to reproduce in the given environment all influence the prevalence of germs. Long-term obstructive pulmonary disease and cystic fibrosis are two lung conditions that seem to have dysregulated lung microbiome [151-154]. One research topic is the connection between immunological response to medication and gut flora. Recent research indicates that the gut microbiota's composition may influence how effective immunotherapy is [155]. Probiotics may increase the effectiveness of immunotherapy and help to overcome some of its drawbacks by enhancing gut bacteria diversity and fostering a supportive immunological milieu [7,154-156].

### 4. FUTURE OF PROBIOTICS

The future of probiotic research is promising and multifaceted with numerous important areas poised for significant advancement. Though

antibiotic resistance is very common nowadays, so probiotics can be replaceable treatments for the future. Identifying and utilizing probiotics from diverse and sustainable sources, including soil and marine environments, can provide new strains with unique benefits. This approach also supports biodiversity and environmental sustainability. Innovations in delivery mechanisms such as encapsulation technologies and smart delivery systems, can improve the survival and efficacy of probiotics as they pass through the digestive system, ensuring they reach their target site in the gut. Advances in genetic engineering could enable the creation of genetically modified probiotics with enhanced functionality such as targeted delivery of therapeutic compounds or improved survival and colonization in the gut. Further research into the precise mechanisms of action of different probiotic strains can provide deeper insights into their role in health and disease, facilitating more targeted and effective applications.

### 5. CONCLUSION

Probiotics are an exciting and rapidly developing field of study that significantly affects human health. These good bacteria play a major role in preserving and improving the balance of the gut microbiota. They are primarily obtained via fermented foods and nutritional supplements. Their potential for therapeutic use is highlighted by their advantageous activities which include enhancing the intestinal barrier, regulating the immunological response and generating vital metabolites. Probiotics have been shown to help treat various illnesses, including mental health problems, systemic diseases like diabetes and gastrointestinal ailments. Although the efficacy of probiotics varies greatly depending on the strain and the individual's health status, the isolation and characterization of particular probiotic strains have permitted focused therapeutic uses. The growing understanding of the gut microbiome's critical role in general health fuels the field's expansion. More studies are necessary to comprehend and maximize the use of probiotics in clinical settings completely.

### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of manuscripts.



## ACKNOWLEDGEMENT

We would like to express our deepest gratitude to all the researchers whose work on probiotics has significantly contributed to the advancement of knowledge in this field. We have cited as many references as permitted and apologize to the authors of those publications that we have not cited due to limitation of references. We apologize to other authors who have worked on several aspects of probiotics but whom we have unintentionally overlooked. Finally, we would like to extend our heartfelt thanks to the Chairman of the Department of Pharmacy, and the Chairman of the Department of Biotechnology & Genetic Engineering, Islamic University, Kushtia-7003, Bangladesh.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Mohr KI. History of antibiotics research. *Curr. Top. Microbiol. Immunol.* 2016;398: 237–272.  
DOI: 10.1007/82\_2016\_499
2. Blondeau LD, Blondeau JM. Antimicrobial Resistance. *Diagnostics Ther. Vet. Dermatology.* 2021;163–174.  
DOI: 10.1002/9781119680642.ch16
3. Salahi A, Abd El-Ghany WA. Beyond probiotics, uses of their next-generation for poultry and humans: A review. *J. Anim. Physiol. Anim. Nutr. (Berl).* 2024;1–12.  
DOI: 10.1111/jpn.13972
4. Wieggers C, Van Beek EHT, Larsen OFA. Clinical research with probiotics as an indicator of global valorization since the year 2000. *Front. Microbiol.* 2023;14 :1–9.  
DOI: 10.3389/fmicb.2023.1323920
5. Mackowiak PA. Recycling Metchnikoff: Probiotics, the intestinal microbiome and the quest for long life. *Front. Public Heal.* 2013;1:1–3.  
DOI: 10.3389/fpubh.2013.00052
6. Morelli L, Capurso L. FAO/WHO Guidelines on Probiotics: 10 Years Later. *J. Clin. Gastroenterol.* 2012;46.  
Available: [https://journals.lww.com/jcge/fulltext/2012/10001/fao\\_who\\_guidelines\\_on\\_probiotics\\_\\_10\\_years\\_later.2.aspx](https://journals.lww.com/jcge/fulltext/2012/10001/fao_who_guidelines_on_probiotics__10_years_later.2.aspx)
7. Hill C. et al. Expert consensus document: The international scientific association for probiotics and prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat. Rev. Gastroenterol. Hepatol.* 2014;11(8):506–514.  
DOI: 10.1038/nrgastro.2014.66
8. Adel M, El-Sayed AFM, Yeganeh S, Dadar M, Giri SS. Effect of potential probiotic *Lactococcus lactis* Subsp. *lactis* on Growth Performance, Intestinal Microbiota, Digestive Enzyme Activities, and Disease Resistance of *Litopenaeus vannamei*. *Probiotics Antimicrob. Proteins.* 2017;9(2): 150–156.  
DOI: 10.1007/s12602-016-9235-9
9. Azcárate-Peril MA, Sikes M, Bruno-Bárcena JM. The intestinal microbiota, gastrointestinal environment and colorectal cancer: A putative role for probiotics in prevention of colorectal cancer? *Am. J. Physiol. - Gastrointest. Liver Physiol.* 2011;301(3).  
DOI: 10.1152/ajpgi.00110.2011.
10. Maldonado Galdeano C, Cazorla SI, Lemme Dumit JM, Vélez E, Perdígón G. Beneficial effects of probiotic consumption on the immune system. *Ann. Nutr. Metab.* 2019;74(2):115–124.  
DOI: 10.1159/000496426
11. De Melo Pereira GV, De Oliveira Coelho B, Magalhães Júnior AI, Thomaz-Soccol V, Soccol CR. How to select a probiotic? A review and update of methods and criteria. *Biotechnol. Adv.* 2018;36(8):2060–2076.  
Available: <https://doi.org/10.1016/j.biotechadv.2018.09.003>.
12. Plaza-Díaz J, et al. Gene expression profiling in the intestinal mucosa of obese rats administered probiotic bacteria. *Sci. Data.* 2017;4(1):170186.  
DOI: 10.1038/sdata.2017.186
13. Collado MC, Gueimonde M, Salminen S. Chapter 23 - Probiotics in Adhesion of Pathogens: Mechanisms of Action. Watson RR, VRB TBF. in P. H. Preedy, Eds., Boston: Academic Press. 2010;353–370.  
Available: <https://doi.org/10.1016/B978-0-12-374938-3.00023-2>.
14. Hemaiswarya S, Raja R, Ravikumar R, Carvalho IS. Mechanism of action of probiotics. *Brazilian Arch. Biol. Technol.* 2013;56(1):113–119.  
DOI: 10.1590/S1516-89132013000100015
15. Trush EA, Poluektova EA, Beniashvili AG, Shifrin OS, Poluektov YM, Ivashkin VT. The evolution of human probiotics: Challenges and prospects. *Probiotics*

- Antimicrob. Proteins. 2020;12(4):1291–1299.  
DOI: 10.1007/s12602-019-09628-4
16. Gram-positive bacteria with probiotic potential for the *Apis mellifera L.* honey bee: The experience in the northwest of Argentina. *Probiotics and antimicrobial proteins*; 2022.
  17. Reid G, Burton J. Use of *Lactobacillus* to prevent infection by pathogenic bacteria. *Microbes Infect.* 2002;4(3):319–324.  
DOI: 10.1016/S1286-4579(02)01544-7
  18. Yunus RA, et al. A Multi-strain Potential Probiotic Formulation of GABA-Producing *Lactobacillus plantarum* 90sk and *Bifidobacterium adolescentis* 150 with Antidepressant Effects. *Probiotics Antimicrob. Proteins.* 2020;12(3):973–979.  
DOI: 10.1007/s12602-019-09601-1
  19. Fijan S, Ter Haar JA, Varga L. Probiotic microorganisms and their benefit to human health. *Adv. Probiotics.* 2021;3–22.  
DOI: 10.1016/b978-0-12-822909-5.00001-0
  20. Mazziotta C, Tognon M, Martini F, Torreggiani E, Rotondo JC. Probiotics Mechanism of Action on Immune Cells and Beneficial Effects on Human Health. 2023;12(1).  
DOI: 10.3390/cells12010184
  21. Fijan S. Microorganisms with claimed probiotic properties: An overview of recent literature. *Int. J. Environ. Res. Public Health.* 2014;11(5):4745–4767.  
DOI: 10.3390/ijerph110504745
  22. Fijan S. Probiotics and their antimicrobial effect. *Microorganisms.* 2023;11(2):17–20.  
DOI: 10.3390/microorganisms11020528
  23. Karimi S, Azizi F, Nayeb-Aghaee M, Mahmoodnia L. The antimicrobial activity of probiotic bacteria *Escherichia coli* isolated from different natural sources against hemorrhagic *E. coli* O157:H7. *Electron. Physician.* 2018;10(3):6548–6553.  
DOI: 10.19082/6548
  24. Karimi S, Rashidian E, Birjandi M, Mahmoodnia L. Antagonistic effect of isolated probiotic bacteria from natural sources against intestinal *Escherichia coli* pathotypes. *Electron. Physician.* 2018;10(3):6534–6539.  
DOI: 10.19082/6534
  25. Yuksel N, Gelmez B, Yildiz-Pekoz A. Lung Microbiota: Its relationship to respiratory system diseases and approaches for lung-targeted probiotic bacteria delivery. *Mol. Pharm.* 2023;20(7):3320–3337.  
DOI: 10.1021/acs.molpharmaceut.3c00323
  26. Zhai Q, Feng S, Arjan N, Chen W. A next generation probiotic, *Akkermansia muciniphila*. *Crit. Rev. Food Sci. Nutr.* 2019;59(19):3227–3236.  
DOI: 10.1080/10408398.2018.1517725
  27. Barbosa JC, et al. Chapter 24 - Next-generation probiotics. A. B. T.-P. Brandelli, Ed., Academic Press. 2022;483–502.  
Available: <https://doi.org/10.1016/B978-0-323-85170-1.00012-9>.
  28. O'Toole PW, Marchesi JR, Hill C. Next-generation probiotics: The spectrum from probiotics to live biotherapeutics. *Nat. Microbiol.* 2017;2(5):17057.  
DOI: 10.1038/nmicrobiol.2017.57
  29. Tan H, Zhai Q, Chen W. Investigations of *Bacteroides* spp. towards next-generation probiotics. *Food Res. Int.* 2019;116:637–644.  
Available: <https://doi.org/10.1016/j.foodres.2018.08.088>.
  30. Article R. *Pakistan Veterinary Journal.* 2012;8318:303–308.
  31. Latif A, et al. Probiotics: Mechanism of action, health benefits and their application in food industries. *Front. Microbiol.* 2023;14.  
DOI: 10.3389/fmicb.2023.1216674
  32. Ohland CL, MacNaughton WK. Probiotic bacteria and intestinal epithelial barrier function. *Am. J. Physiol. - Gastrointest. Liver Physiol.* 2010;298(6).  
DOI: 10.1152/ajpgi.00243.2009.
  33. Bermudez-Brito M, Plaza-Díaz J, Muñoz-Quezada S, Gómez-Llrente C, Gil A. Probiotic mechanisms of action. *Ann. Nutr. Metab.* 2012;61(2):160–174.  
DOI: 10.1159/000342079
  34. Van Tassell ML, Miller MJ. *Lactobacillus* adhesion to mucus. *Nutrients.* 2011;3(5):613–636.  
DOI: 10.3390/nu3050613
  35. Petruzzello C, Saviano A, Ojetti V. Probiotics, the immune response and acute appendicitis: A review. *Vaccines.* 2023;11(7).  
DOI: 10.3390/vaccines11071170
  36. Sajedi D, Shabani R, Elmieh A. Changes in leptin, serotonin, and cortisol after eight weeks of aerobic exercise with probiotic intake in a cuprizone-induced demyelination mouse model of multiple sclerosis. *Cytokine.* 2021;144:155590.

- Available:<https://doi.org/10.1016/j.cyto.2021.155590>.
37. Gangaraju D, Raghu AV, Siddalingaiya Gurudutt P. Green synthesis of  $\gamma$ -aminobutyric acid using permeabilized probiotic *Enterococcus faecium* for biocatalytic application. *Nano Sel.* 2022;3(10):1436–1447. DOI: 10.1002/nano.202200059
  38. Srivastav S, et al. Probiotics mixture increases butyrate, and subsequently rescues the nigral dopaminergic neurons from MPTP and rotenone-induced neurotoxicity. *J. Nutr. Biochem.* 2019;69:73–86. Available:<https://doi.org/10.1016/j.jnutbio.2019.03.021>.
  39. Quintieri L, Fanelli F, Monaci L, Fusco V. Milk and its derivatives as sources of components and microorganisms with health-promoting properties: Probiotics and bioactive peptides. *Foods.* 2024;13(4). DOI: 10.3390/foods13040601
  40. Li T, Teng D, Mao R, Hao Y, Wang X, Wang J. A critical review of antibiotic resistance in probiotic bacteria. *Food Res. Int.* 2020;136:109571. Available:<https://doi.org/10.1016/j.foodres.2020.109571>.
  41. Zommiti M, Feuilloley MGJ, Connil N. Update of probiotics in human world: A nonstop source of benefactions till the end of time. *Microorganisms.* 2020;8(12):1–33. DOI: 10.3390/microorganisms8121907
  42. Fusco V, Quero GM. Culture-dependent and culture-independent nucleic-acid-based methods used in the microbial safety assessment of milk and dairy products. *Compr. Rev. food Sci. food Saf.* 2014;13(4):493–537. DOI: 10.1111/1541-4337.12074
  43. Fontana L, Bermudez-Brito M, Plaza- Diaz J, Muñoz-Quezada S, Gil A. Sources, isolation, characterisation and evaluation of probiotics. *Br. J. Nutr.* 2013;109(S2):S35–S50. DOI: 10.1017/S0007114512004011
  44. Cunningham M, et al. Shaping the future of probiotics and prebiotics. *Trends Microbiol.* 2021;29(8):667–685. DOI: 10.1016/j.tim.2021.01.003
  45. Wang S, et al. Nonbovine milk and its products as sources of probiotics delivery: An overview of its viability, functionality and product quality characteristics. *Int. J. Dairy Technol.* 2023;76(3):482–511. DOI: 10.1111/1471-0307.12976
  46. Baranenko D, et al. Probiotic Emulsified Meat Products BT - Functional Meat Products. Verruck S, Teixeira Marsico E. Eds., New York, NY: Springer US. 2024;13–24. DOI: 10.1007/978-1-0716-3573-5\_2
  47. Sayes C, Leyton Y, Riquelme C. Probiotic Bacteria as an Healthy Alternative for Fish Aquaculture. Savic S, Ed., Rijeka: IntechOpen. 2017;Ch. 7. DOI: 10.5772/intechopen.71206
  48. Pourjafar H, Pimentel TC, Baú TR. Probiotic Fermented Vegetables BT - Probiotic Foods and Beverages: Technologies and Protocols. Gomes da Cruz A, Silva MC, Colombo Pimentel T, Esmerino EA, Verruck S, Eds., New York, NY: Springer US. 2023;119–132. DOI: 10.1007/978-1-0716-3187-4\_8
  49. Bernal-Castro C, Espinosa-Poveda E, Gutiérrez-Cortés C, Díaz-Moreno C. Vegetable substrates as an alternative for the inclusion of lactic acid bacteria with probiotic potential in food matrices. *J. Food Sci. Technol.* 2024;61(5):833–846. DOI: 10.1007/s13197-023-05779-z
  50. Rahman MS, et al. Recent advances in probiotication of fruit and vegetable juices. *J. Adv. Vet. Anim. Res.* 2023;10(3):522–537. DOI: 10.5455/javar.2023.j706
  51. Bodke H, Jogdand S. Role of probiotics in human health. *Cureus.* 2022;14(11):e31313. DOI: 10.7759/cureus.3131
  52. Ricci A, et al. Scientific Opinion on the update of the list of QPS-recommended biological agents intentionally added to food or feed as notified to EFSA. *EFSA J.* 2017;15(3). DOI: 10.2903/J.EFSA.2017.4664
  53. The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2016. *EFSA journal. Eur. Food Saf. Auth.* 2017;15(12):e05077. DOI: 10.2903/j.efsa.2017.5077
  54. Sionek B, Szydłowska A, Zielińska D, Neffe-Skocińska K, Kołożyn-Krajewska D. beneficial bacteria isolated from food in relation to the next generation of probiotics. *Microorganisms.* 2023;11(7):1–21. DOI: 10.3390/microorganisms11071714
  55. Singh K, Rao A. Probiotics: A potential

- immunomodulator in COVID-19 infection management. *Nutr. Res.* 2021;87:1–12. Available:<https://doi.org/10.1016/j.nutres.2020.12.014>.
56. Bunešová V, et al. Bifidobacterium animalis subsp. lactis strains isolated from dog faeces. *Vet. Microbiol.* 2012;160(3):501–505. Available:<https://doi.org/10.1016/j.vetmic.2012.06.005>.
  57. Abbasi Z, Ghasemi SM, Ahmadi Y, Shokri D. Isolation and identification of effective probiotics on drug-resistant acinetobacter baumannii strains and their biofilms. *Can. J. Infect. Dis. Med. Microbiol.* 2024;2024. DOI: 10.1155/2024/8570521
  58. SEA, et al. Isolation of Lactobacillus salivarius 1077 (NRRL B-50053) and Characterization of Its Bacteriocin, Including the Antimicrobial Activity Spectrum. *Appl. Environ. Microbiol.* 2011;77(8):2749–2754. DOI: 10.1128/AEM.02481-10
  59. Ghiaei A, Ghasemi SM, Shokri D. Investigating the antagonistic effect of indigenous probiotics on carbapenem-resistant pseudomonas aeruginosa strains. *Biomed Res. Int.* 2023;2023. DOI: 10.1155/2023/6645657
  60. Xiong Y, Xu G, Chen M, Ma H. Intestinal uptake and tolerance to food antigens. *Front. Immunol.* 2022;13:1–10. DOI: 10.3389/fimmu.2022.906122
  61. Sicherer SH, Warren CM, Dant C, Gupta RS, Nadeau KC. Food allergy from infancy through adulthood. *J. Allergy Clin. Immunol. Pract.* 2020;8(6):1854–1864. Available:<https://doi.org/10.1016/j.jaip.2020.02.010>.
  62. Peters RL, Krawiec M, Koplin JJ, Santos AF. Update on food allergy. *Pediatr. Allergy Immunol.* 2021;32(4):647–657. Available:<https://doi.org/10.1111/pai.13443>.
  63. Goldin BR. Health benefits of probiotics. *Br. J. Nutr.* 1998;80(SUPPL. 2). DOI: 10.1017/s0007114500006036
  64. Naeem H, et al. Role of probiotics against human cancers, inflammatory diseases, and other complex malignancies. *J. Food Biochem.* 2024;2024. DOI: 10.1155/2024/6632209
  65. Lee YK, Mazmanian SK. Has the microbiota played a critical role in the evolution of the adaptive immune system? *Science* (80-. ). 2010;330(6012):1768–1773. DOI: 10.1126/science.1195568
  66. Spacova I, Ceuppens JL, Seys SF, Petrova MI, Lebeer S. Probiotics against airway allergy: host factors to consider. *DMM Dis. Model. Mech.* 2018;11(7). DOI: 10.1242/DMM.034314
  67. Knipe H, Temperton B, Lange A, Bass D, Tyler CR. Probiotics and competitive exclusion of pathogens in shrimp aquaculture. *Rev. Aquac.* 2021;13(1):324–352. DOI: 10.1111/raq.12477
  68. Silva DR, De J, Sardi CO, De N, Pitangui S, Roque SM, Da Silva ACB, Rosalen PL. Probiotics as an alternative antimicrobial therapy: Current reality and future directions. *J. Funct. Foods.* 2020;73:104080. Available:<https://doi.org/10.1016/j.jff.2020.104080>.
  69. Prabhurajeshwar C, Chandrakanth K. Evaluation of antimicrobial properties and their substances against pathogenic bacteria *In-vitro* by probiotic Lactobacilli strains isolated from commercial yoghurt. *Clin. Nutr. Exp.* 2019;23:97–115. Available:<https://doi.org/10.1016/j.yclnex.2018.10.001>.
  70. Plaza-Díaz J, Ruiz-Ojeda FJ, Vilchez-Padial LM, Gil A. Evidence of the anti-inflammatory effects of probiotics and synbiotics in intestinal chronic diseases. *Nutrients.* 2017;9(6). DOI: 10.3390/nu9060555.
  71. La Storia A, Di Giuseppe FA, Volpe S, Oliviero V, Villani F, Torrieri E. Physical properties and antimicrobial activity of bioactive film based on whey protein and Lactobacillus curvatus 54M16 producer of bacteriocins. *Food Hydrocoll.* 2020;108:105959. Available:<https://doi.org/10.1016/j.foodhyd.2020.105959>.
  72. Pérez-Ramos A, Madi-Moussa D, Coucheney F, Drider D. Current knowledge of the mode of action and immunity mechanisms of lab-bacteriocins. *Microorganisms.* 2021;9(10). DOI: 10.3390/microorganisms9102107.
  73. Martinez FAC, Balciunas EM, Converti A, Cotter PD, De Souza Oliveira RP. Bacteriocin production by Bifidobacterium spp. A review. *Biotechnol. Adv.* 2013;31(4):482–488.

- Available:<https://doi.org/10.1016/j.biotechadv.2013.01.010>.
74. Paterniti I, Scuderi SA, Cambria L, Nostro A, Esposito E, Marino A. Protective effect of probiotics against pseudomonas aeruginosa infection of human corneal epithelial cells. *Int. J. Mol. Sci.* 2024;25(3). DOI: 10.3390/ijms25031770.
  75. Rejaili M, Sani AM. Role of probiotics in health and disease. *Int. J. PharmTech Res.* 2016;9(7):351–356.
  76. Wong S, et al. Lactobacillus casei Shirota probiotic drinks reduce antibiotic associated diarrhoea in patients with spinal cord injuries who regularly consume proton pump inhibitors: A subgroup analysis of the ECLISP multicentre RCT. *Spinal Cord.* 2024;62(5):255–263. DOI: 10.1038/s41393-024-00983-w.
  77. Szajewska H, Kołodziej M. Systematic review with meta-analysis: Lactobacillus rhamnosus GG in the prevention of antibiotic-associated diarrhoea in children and adults. *Aliment. Pharmacol. Ther.* 2015;42(10):1149–1157. Available:<https://doi.org/10.1111/apt.13404>
  78. Bareerah Haidry, Naheem Ahmed, Tayyaba Saddique, Maryam Abbasi, Mashal Yaqoob, and Nuzhat Rasheed. Clinical efficacy of use of probiotics in children with acute watery Diarrhea. *Ann. PIMS-Shaheed Zulfiqar Ali Bhutto Med. Univ.* 2024;20(2):186–191. DOI: 10.48036/apims.v20i2.868.
  79. Daniali M, Nikfar S, Abdollahi M. A brief overview on the use of probiotics to treat overweight and obese patients. *Expert Rev. Endocrinol. Metab.* 2020;15(1):1–4. DOI: 10.1080/17446651.2020.1719068.
  80. Ejtahed HS, et al. Probiotics supplementation for the obesity management; A systematic review of animal studies and clinical trials. *J. Funct. Foods.* 2019;52:228–242. Available:<https://doi.org/10.1016/j.jff.2018.10.039>.
  81. Chen L, et al. Lactobacillus acidophilus suppresses colitis-associated activation of the IL-23/Th17 axis. *J. Immunol. Res.* 2015;2015. DOI: 10.1155/2015/909514
  82. Barendolts E. Gut Microbiota, Prebiotics, Probiotics, and synbiotics in management of obesity and prediabetes: Review of randomized controlled trials. *Endocr. Pract.* 2016;22(10):1224–1234. Available:<https://doi.org/10.4158/EP151157.RA>.
  83. Kadooka Y, et al. Regulation of abdominal adiposity by probiotics (Lactobacillus gasseri SBT2055) in adults with obese tendencies in a randomized controlled trial. *Eur. J. Clin. Nutr.* 2010;64(6):636–643. DOI: 10.1038/ejcn.2010.19
  84. Ben Othman R, et al. A clinical trial about effects of prebiotic and probiotic supplementation on weight loss, psychological profile and metabolic parameters in obese subjects. *Endocrinol. Diabetes Metab.* 2023;6(2):1–7. DOI: 10.1002/edm2.402
  85. Hasani-Ranjbar S, Hoseini Tavassol Z, Malmir H, Ejtahed HS, Tajabadi Ebrahimi M, Larijani B. Investigation of the probiotic supplement's effect on obese adults demonstrated a reduction in fasting insulin levels: A double-blind randomized clinical trial. *J. Diabetes Metab. Disord.* 2024;23(1):1141–1149. DOI: 10.1007/s40200-024-01400-y
  86. Sun J, Buys NJ. Glucose- and glycaemic factor-lowering effects of probiotics on diabetes: A meta-analysis of randomised placebo-controlled trials. *Br. J. Nutr.* 2016;115(7):1167–1177. DOI: 10.1017/S0007114516000076
  87. Rosa CP, Brancaglioni GA, Miyauchi-Tavares TM, Corsetti PP, De Almeida LA. Antibiotic-induced dysbiosis effects on the murine gastrointestinal tract and their systemic repercussions. *Life Sci.* 2018;207:480–491. Available:<https://doi.org/10.1016/j.lfs.2018.06.030>
  88. Mishima Y, Sartor RB. Manipulating resident microbiota to enhance regulatory immune function to treat inflammatory bowel diseases. *J. Gastroenterol.* 2020;55(1):4–14. DOI: 10.1007/s00535-019-01618-1
  89. Yakovlieva M, et al. Influence of Lactobacillus brevis 15 and Lactobacillus plantarum 13 on blood glucose and body weight in rats after high-fructose diet. *Benef. Microbes.* 2015;6(4):505–512. DOI: 10.3920/BM2014.0012
  90. Kobylak N, Falalyeyeva T, Mykhalchyshyn G, Kyriienko D, Komissarenko I. Effect of alive probiotic on insulin resistance in type 2 diabetes patients: Randomized clinical

- trial. *Diabetes Metab. Syndr. Clin. Res. Rev.* 2018;12(5):617–624.  
DOI: 10.1016/j.dsx.2018.04.015
91. Farida E, Nuraida L, Giriwono PE, Jenie BSL. Lactobacillus rhamnosus reduces blood glucose level through downregulation of gluconeogenesis gene expression in streptozotocin-induced diabetic rats. *Int. J. Food Sci.* 2020;2020.  
DOI: 10.1155/2020/6108575
92. Fang B, Zhang M, Dong L, Zhou X, Ren F, Ge S. Probiotic camel milk powder improves glycemic control, dyslipidemia, adipose tissue and skeletal muscle function in T2DM patients: A randomized trial; 2020.
93. Sabico S, et al. Effects of a 6-month multi-strain probiotics supplementation in endotoxemic, inflammatory and cardiometabolic status of T2DM patients: A randomized, double-blind, placebo-controlled trial. *Clin. Nutr.* 2019;38(4): 1561–1569.  
DOI: 10.1016/j.clnu.2018.08.009
94. Sandberg AS, Önning G, Engström N, Scheers N. Iron supplements containing Lactobacillus plantarum 299v increase ferric iron and up-regulate the ferric reductase DCYTB in human Caco-2 /HT29 MTX co-cultures. *Nutrients.* 2018; 10(12).  
DOI: 10.3390/nu10121949
95. Hadadi N, Berweiler V, Wang H, Trajkovski M. Intestinal microbiota as a route for micronutrient bioavailability. *Curr. Opin. Endocr. Metab. Res.* 2021;20:100285.  
Available: <https://doi.org/10.1016/j.coemr.2021.100285>.
96. Ohtani N, Yoshimoto S, Hara E. Obesity and cancer: A gut microbial connection. *Cancer Res.* 2014;74(7):1885–1889.  
DOI: 10.1158/0008-5472.CAN-13-3501
97. Increases Iron Absorption in Young Females- Double Isotope Sequential Single-Blind Studies in Menstruating Women.
98. Korcok DJ, Tršić-Milanovic NA, Ivanovic ND, Dordevic BI. Development of probiotic formulation for the treatment of iron deficiency anemia. *Chem. Pharm. Bull.* 2018;66(4):347–352.  
DOI: 10.1248/cpb.c17-00634
99. Rosen GM, Morrissette S, Larson A, Stading P, Griffin KH, Barnes TL. Use of a probiotic to enhance iron absorption in a randomized trial of pediatric patients presenting with iron deficiency. *J. Pediatr.* 2019;207:192-197.e1.  
DOI: 10.1016/j.jpeds.2018.12.026
100. Ballini A, et al. Probiotics Improve Urogenital Health in Women. Open access Maced. *J. Med. Sci.* 2018;6(10):1845–1850.  
DOI: 10.3889/oamjms.2018.406
101. Gupta V, Mastromarino P, Garg R. Effectiveness of prophylactic oral and/or vaginal probiotic supplementation in the prevention of recurrent urinary tract infections: A randomized, double-blind, placebo-controlled trial. *Clin. Infect. Dis.* 2024;78(5):1154–1161.  
DOI: 10.1093/cid/ciad766
102. Farhana F, et al. Effectiveness of probiotics for the treatment of urinary tract infection among women in Bangladesh. *Med. Res. Its Appl.* 2024;3:25–43.  
DOI: 10.9734/bpi/mria/v3/181
103. Cagetti MG, Mastroberardino S, Milia E, Cocco F, Lingström P, Campus G. The use of probiotic strains in caries prevention: A systematic review. *Nutrients.* 2013;5(7):2530–2550.  
DOI: 10.3390/nu5072530
104. Hedayati-Hajikand T, Lundberg U, Eldh C, Twetman S. Effect of probiotic chewing tablets on early childhood caries - a randomized controlled trial. *BMC Oral Health.* 2015;15(1):1–5.  
DOI: 10.1186/s12903-015-0096-5
105. Jørgensen MR, Castiblanco G, Twetman S, Keller MK. Prevention of caries with probiotic bacteria during early childhood. Promising but inconsistent findings. *Am. J. Dent.* 2016;29(3):127–131.  
Available: <http://europepmc.org/abstract/MED/27505986>
106. Pujia AM, et al. The probiotics in dentistry: A narrative review. *Eur. Rev. Med. Pharmacol. Sci.* 2017;21(6):1405–1412.
107. Wongsupa P, Tunkura K, Jantama R. The Effectiveness of Probiotic Milk Tablets in Preventing Dental Caries Among Young Children: A Cluster Randomized Control Trial- 12-Months Results. 2024;139–145.  
DOI: 10.14456/jdat.2024.14
108. Jach ME, et al. The Role of probiotics and their metabolites in the treatment of depression. *Molecules.* 2023; 28(7).  
DOI: 10.3390/molecules28073213
109. Tian P, et al. Bifidobacterium breve CCFM1025 attenuates major depression disorder via regulating gut microbiome and tryptophan metabolism: A randomized



- clinical trial. *Brain. Behav. Immun.* 2022;100:233–241.  
Available: <https://doi.org/10.1016/j.bbi.2021.11.023>.
110. Oniszczyk A, Oniszczyk T, Gancarz M, Szymańska J. Role of gut microbiota, probiotics and prebiotics in the cardiovascular diseases. *Molecules.* 2021;26(4):1–15.  
DOI: 10.3390/molecules26041172
111. Khongrum J, et al. Safety and Effects of *Lactobacillus paracasei* TISTR 2593 supplementation on improving cholesterol metabolism and atherosclerosis-related parameters in subjects with hypercholesterolemia: A randomized, double-blind, placebo-controlled clinical trial. *Nutrients.* 2023;15(3).  
DOI: 10.3390/nu15030661
112. Parvin A. COVID-19 Dynamics in Rajshahi Division: An RT-qPCR-Based Study. *J. Agric. Food Environ. (JAFE)| ISSN (Online Version) 2708-5694.* 2024;5(2):35–41.
113. Kaur H, Kabra SK. Upper respiratory tract infectio. *Essent. Pediatr. Pulmonol.* 2023;103.
114. Arroll B. Common cold. *BMJ Clin. Evid.* 2011;2011.
115. Vos T, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: A systematic analysis for the Global Burden of Disease Study 2015. *Lancet.* 2016;388(10053):1545–1602.  
DOI: 10.1016/S0140-6736(16)31678-6.
116. Furushima D, et al. Prevention of acute upper respiratory infections by consumption of catechins in healthcare workers: A randomized, placebo-controlled trial. *Nutrients.* 2020;12(1):1–12.  
DOI: 10.3390/nu12010004
117. Zhao Y, Dong BR, Hao Q. Probiotics for preventing acute upper respiratory tract infections. *Cochrane Database Syst. Rev.* 2022;8.  
DOI: 10.1002/14651858.CD006895.pub4
118. Shida K. et al. Daily intake of fermented milk with *Lactobacillus casei* strain Shirota reduces the incidence and duration of upper respiratory tract infections in healthy middle-aged office workers. *Eur. J. Nutr.* 2017;56(1):45–53.  
DOI: 10.1007/s00394-015-1056-1
119. Singh A, Agarwal A, Abhishek E, Jain S, Jawarker R, Raza M. Probiotics as an adjuvant treatment for recurrent aphthous ulcer. *Int. J. Health Sci. (Qassim);* 2022.  
DOI: 10.53730/ijhs.v6ns5.9689
120. Dou X, et al. Probiotic-loaded calcium alginate/fucoidan hydrogels for promoting oral ulcer healing. *Int. J. Biol. Macromol.* 2023;244:125273.  
Available: <https://doi.org/10.1016/j.ijbiomac.2023.125273>.
121. Nirmala M, Smitha SG, Kamath GJ. A Study to assess the efficacy of local application of oral probiotic in treating recurrent aphthous ulcer and oral candidiasis. *Indian J. Otolaryngol. Head Neck Surg.* 2019;71(1): 113–117.  
DOI: 10.1007/s12070-017-1139-9.
122. Lanas A, Chan FKL. Peptic ulcer disease. *Lancet.* 2017;390(10094):613–624.  
Available: [https://doi.org/10.1016/S0140-6736\(16\)32404-7](https://doi.org/10.1016/S0140-6736(16)32404-7).
123. Liang B, Yuan Y, Peng XJ, Liu XL, Hu XK, Xing DM. Current and future perspectives for *Helicobacter pylori* treatment and management: From antibiotics to probiotics. *Front. Cell. Infect. Microbiol.* 2022;12(1–13).  
DOI: 10.3389/fcimb.2022.1042070
124. Siegel RL, Miller KD, Jemal A. Cancer statistics. *CA. Cancer J. Clin.* 2018;68(1):7–30.  
DOI: 10.3322/caac.21442
125. Jemal A, Siegel R, Ward E, Hao Y, Xu J, Thun MJ. Cancer Statistics , 2009 Both Sexes Female Both Sexes Estimated Deaths. *CA Cancer J Clin.* 2009;59(4):1–25.  
DOI: 10.1002/caac.20073.Available.
126. Anderson KN, Schwab RB, Martinez ME. Reproductive risk factors and breast cancer subtypes: a review of the literature. *Breast Cancer Res. Treat.* 2014;144(1):1–10.  
DOI: 10.1007/s10549-014-2852-7
127. Reid G, et al. New scientific paradigms for probiotics and prebiotics. *J. Clin. Gastroenterol.* 2003;37(2):105–118.  
DOI: 10.1097/00004836-200308000-00004
128. Azam R, et al. *Lactobacillus acidophilus* and *Lactobacillus crispatus* culture supernatants downregulate expression of cancer-testis genes in the MDA-MB-231 cell line. *Asian Pacific J. Cancer Prev.* 2014;15(10):4255–4259.  
DOI: 10.7314/APJCP.2014.15.10.4255
129. Tabung FK, Brown LS, Fung TT. Dietary patterns and colorectal cancer risk: A review of 17 Years of Evidence (2000–

- 2016). *Curr. Colorectal Cancer Rep.* 2017;13(6):440–454.  
DOI: 10.1007/s11888-017-0390-5
130. Meng C, Bai C, Brown TD, Hood LE, Tian Q. Human gut microbiota and gastrointestinal cancer. *Genomics. Proteomics Bioinformatics.* 2018;16(1):33–49.  
DOI: 10.1016/j.gpb.2017.06.002
131. Hendler R, Zhang Y. Probiotics in the treatment of colorectal cancer. *Medicines.* 2018;5(3):101.  
DOI: 10.3390/medicines5030101
132. Selvamani S, et al. Efficacy of probiotics-based interventions as therapy for inflammatory bowel disease: A recent update. *Saudi J. Biol. Sci.* 2022;29(5):3546–3567.  
Available: <https://doi.org/10.1016/j.sjbs.2022.02.044>
133. Sivamaruthi BS, Kesika P, Chaiyasut C. A mini-review of human studies on cholesterol-lowering properties of probiotics. *Sci. Pharm.* 2019;87(4).  
DOI: 10.3390/scipharm87040026
134. Bassaganya-Riera J, Viladomiu M, Pedragosa M, De Simone C, Hontecillas R. Immunoregulatory mechanisms underlying prevention of colitis-associated colorectal cancer by probiotic bacteria. *Plos One.* 2012;7(4):e34676.  
Available: <https://doi.org/10.1371/journal.pone.0034676>
135. Faghfoori Z, Pourghassem Gargari B, Saber Gharamaleki A, Bagherpour H, Yari Khosroushahi A. Cellular and molecular mechanisms of probiotics effects on colorectal cancer. *J. Funct. Foods.* 2015;18:463–472.  
Available: <https://doi.org/10.1016/j.jff.2015.08.013>.
136. Tjalsma H, Boleij A, Marchesi JR, Dutilh BE. A bacterial driver–passenger model for colorectal cancer: Beyond the usual suspects. *Nat. Rev. Microbiol.* 2012;10(8):575–582.  
DOI: 10.1038/nrmicro2819
137. Zhu Q, Gao R, Wu W, Qin H. The role of gut microbiota in the pathogenesis of colorectal cancer. *Tumor Biol.* 2013;34(3):1285–1300.  
DOI: 10.1007/s13277-013-0684-4.
138. Verma A, Shukla G. Probiotics *Lactobacillus rhamnosus* GG, *Lactobacillus acidophilus* Suppresses DMH-Induced Procarcinogenic Fecal Enzymes and Preneoplastic Aberrant Crypt Foci in Early Colon Carcinogenesis in Sprague Dawley Rats. *Nutr. Cancer.* 2013;65(1):84–91.  
DOI: 10.1080/01635581.2013.741746
139. Mohania D, Kansal VK, Kruzliak P, Kumari A. Probiotic Dahi Containing *Lactobacillus acidophilus* and *bifidobacterium bifidum* modulates the formation of aberrant crypt foci, mucin-depleted foci, and cell proliferation on 1,2-dimethylhydrazine-induced colorectal carcinogenesis in wistar rats. *Rejuvenation Res.* 2014;17(4):325–333.  
DOI: 10.1089/rej.2013.1537
140. Nowak A, Śliżewska K, Błasiak J, Libudzisz Z. The influence of *Lactobacillus casei* DN 114 001 on the activity of faecal enzymes and genotoxicity of faecal water in the presence of heterocyclic aromatic amines. *Anaerobe.* 2014;30:129–136.  
Available: <https://doi.org/10.1016/j.anaerobe.2014.09.014>.
141. Dos Reis SA, Da Conceição LL, Siqueira NP, Rosa DD, Da Silva LL, Do M, Peluzio CG. Review of the mechanisms of probiotic actions in the prevention of colorectal cancer. *Nutr. Res.* 2017;37:1–19.  
Available: <https://doi.org/10.1016/j.nutres.2016.11.009>.
142. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA. Cancer J. Clin.* 2018;68(6):394–424.  
DOI: 10.3322/caac.21492
143. Sia D, Villanueva A, Friedman SL, Llovet JM. Liver cancer cell of origin, molecular class, and effects on patient prognosis. *Gastroenterology.* 2017;152(4):745–761.  
Available: <https://doi.org/10.1053/j.gastro.2016.11.048>.
144. Ng CKY, Piscuoglio S, Terracciano LM. Molecular classification of hepatocellular carcinoma: The view from metabolic zonation. *Hepatology (Baltimore, Md.)*, vol. 66, no. 5. United States. 2017;1377–1380.  
DOI: 10.1002/hep.29311
145. Ko KL, Mak LY, Cheung KS, Yuen MF. Hepatocellular carcinoma: Recent advances and emerging medical therapies. *F1000Research.* 2020;9.  
DOI: 10.12688/f1000research.24543.1.
146. Gupta H, Youn GS, Shin MJ, Suk KT. Role of gut microbiota in

- hepatocarcinogenesis. *Microorganisms*. 2019;7(5).  
DOI: 10.3390/microorganisms7050121
147. Wan MLY, El-Nezami H. Targeting gut microbiota in hepatocellular carcinoma: Probiotics as a novel therapy. *Hepatobiliary Surg. Nutr.* 2018;7(1):11–20.  
DOI: 10.21037/hbsn.2017.12.07.
148. Liu Q, et al. Surface components and metabolites of probiotics for regulation of intestinal epithelial barrier. *Microb. Cell Fact.* 2020;19(1):23.  
DOI: 10.1186/s12934-020-1289-4
149. Chávez-Tapia NC, et al. Current evidence on the use of probiotics in liver diseases. *J. Funct. Foods.* 2015;17:137–151.  
Available:https://doi.org/10.1016/j.jff.2015.05.009.
150. Zhao Y, et al. Role of lung and gut microbiota on lung cancer pathogenesis. *J. Cancer Res. Clin. Oncol.* 2021;147(8):2177–2186.  
DOI: 10.1007/s00432-021-03644-0.
151. Dickson RP, Huffnagle GB. The Lung microbiome: New principles for respiratory bacteriology in health and disease. *Plos Pathog.* 2015;11(7):e1004923.  
Available:https://doi.org/10.1371/journal.ppat.1004923
152. Dickson RP, Martinez FJ, Huffnagle GB. The role of the microbiome in exacerbations of chronic lung diseases. *Lancet.* 2014;384(9944):691–702.  
Available:https://doi.org/10.1016/S0140-6736(14)61136-3.
153. Dickson RP, Erb-Downward JR, Huffnagle GB. The role of the bacterial microbiome in lung disease. *Expert Rev. Respir. Med.* 2013;7(3):245–257.  
DOI: 10.1586/ers.13.24
154. Dickson RP, Huang YJ, Martinez FJ, Huffnagle GB. The lung microbiome and viral-induced exacerbations of chronic obstructive pulmonary disease: New observations, novel approaches. *American Journal of Respiratory and Critical Care Medicine.* United States. 2013;188 (10): 1185–1186.  
DOI: 10.1164/rccm.201309-1573ED
155. Sadrekarimi H, et al. Emerging role of human microbiome in cancer development and response to therapy: Special focus on intestinal microflora. *J. Transl. Med.* 2022;20(1):301.  
DOI: 10.1186/s12967-022-03492-7
156. Hossain MS, Rupa SI, Sarkar MS, Al Amin M, Khatun MT, Shamim M, Islam MZ. Gender differences-based bioinformatics analysis to identify hub genes and key pathways in type 2 diabetes. *Informatics in Medicine Unlocked.* 2023;40:101302.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/119570>