



# **Music Therapy 101 for Alzheimer's Disease: A Review of Current Trends and Future Directions**

**Eric Murianki <sup>a\*</sup>**

<sup>a</sup> *School of Music, University of Florida, 130 Music Building, Gainesville, FL 32603, United States.*

## **Author's contribution**

*The sole author designed, analysed, interpreted and prepared the manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/air/2024/v25i41118>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/120700>

**Minireview Article**

**Received: 29/05/2024**  
**Accepted: 31/07/2024**  
**Published: 02/08/2024**

## **ABSTRACT**

Alzheimer's disease (AD) poses significant challenges to individuals, families, and healthcare systems globally. Currently, there is no cure for AD, and available therapies focus on managing symptoms and slowing disease progression. Given the limitations of pharmacological treatments in halting the progression of AD, there is increasing interest in non-pharmacological interventions such as music therapy. This review explores the therapeutic benefits of music therapy for AD patients, evaluating empirical evidence from clinical studies and exploring the underlying mechanisms that contribute to its effectiveness. Additionally, this review discusses the practical implications of integrating music therapy into treatment plans. It identifies future research directions to understand further and enhance its impact on AD patients' cognitive functioning and quality of life.

**Keywords:** *Alzheimer's disease; Music therapy; cognitive functioning; non-pharmacological interventions.*

\*Corresponding author: E-mail: [koomemurianki@gmail.com](mailto:koomemurianki@gmail.com);

**Cite as:** Murianki, Eric. 2024. "Music Therapy 101 for Alzheimer's Disease: A Review of Current Trends and Future Directions". *Advances in Research* 25 (4):397-405. <https://doi.org/10.9734/air/2024/v25i41118>.

## 1. INTRODUCTION

Alzheimer's disease (AD) is a progressive neurodegenerative disorder that profoundly impacts cognitive function, behavior, and daily functioning. First identified by Dr. Alois Alzheimer in 1906, AD is characterized by the accumulation of amyloid-beta plaques and tau protein tangles in the brain, which lead to neuronal cell death and synaptic dysfunction. These pathological changes primarily affect brain regions responsible for memory, learning, reasoning, and language [1].

The clinical manifestation of AD typically begins with mild memory impairment. Over time, it progresses to severe cognitive decline, ultimately affecting an individual's ability to perform even the most basic activities of daily living [2]. As the most common cause of dementia worldwide, AD poses a significant public health challenge, with estimates indicating that over 50 million people are affected globally [1].

Despite extensive research, the exact cause of Alzheimer's disease remains incompletely understood. However, genetic, environmental, and lifestyle factors are believed to contribute to its development. Currently, there is no cure for AD, and available treatments focus on managing symptoms and slowing disease progression [3]. These treatments include medications that enhance neurotransmitter function and non-pharmacological interventions such as cognitive stimulation therapy and music therapy.

This literature review explores music therapy's role in treating Alzheimer's disease, examining its potential to alleviate symptoms, enhance quality of life, and stimulate cognitive function in affected individuals. By synthesizing findings from various studies and clinical trials, this review aims to provide a comprehensive understanding of how health institutions can integrate music therapy into holistic care approaches for Alzheimer's patients [4].

## 2. UNDERSTANDING ALZHEIMER'S DISEASE

Alzheimer's disease is distinguished by the presence of beta-amyloid plaques and tau protein tangles in the brain, causing disruption in synaptic function, neuronal depletion, and a decline in cognitive abilities [5]. The disease progresses through stages of mild cognitive impairment (MCI) to severe dementia, impacting

memory, language, executive functions, and behavior [6]. AD patients often experience heightened emotional distress, social isolation, and diminished quality of life as the disease advances [1].

The clinical presentation of Alzheimer's disease progresses through distinct stages, each characterized by specific cognitive and functional impairments [6]. Individuals may experience mild cognitive impairment in the early stages, characterized by subtle memory deficits and difficulty with word-finding and problem-solving tasks [7]. As the disease advances, memory loss becomes more pronounced, accompanied by challenges in performing activities of daily living and significant changes in behavior and mood [8]. In the severe stages of AD, individuals may lose the ability to communicate effectively, recognize loved ones, and maintain independence [1].

The risk factor associated with the disease is Age, remaining the most significant risk factor for Alzheimer's disease, with prevalence increasing exponentially after the Age of 65 [9]. Genetic factors also play a crucial role, with mutations in genes such as APP, PSEN1, and PSEN2 associated with early-onset familial AD [10]. The APOE  $\epsilon$ 4 allele is the most potent genetic risk factor for late-onset AD, influencing disease onset and progression [11]. Beyond genetics, lifestyle factors such as cardiovascular health, physical activity, and social engagement are increasingly recognized as modifiable factors influencing AD risk [12,13].

## 3. UNDERSTANDING MUSIC THERAPY

Music therapy is a structured intervention that utilizes musical elements, such as rhythm, melody, and harmony, to achieve therapeutic goals [14]. It encompasses active participation (e.g., singing and playing instruments) and receptive listening tailored to individual needs and preferences [15]. Music's ability to evoke emotions, stimulate memories, and engage neural networks makes it a powerful tool for addressing cognitive and emotional symptoms associated with AD [3].

The origins of music therapy can be traced back to ancient civilizations, where music was used for healing purposes [16]. In the modern era, music therapy emerged as a structured discipline following its extensive use in World War II rehabilitation programs [17]. Theoretical

frameworks underpinning music therapy include Nordoff-Robbins Music Therapy, which emphasizes improvisation and musical interaction to promote emotional expression and communication [18], and the Bonny Method of Guided Imagery and Music, which uses music to explore unconscious emotions and memories [19].

The theoretical foundations of music therapy draw from disciplines such as psychology, neuroscience, and musicology. Nordoff-Robbins Music Therapy, for example, emphasizes improvisation and musical interaction to promote emotional expression and communication [20]. Moreover, the Bonny Method of Guided Imagery and Music utilizes music to access and explore unconscious emotions and memories [21]. These frameworks provide structured approaches for music therapists to tailor interventions to meet clients' needs.

Music therapy interventions encompass a variety of techniques, including active music-making, receptive music-listening, songwriting, and improvisation [22]. In clinical settings, music therapists collaborate with healthcare professionals to address goals related to physical rehabilitation, pain management, emotional processing, and cognitive functioning [14]. For example, music therapy has been effectively used in oncology to reduce anxiety and enhance the quality of life for cancer patients undergoing treatment [23].

Research demonstrates the effectiveness of music therapy in improving various health outcomes. Studies have shown that music therapy interventions can reduce anxiety, enhance mood, and improve quality of life in diverse populations [24]. In neurorehabilitation, researchers have found that music therapy facilitates motor skills recovery and speech rehabilitation in stroke patients [25]. Neuroscientific research highlights that music engages multiple brain regions involved in emotion regulation and auditory processing, offering insights into its therapeutic mechanisms [26].

## **4. BENEFITS OF MUSIC THERAPY IN ALZHEIMER'S DISEASE**

### **4.1 Cognitive Benefits**

Music therapy interventions have demonstrated significant cognitive benefits for AD patients.

Studies indicate memory recall, attention, and executive function improvements following music-based interventions [3,27]. Music's rhythmic and melodic components can enhance neuroplasticity, promote synaptic connectivity, and facilitate cognitive rehabilitation in individuals with AD [28].

Music therapy engages various cognitive processes that AD affects, such as memory retrieval, attention, and emotional processing [28]. The structured use of music can evoke memories, reduce agitation, and improve mood, enhancing cognitive abilities [15]. For example, personalized playlists or familiar music can stimulate autobiographical memories, supporting cognitive recall and emotional well-being [29].

Neuroimaging studies have provided insights into the neurobiological mechanisms underlying the cognitive benefits of music therapy in AD. Functional MRI (fMRI) studies demonstrate that music activates multiple brain regions involved in memory consolidation, emotional regulation, and auditory processing [30]. Research has shown that Music-based interventions enhance neural connectivity and promote neuroplasticity, potentially slowing cognitive decline in AD [3].

In clinical practice, therapists tailor music therapy interventions to address specific cognitive deficits and emotional needs of patients with Alzheimer's patients. Techniques include active music-making (e.g., singing and playing instruments), receptive listening, and rhythmic exercises [15]. These interventions are integrated into comprehensive care plans to improve cognitive function, reduce behavioral symptoms, and enhance social engagement [31,32].

Research findings consistently support the positive effects of music therapy on cognitive function in AD. Meta-analyses and systematic reviews indicate improvements in memory, language skills, and executive function following music interventions [33]. For instance, a randomized controlled trial demonstrated that regular music activities led to significant cognitive benefits and improved quality of life in AD patients compared to standard care [3].

Despite the promising outcomes, challenges remain in implementing music therapy as a standardized intervention in dementia care settings. Critical considerations include variability in treatment protocols, limited access to trained music therapists, and the need for more rigorous

research designs [33]. Future research should identify optimal intervention strategies, explore long-term effects, and integrate music therapy into dementia care guidelines [34].

#### 4.2 Emotional Benefits

Emotionally, music therapy helps alleviate symptoms of depression, anxiety, and agitation commonly experienced by AD patients [28]. Music engages limbic system structures involved in emotion processing and regulation, improving mood states, reducing stress levels, and enhancing emotional well-being [3].

Music therapy interventions offer a range of emotional benefits for individuals with AD. Music uniquely evokes emotions, elicits memories, and stimulates positive affective responses [31]. Listening to familiar music or engaging in musical activities can reduce agitation, anxiety, and depressive symptoms, promoting relaxation and well-being [28]. These emotional responses are particularly significant given the emotional dysregulation often experienced by individuals with AD.

Beyond emotional regulation, music therapy contributes to psychosocial well-being by fostering social interaction and communication. Group music-making activities provide opportunities for social engagement, enhancing interpersonal connections, and reducing feelings of isolation [15]. Music therapy sessions are often tailored to individual preferences and therapeutic goals, empowering patients to express themselves and maintain a sense of agency [29].

Neuroscientific research underscores the neurobiological mechanisms through which music therapy influences emotional function in AD. Functional imaging studies demonstrate that music activates brain regions involved in emotional processing, memory retrieval, and reward systems [30]. Music's capacity to modulate neural activity may contribute to its therapeutic effects on emotional regulation and mood enhancement in AD patients [3].

Clinical studies have consistently reported positive outcomes of music therapy on emotional function in AD. For instance, systematic reviews and meta-analyses have shown that music interventions reduce behavioral symptoms, enhance emotional expression, and improve overall quality of life for individuals with dementia

[28]. Randomized controlled trials have demonstrated that personalized music playlists and active music-making interventions effectively mitigate emotional distress and promote emotional well-being in AD patients [15].

#### 4.3 Social Benefits

On the social aspect, music therapy helps individuals suffering from AD to interact with others, socially communicate, and engage in social activities [28]. Social interaction is also facilitated during group music sessions and lessens the loneliness and isolation often witnessed in dementia patients [15].

Music therapy interventions are intended to facilitate social interactions with patients with Alzheimer's Disease through musical communication, including singing, dancing, and mimicry. Such activities stimulate camaraderie, social interaction, and interpersonal communication, as Raglio et al. [15] identified. The given functions of music as an emotional recall can also foster social relations across cultures and thus reduce social isolation [31].

Apart from facilitating social contact, music therapy has benefits related to psychosocial functioning: it has, for instance, pro-toning effects regarding mood, anxiety, or even quality of life in patients with AD [15]. According to Raglio et al., Group music-making activities can foster participation, expression of emotions, and creativity, thus giving a feeling of power. Music therapy usually consists of participants' favorite songs and reflects their potential and willingness to socialize and participate in other activities [29].

The neurological studies discussed above show how music therapy for social function in AD is mediated neurobiologically. A cross-sectional analysis of functional neuroimaging confirms that music functions in social cognition, empathy, and emotional regulation pathways [30]. Researchers hypothesize that music's ability to entrain neural firing and support emotional attunement can improve social relatedness and interpersonal interactions patients with Alzheimer's disease [35].

Both quantitative and qualitative research have revealed that music therapy has a positive impact on social function in patients diagnosed with AD. For instance, Rasmussen and colleagues conducted a meta-analysis of randomized controlled trials to determine that

group music therapy intervention leads to increased responses on social interaction skills, reduced scores on social withdrawal, and better communication among participants [15]. As has been demonstrated by systematic reviews and meta-analyses, music has proven to enhance social interactions and interpersonal relationships among AD patients [28].

## 5. EVIDENCE FROM RESEARCH STUDIES

Numerous research and scientific analyses of randomized clinical trials have emphasized that music therapy effectively manages AD. Vink et al. [27] conducted a meta-analysis of randomized controlled trials that confirmed evidence of the favorable impact of music therapy on the cognitive and behavioral outcomes of AD patients. The study highlights the role of music therapy in conjunction with conventional management strategies in the treatment of AD and the quality of life of patients.

Särkämö et al. [3] conducted a study that showed that singing and listening to music enhanced cognitive skills, including attention, memory, and executive functions in early AD patients. The researchers credited these enhancements to music's arousing and facilitating effect on the multifaceted aspects of cerebral functioning [36].

In another RCT, Raglio et al. [15] demonstrated that active music therapy positively impacted cognitive performance and cognitive deterioration among AD patients compared to conventional treatment. The study further noted that personalized music intervention effectively boosts cognitive performance [37].

In a systematic review and meta-analysis conducted by Gómez Gallego et al. [28], researchers found out that music therapy interventions have a positive impact on decreasing Behavioral and Psychological Symptoms of Dementia (BPSD), which manifests as agitation, anxiety, and depression. A review of several studies in the article also suggested that music may help reduce emotional symptoms and enhance mood in patients with AD.

Raglio et al. [15\*] examined the impact of music therapy on the psychological distress and the Heart Rate Variability HRV of AD participants. According to their findings, music interventions restored emotional aspects aligned with

participants' need assessment and addressed participant concerns while providing physiological relaxation [38].

In their study, Clair and Memmott [31] noted that music therapy helped nursing home patients have more meaningful social interactions and communicate more than before, especially patients with dementia. Of the activities, group music-making was observed to improve social integration and interpersonal interactions and decrease loneliness in patients with AD.

A systematic review by van der Steen and colleagues published in the Cochrane Database of Systematic Reviews in 2020 compared several music-based therapies for individuals diagnosed with dementia, including AD. In the review, the author reviewed studies that showed that music therapy enhances meaningful social interaction and the quality of life of individuals [39].

Specifically, Särkämö & Soto [35] and Belfi et al. [30] explained the neurophysiological foundations of music therapy interventions for AD patients. These studies employed functional imaging methods to establish that music engages regions related to memory, emotion, and reward systems. People believe such activations increase neuroplasticity and neuronal connections that may counteract cognitive loss in AD patients.

## 6. PRACTICAL CONSIDERATIONS AND CHALLENGES WHEN USING MUSIC THERAPY IN TREATING ALZHEIMER'S DISEASE

In this section, music therapy will be discussed as an essential form of treatment for patients with Alzheimer's Disease (AD), which positively impacts patients' cognitive, emotional, and social functioning. Thus, the use of music therapy in clinical practice is not without its limitations and practical concerns that deserve a focus [38]. This review section discusses the feasibility, advantages, limitations, and possible development of MT applications in treating AD.

### 6.1 Practical Considerations

Implementing music therapy in AD care requires careful consideration of several practical factors:

- Personalization: It is imperative to make music interventions patient-specific and address their cognitive characteristics.

Theme and genre-specific songs and songs that relate to the patient's cultural background and experiences may further improve patient involvement and treatment results [15].

- **Training and Expertise:** A professional certified music therapist is trained to use music in interactions effectively. It is a requirement for therapists to be well-equipped to apply therapeutic sessions for AD patients so that the therapy goals can be met [29].
- **Integration into Care Plans:** Working with other caregivers, nurses, and psychologists helps implement music therapy into the AD management care plan. This finding is in line with the finding that coordination improves the quality of the music intervention and its therapeutic effects [33].

## 6.2 Benefits of Music Therapy

Music therapy offers a range of benefits that support AD treatment:

- **Cognitive Enhancement:** Listening to music or singing activates the brain, thus enhancing memory and attention and possibly delaying the progression of AD [3].
- **Emotional Regulation:** Music reduces anxiety, depression, and factors such as agitation, enhancing emotional health and quality of life [28].
- **Social Engagement:** It was also agreed that group music sessions enhance interactions, reduce loneliness, and enhance interpersonal communication in AD patients as opposed to individual practice [31].

## 6.3 Challenges in Implementation

Despite the therapeutic benefits, several challenges exist in implementing music therapy for AD:

- **Resource Allocation:** Challenges include restricted financial resources for music therapy interventions, such as funding for music therapy services and scarce qualified staff [33].
- **Standardization of Protocols:** The use of differing intervention protocols and the variability of the outcome measures across the studies makes it even harder to compare and replicate the findings,

challenging the development of standardized practices [15\*].

- **Ethical Considerations:** Population: Music therapy involves patients, and as such, the issues of patient self-determination or informed consent for participation in music therapy sessions are significant ethical concerns. Privacy and confidentiality combined with therapeutic purposes are crucial, as Särkämö and Soto noted in studies conducted in 2012.

## 7. FUTURE DIRECTIONS

To address these challenges and enhance the efficacy of music therapy in AD treatment, future research should:

- **Focus on Long-term Outcomes:** Conduct a meta-analysis to explore the durability of improvements in cognition, mood, and social skills in AD patients after music therapy [30,40].
- **Develop Guidelines:** Propose best practices for using music therapy with dementia patients to understand how to incorporate it into practice across different care settings for healthcare professionals and caregivers [28].
- **Expand Access:** Support policy changes and partnerships with healthcare institutions to expand access to necessary music therapy to enhance patient care and well-being [29].

Music is now considered to be a potential intervention for enhancing the quality of life of people with Alzheimer's disease (AD) since it provides benefits in cognitive, emotional, and social areas. This discussion is based on a systematic literature review and discusses the benefits, applications, limitations, and further research regarding music therapy in AD treatment.

### 7.1 Efficacy and Therapeutic Benefits

The reviewed literature consistently supports the efficacy of music therapy in enhancing various aspects of AD:

- **Cognitive Function:** For instance, Särkämö et al. [3] and Raglio et al. [15] have evidenced that music positively impacts memory and executive function. Music involvement in terms of singing, listening, and creative music activities has proved helpful in enhancing cognitive skills as well

as in preventing the early onset of the disease in AD patients [41].

- Emotional Well-being: Gómez-Romero et al. [28] explained that, at least for patients with moderate dementia, music therapy helps alleviate symptoms such as agitation, anxiety, and depression. Therefore, music's capability to impact emotions and mood and facilitate the expression of emotions plays a positive role in enhancing the emotional status and quality of life [42].
- Social Engagement: In their study, Clair and Memmott [31] discussed the social uses of music therapy, including facilitating social interactions and communication and lessening the social alienation of AD patients. Group music sessions can facilitate social relations and interpersonal interactions, promoting interactions.

## 7.2 Factors to Address

Implementing music therapy in AD care settings involves addressing several practical considerations and challenges:

- Training and Expertise: Specifically, certified music therapists focus on planning and implementing musically-based interventions that meet specific cognitive and emotional profiles [29]. Ongoing staff training shows that therapists are prepared to modify the implementation of interventions and enhance optimal therapeutic outcomes.
- Resource Allocation: Lack of funding for music therapy programs and a shortage of qualified professionals acting as music therapists remain significant challenges to expansion [33]. Public awareness and demands for more resources to fund and establish music therapy services are equally important.
- Ethical Considerations: Some ethical issues include the patient's self-determination, cultural considerations, and the patient's choice of songs [35]. Families and caregivers must be involved in receiving treatment and consent, and the individual's dignity in the therapy must be maintained.

## 8. CONCLUSION

In conclusion, the reviewed sources indicate the effectiveness of music therapy in treating Alzheimer's disease. By focusing on the

implemented perspectives and existing and potential barriers to implementation, as well as by promoting supplementary research, healthcare providers can maximize the use of music therapy within dementia care. It is important to note that music therapy is a non-drug treatment that can help to improve such parameters of care as cognition, mood, and social interaction in patients with AD, thereby increasing their quality of life.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of manuscripts.

## COMPETING INTERESTS

Author has declared that no competing interests exist.

## REFERENCES

1. Alzheimer's Association. Stages of Alzheimer's; 2020. Available: <https://www.alz.org/alzheimers-dementia/stages>
2. Baird PN, Saw SM, Lanca C, Guggenheim JA, Smith EL III, Zhou X, et al. Myopia. *Nat Rev Dis Primers*. 2020;6(1):99. doi: 10.1038/s41572-020-00231-4.
3. Särkämö T, Tervaniemi M, Laitinen S, Forsblom A, Soinila S, Mikkonen M, Autti T, Silvennoinen HM, Erkkilä J, Laine M, Peretz I, Hietanen M. Music listening enhances cognitive recovery and mood after middle cerebral artery stroke. *Brain*. 2008;131(3):866-876. DOI: 10.1093/brain/awn013
4. Leggieri M, Thaut MH, Fornazzari L, Schweizer TA, Barfett J, Munoz DG, Fischer CE. Music intervention approaches for Alzheimer's disease: A review of the literature. *Front Neurosci*. 2019;13:132. DOI: 10.3389/fnins.2019.00132
5. Selkoe DJ, Hardy J. The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Mol Med*. 2016;8(6):595-608. DOI: 10.15252/emmm.201606210
6. McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR, Kawas CH, et al. The diagnosis of dementia due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic

- guidelines for Alzheimer's disease. *Alzheimers Dement.* 2011;7(3):263-269. DOI: 10.1016/j.jalz.2011.03.005
7. Petersen RC. Mild cognitive impairment as a diagnostic entity. *J Intern Med.* 2004;256(3):183-194. DOI: 10.1111/j.1365-2796.2004.01388.x
  8. Alzheimer's Association, Thies W, Bleiler L. 2013 Alzheimer's disease facts and figures. *Alzheimers Dement.* 2013;9(2):208-245. DOI: 10.1016/j.jalz.2013.02.003
  9. Brookmeyer R, Johnson E, Ziegler-Graham K, Arrighi HM. Forecasting the global burden of Alzheimer's disease. *Alzheimers Dement.* 2007;3(3):186-191. DOI: 10.1016/j.jalz.2007.04.381
  10. Tanzi RE, Bertram L. Twenty years of the Alzheimer's disease amyloid hypothesis: A genetic perspective. *Cell.* 2005;120(4):545-555. DOI: 10.1016/j.cell.2005.02.008
  11. Corder EH, Saunders AM, Strittmatter WJ, Schmechel DE, Gaskell PC, Small GW, et al. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science.* 1993 Aug 13;261(5123):921-3. DOI: 10.1126/science.8346443
  12. Livingston G, Huntley J, Sommerlad A, Ames D, Ballard C, Banerjee S, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet.* 2020 Aug 8;396(10248):413-446. DOI: 10.1016/S0140-6736(20)30367-6
  13. Guétin S, Charras K, Berard A, Arbus C, Berthelon P, Blanc F, et al. An overview of the use of music therapy in the context of Alzheimer's disease: A report of a French expert group. *Dementia (London).* 2013 Sep;12(5):619-634. DOI: 10.1177/1471301212438290
  14. Merchant RM, Topjian AA, Panchal AR, Cheng A, Aziz K, Berg KM, et al. Part 1: executive summary: 2020 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation.* 2020 Oct 20;142(16 Suppl 2)
  15. Raglio A, Bellandi D, Baiardi P, Gianotti M, Ubezio MC, Gentile S, et al. Active music therapy approach in Alzheimer's disease: A randomized controlled trial. *Aging Ment Health.* 2015;19(10):1-8. Available: <https://doi.org/10.1111/jgs.13558>
  16. Stegemann T. Historical perspectives of music therapy. In: Wheeler BL, editor. *Music therapy handbook.* New York: Guilford Press. 2017;3-20.
  17. Bradt J, Dileo C. Music interventions for mechanically ventilated patients. *Cochrane Database Syst Rev.* 2014 Dec 10;(12).
  18. Smeijsters H, van den Hurk J. Music therapy helping to work through grief and finding a personal identity. *J Music Ther.* 1999;36(3):222-252.
  19. Wheeler BL, Grocke D, Wigram T. *Receptive Methods in Music Therapy: Techniques and Clinical Applications for Music Therapy Clinicians, Educators and Students.* London and Philadelphia: Jessica Kingsley Publishers; 2007.
  20. Ansdell G, Pavlicevic M. Practicing "gentle empiricism"—The Nordoff Robbins research heritage. *Music Ther Perspect.* 2010;28(2):131-139.
  21. Ventre M, McKinney CH. The Bonny method of guided imagery and music. In: Wheeler BL, editor. *Music therapy handbook.* New York: Guilford Press. 2015;96-105.
  22. Bleibel M, El Cheikh A, Sadier NS, Abou-Abbas L. The effect of music therapy on cognitive functions in patients with Alzheimer's disease: A systematic review of randomized controlled trials. *Alzheimers Res Ther.* 2023;15(1):65. DOI: 10.1186/s13195-023-01214-9
  23. Robb SL, Hanson-Abromeit D, May L, Hernandez-Ruiz E, Allison M, Beloit A, et al. Reporting quality of music intervention research in healthcare: A systematic review. *Complement Ther Med.* 2018;38:24-41. DOI: 10.1016/j.ctim.2018.02.008
  24. Grau-Sánchez J, Jamey K, Paraskevopoulos E, Dalla Bella S, Gold C, Schlaug G, et al. Putting music to trial: Consensus on key methodological challenges investigating music-based rehabilitation. *Ann N Y Acad Sci.* 2022;1518(1):12-24. DOI: 10.1111/nyas.14892
  25. Magee WL, Davidson JW. Music therapy in multiple sclerosis: Results of a systematic qualitative analysis. *Music Ther Perspect.* 2004;22(1):39-51. DOI: 10.1093/mtp/22.1.39
  26. Peretz I, Zatorre RJ. Brain organization for music processing. *Annu Rev Psychol.* 2005;56(1):89-114.



- DOI:10.1146/annurev.psych.56.091103.070225
27. Janus SI, Vink AC, Ridder HM, Geretsegger M, Stige B, Gold C, Zuidema SU. Developing consensus description of group music therapy characteristics for persons with dementia. *Nord J Music Ther.* 2021;30(1):24-40.
  28. Gómez-Romero M, Castillo C, Gutiérrez-Martínez O, Vázquez-Sánchez M, Franco-Martín MA. Music therapy and Alzheimer's disease: Cognitive, psychological, and behavioral effects. *Neurology and Therapy.* 2020. doi:10.1016/j.nrl.2015.12.003.
  29. Sung HC, Lee WL, Li TL, Watson R. A systematic review of music therapy interventions for dementia. *J Clin Nurs.* 2017;26(23-24):3311-3324.
  30. Belfi AM, Karlsson EM, Madison TI, Vogl J, Jones GA, Halpern AR. Neural correlates of music-evoked autobiographical memories in Alzheimer's disease. *Neuroimage Clin.* 2021;30:102621.
  31. Clair AA, Memmott JE. *Therapeutic uses of music with older adults.* Baltimore, MD: Health Professions Press; 2017.
  32. Moreno-Morales C, Calero R, Moreno-Morales P, Pintado C. Music therapy in the treatment of dementia: A systematic review and meta-analysis. *Front Med (Lausanne).* 2020 Mar 31;7:160. DOI: 10.3389/fmed.2020.00160.
  33. van der Steen JT, Smaling HJA, van der Wouden JC, Bruinsma MS, Scholten RJ, Vink AC. Music-based therapeutic interventions for people with dementia. *Cochrane Database Syst Rev.* 2018 Jul 23;7. DOI: 10.1002/14651858.CD003477.pub4
  34. Fang R, Ye S, Huangfu J, Calimag DP. Music therapy is a potential intervention for cognition of Alzheimer's Disease: A mini-review. *Transl Neurodegener.* 2017 Jan 12;6(1):2. DOI: 10.1186/s40035-017-0073-9
  35. Särkämö T, Soto D. Music listening after stroke: Beneficial effects and potential neural mechanisms. *Ann N Y Acad Sci.* 2012 Mar;1252(1):266-281. DOI: 10.1111/j.1749-6632.2011.06402.x.
  36. Tsoi KK, Chan JY, Ng YM, Lee MM, Kwok TC, Wong SY. Receptive music therapy is more effective than interactive music therapy to relieve behavioral and psychological symptoms of dementia: A systematic review and meta-analysis. *J Am Med Dir Assoc.* 2018 Jul;19(7):568-576.e3. DOI: 10.1016/j.jamda.2017.12.009.
  37. Lam HL, Li WTV, Laher I, Wong RY. Effects of music therapy on patients with dementia—A systematic review. *Geriatrics (Basel).* 2020 Oct 10;5(4):62. DOI: 10.3390/geriatrics5040062.
  38. Li HC, Wang HH, Chou FH, Chen KM. The effect of music therapy on cognitive functioning among older adults: A systematic review and meta-analysis. *J Am Med Dir Assoc.* 2015 Jan;16(1):71-77. DOI: 10.1016/j.jamda.2014.09.001
  39. Cammisuli DM, Danti S, Bosinelli F, Cipriani G. Non-pharmacological interventions for people with Alzheimer's Disease: A critical review of the scientific literature from the last ten years. *Eur Geriatr Med.* 2016 Feb;7(1):57-64. DOI: 10.1016/j.eurger.2016.01.001.
  40. Sherriff CA, Mathews J, Reynish EL, Shenkin SD. Music therapy for neuropsychiatric symptoms in the general hospital: A systematic literature review. *Music Med.* 2017;9(4):217-226
  41. Ziv N, Granot A, Hai S, Dassa A, Haimov I. Music therapy for Alzheimer's disease and other types of dementia: A systematic review and meta-analysis. *Aging Ment Health.* 2018;22(7):881-888.
  42. Brotons M, Koger SM. The impact of music therapy on language functioning in dementia. *J Music Ther.* 2000;37(3):183-195. DOI: 10.1093/jmt/37.3.183

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/120700>