



Effect of Bio-fertilizers on Shoot Growth of Dragon Fruit Cuttings (*Hylocereus undatus* L.)

Mahandra Kumar Dhanka ^a, R.S.Verma ^{a*}, Anup Baitha ^a, Shikha Choudhary ^a, Saurabh Verma ^a and Bipin Kumar ^a

^a *Department of Horticulture, School of Agricultural sciences and Technology, Babasaheb Bhimrao Ambedkar University, Vidya-Vihar, Rae Bareilly Road, Lucknow 226025 (U.P), India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/arja/2024/v17i4542>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123958>

Original Research Article

Received: 23/07/2024

Accepted: 25/09/2024

Published: 04/10/2024

ABSTRACT

The present investigation was carried out under open field condition at Horticulture Research Farm, Babasaheb Bhimrao Ambedkar University, Lucknow Uttar Pradesh., India to determine the effect of bio-fertilizers on the shoot growth of dragon fruit cutting (*Hylocereus undatus* L.) during the year of 2022-23. The experiment was laid out in a randomized complete block design (RCBD) with three replications. There were 09 treatments, viz., T₁-Control, T₂-Azotobacter (1%), T₃-Phosphate-solubilizing bacteria-1%), T₄-Azotobacter (2%), T₅- Phosphate-solubilizing bacteria (2%), T₆-Azotobacter (1%) + Phosphate-solubilizing bacteria (1%), T₇- Azotobacter (1%) + Phosphate-solubilizing bacteria (2%), T₈- Azotobacter (2%) + Phosphate-solubilizing bacteria (1%) and T₉-Azotobacter (2%) + Phosphate-solubilizing bacteria (2%). Treatments have showed significant differences among the growth parameters in terms of number of days taken for sprouting, percent

*Corresponding author: E-mail: ravihort.009@gmail.com;

Cite as: Dhanka, Mahandra Kumar, R.S.Verma, Anup Baitha, Shikha Choudhary, Saurabh Verma, and Bipin Kumar. 2024. "Effect of Bio-Fertilizers on Shoot Growth of Dragon Fruit Cuttings (*Hylocereus Undatus* L.)". *Asian Research Journal of Agriculture* 17 (4):417-22. <https://doi.org/10.9734/arja/2024/v17i4542>.

sprouting, number of sprouts per cutting, length of the newly emerged shoot, diameter of shoot, average number of spine/areoles, diameter of shoot average number of spine/areoles.

Among the treatment combination least number of days taken for sprouting ,maximum percent sprouting , maximum number of sprouts per cutting, maximum increase in length of the newly emerged shoot, maximum number of spine/areoles, maximum increase in diameter of shoot, maximum fresh weight of shoot and maximum dry weight of shoot was found in treatment T₈-Azotobacter (2%) + Phosphate-solubilizing bacteria (1%) followed by T₉-Azotobacter (2%)+ Phosphate-solubilizing bacteria (2%) while minimum has recorded in T₁(Control).

Keywords: Bio-fertilizers; shoot growth; dragon fruit cuttings; growth parameters.

1. INTRODUCTION

“Dragon fruit (*Hylocereus undatus*) is a cactus, belonging to family cactaceae. Recently, dragon fruit has been introduced as super fruit in India. Its considered to be a promising and remunerative fruit crop. Dragon fruit is a long day plant with beautiful night-blooming flowers that is nicknamed Noble Woman or Queen of the Night. The fruit is also known as strawberry, pear, pitahaya, night blooming cereus, Belle of the Night, Cinderella plant, and Jesus in the Cradle. The fruit is named as Pitaya because of the bracts or scales on the fruit skin and he pitaya means the scaly fruit. The fruit has a very attractive colour and mellow mouth-melting pulp with a black colour edible seed embedded in the pulp along with tremendous nutritive properties. It attracts the growers from different parts of India to cultivate this fruit crop. Dragon fruit is native to tropical and subtropical forest regions of Mexico and Central South America” [1]. “It is a nutritious fruit with a variety of uses. The fruit pulp can be eaten as fresh and can also be made into various valuable processed products. The fruit possesses some medicinal properties. It is known to prevent colon cancer and diabetes, neutralizes toxic substances such as heavy metals, reduce cholesterol and high blood pressure. It is also reported to control high sugar levels. The fruit is rich in vitamin C, phosphorous and calcium which help to develop strong bones, teeth and skin. It is considered as ‘healthfruit’ [2-4]. Betalains have a great potential in colouring a broad array of food. In this view, betacyanins from red coloured dragon fruit are most promising, not only as colouring agents but also in possessing antiradical potential” [5-7]. “Dragon fruit was introduced in 1990 for its commercial cultivation in South Asian tropical countries. At present, significant production and expansion of fruit are occurring in many countries viz., Australia, Cambodia, China, Columbia, Ecuador, Guatemala, Hawaii, Indonesia, Israel, Japan, Laos, Malaysia, Mexico, New Zealand,

Nicaragua, Peru, Philippines, Spain, Sri Lanka, Taiwan, Thailand, South Western USA and Vietnam” (Barbeau,1990, Wu and Chen, 1997). “The plants propagated through stem cuttings starts flowering within 12 to 18 months after planting. Growing media is the important factor for the plants that give anchorage to the plants and provide essential nutrients required by the plants. The growing media enriched with bio-fertilizers possess the advantages like more availability of nutrients in the available forms through natural process like nitrogen fixing, phosphorus solubilizing and stimulate plant growth through the synthesis of growth promoting substances. They build up soil microflora and thereby maintains soil health” (Barbeau,1990). Therefore, the experiment was conducted to study the effect of bio-fertilizers on the shoot growth of dragon fruit cutting.

2. MATERIALS AND METHODS

The present investigation was carried out at Horticulture Research Farm, Department of Horticulture, Babasaheb Bhimrao Ambedkar University (A Central University), Lucknow, Uttar Pradesh, India during the year of 2022-23 to study the effect of bio-fertilizers on shoot growth of dragon fruit cuttings under open field condition. The experiment was laid out in randomized complete block design (RCBD) with three replications. The various treatment combinations of bio-fertilizers were T₁-Control, T₂-Azotobacter (1%), T₃- Phosphate-solubilizing bacteria (1%), T₄-Azotobacter (2%), T₅- Phosphate-solubilizing bacteria (2%), T₆-Azotobacter (1%) + Phosphate-solubilizing bacteria (1%), T₇- Azotobacter (1%) + Phosphate-solubilizing bacteria (2%), T₈-Azotobacter (2%) + Phosphate-solubilizing bacteria (1%) and T₉-Azotobacter (2%) + Phosphate-solubilizing bacteria (2%) [8]. Number of cuttings in each replication was two. Thus, a total of 54 cuttings were taken as planting materials in this experiment. The shooted

cuttings of three year old plant were collected from the progressive farmer namely "Shri Ram Sharan Verma" at Rasoolpur, Sultanpur. The shooting media was prepared by mixing of sand, soil and FYM with 2:1:1 ratio. Data on growth parameters of dragon fruit in terms of number of days to sprouting, sprouting percentage, number of sprout per cutting, sprout length, shoot length, shoot diameter, number of spines/areole, shoot fresh weight and shoot dry weight were recorded at 30, 60 and 90 days after planting (DAP). Data recorded from the present studies were subjected to analysis by using standard method suggested by Panse and Sukhatme (1967).

3. RESULTS AND DISCUSSION

3.1 Number of Days to Sprouting

Number of days to sprout initiation varied significantly by the influence of bio-fertilizers and their combination. Bio-fertilizer combination contained Azotobacter (2ml) + PSB (1 ml) (T₈) showed significantly less (27.95 days) time for sprout initiation followed by Azotobacter (2 ml) + PSB (2 ml) (28.15 days), whereas the control treatment (T₁) had taken comparatively more (46.24 days) time for sprout initiation. This may be due to increase level of plant growth regulators in the cutting. Therefore, the physiological processes involved in rooting and sprouting of cuttings were completed earlier as a result of the increased amount of auxins (PGR's). "The results are in agreement with the findings of Slankis (1973) who stated that bio-fertilizer raised the concentration of plant growth regulators in plants. Bio-fertilizers has ability to fix or increase the nitrogen content and is vital for cellular processes, growth, electron transport, and photosynthetic rate. It is also a vital source of proteins needed for metabolic processes that occur during growth and development" (Chaplin and Westwood, 1980). Similar results were also reported by Awasthi et al. (2008) in guava and by Minz [9] in dragon fruit cuttings.

3.2 Percent Sprouting

The data pertaining to sprouting percentage as influenced by various concentrations and combinations of bio-fertilizers with at different stages of growth are presented in Table 1. The highest percentage of sprouting were recorded in cuttings treated with treatment (T₈) Azotobacter (2 ml) + PSB (1ml) at 30 and 60 DAP (33.33% and 49.88%, respectively). While the control treatment (T₁) showed the lowest percentage of sprouting (11.44% and 18.22%, respectively).

The ability to produce more sprouts is due to the use of bio-fertilizers, which assisted in the creation of beneficial hormones and growth factors, which in turn increased cell division, cell multiplication, and increased assimilation and accumulation of food resources. Similar results were observed in apple by Raman (2012) and in shea tree by Abdullahi et al. [10]. This finding was also supported by Kaur [11] in Mango.

3.3 Number of Sprouts per Cutting

At 30, 60 and 90 DAP, significant differences were observed on the number of sprouts per cutting among different combinations of bio-fertilizers. Among the bio-fertilizers, the highest numbers of sprouts per cutting (0.75, 1.53 and 2.05) was recorded in Azotobacter (2ml) + PSB (1ml) (T₈) combination at 30, 60 and 90 DAP, respectively). The lowest numbers of sprouts per cutting (0.32, 1.12 and 1.43) were observed at 30, 60 and 90 DAP, respectively) in the control (T₁). The ability to produce more sprouts is due to the use of bio-fertilizers, which assisted in the creation of beneficial hormones and growth factors, which in turn increased cell division, cell multiplication, and increased assimilation and accumulation of food resources. Similar results were observed in apple by Raman (2012) and in shea tree by Abdullahi et al. [10]. This finding was supported by Kaur [11] in mango.

3.4 Shoot Diameter and Length of Sprout

Bio-fertilizers and their combinations had significant influence on the diameter of shoot. The mixture applied with Azotobacter (2 ml) + PSB (1 ml) (T₈) showed the largest diameter of shoot (2.45, 3.15, and 4.12 mm at 30, 60 and 90 DAP, respectively) followed by Azotobacter (T₉) (2.13, 2.99 and 3.99 cm at 30, 60 and 90 DAP, respectively). The largest length of sprout per stem cutting (2.45, 7.01 and 14.75 cm 30, 60 and 90 DAP respectively) was also given by the same treatment (T₈). Statistically similar results were found in T₉ treatment (2.39, 6.36 and 14.25 cm at 30, 60 and 90 DAP). The un-inoculated seedlings (control) had the smallest diameter of shoot (1.05, 1.44, 1.52 mm at 30, 60 and 90 DAP, respectively). The lowest length of sprouts per stem cutting was recorded (1.06, 1.38 and 2.82 cm at 30, 60 and 90 DAP, respectively) in control. The shoot diameter increased due to the uptake of NPK by the plants which was improved by the bio-fertilizers applied. Similarly, length of sprouting increased because of the increased availability of NPK and uptake of other nutrients.

Table 1. Effect of bio-fertilizers on the shoot growth parameters of dragon fruit cuttings

Treatment	Number of days to sprouting	Sprouting percentage			Number of sprout per cutting			Sprout and shoot length (cm)			Diameter of shoot /cutting (cm)			Number of spines/areole			Shoot fresh weight per cutting (g)			Shoot dry weight per cutting (g)		
		30 DAP	60 DAP	90 DAP	30 DAP	60 DAP	90 DAP	30 DAP	60 DAP	90 DAP	30 DAP	60 DAP	90 DAP	30 DAP	60 DAP	90 DAP	30 DAP	60 DAP	90 DAP	30 DAP	60 DAP	90 DAP
T ₁ -Control	16.24	11.44	18.22	0.32	1.12	1.43	1.06	1.38	2.82	1.05	1.44	1.52	3.23	3.64	3.79	9.51	18.46	24.28	1.85	3.77	6.85	
T ₂ - Azotobacter (1%)	14.45	16.45	28.12	0.45	1.15	1.55	1.42	3.45	7.85	1.25	2.15	2.99	3.42	3.98	4.25	12.85	20.15	31.45	3.15	5.45	9.15	
T ₃ - PSB (1%)	13.25	15.25	27.16	0.42	1.14	1.53	1.39	3.25	8.15	1.56	2.25	3.15	3.37	3.87	4.23	11.75	19.85	30.38	2.95	5.25	9.05	
T ₄ - Azotobacter (2%)	13.12	19.33	35.18	0.56	1.25	1.65	1.55	4.55	9.12	2.05	2.89	3.14	3.45	4.01	4.46	12.95	21.25	35.44	3.25	5.85	9.25	
T ₅ - PSB(2%)	11.25	21.27	36.25	0.58	1.28	1.72	1.65	3.75	8.76	1.69	1.78	2.85	3.56	4.12	4.65	13.25	22.15	37.63	3.45	6.12	9.45	
T ₆ - Azotobacter (1%)+ PSB (1%)	12.36	25.47	34.56	0.61	1.35	1.76	1.75	5.75	9.32	2.00	2.26	3.25	3.65	3.99	4.76	14.15	21.85	41.26	3.75	6.45	9.75	
T ₇ - Azotobacter (1%)+ PSB (2%)	9.46	29.38	39.44	0.65	1.51	1.85	1.95	5.25	12.33	2.21	3.01	3.85	3.78	4.24	4.82	14.35	23.65	46.74	3.96	6.88	10.01	
T ₈ – Azotobacter (2%)+ PSB (1%)	7.95	33.33	49.88	0.75	1.53	2.05	2.45	7.01	14.75	2.45	3.15	4.12	3.99	4.56	5.02	14.97	24.69	51.34	4.12	7.48	10.64	
T ₉ - Azotobacter (2%)+ PSB (2%)	8.15	31.68	48.68	0.72	1.48	1.99	2.39	6.36	14.25	2.13	2.99	3.99	3.88	4.45	4.99	14.75	24.25	50.16	4.01	7.36	10.25	
S.Em(±)	0.132	0.325	0.535	0.008	0.027	0.042	0.023	0.062	0.182	0.03	0.04	0.06	0.036	0.047	0.067	0.183	0.296	0.559	0.077	0.087	0.173	
C.D.at 5%	0.399	0.984	1.618	0.025	0.081	0.128	0.070	0.187	0.550	0.10	0.12	0.19	0.110	0.141	0.202	0.552	0.896	1.691	0.233	0.262	0.522	

The results of experiment as conducted by Verma et al. [12] in dragon fruit and Rana et al. (2020) on sweet orange were in accordance with these findings. The beneficial nutrients provided by the bio-fertilizers, caused the diameter of the seedlings to increase. Ganeshnauth et al. (2018) study on pepper plants and obtained the same findings.

3.5 Fresh Weight of Shoot

Shoot fresh weight of dragon fruit cuttings as influenced by different concentrations of bio-fertilizers are furnished in Table 1. The shoot fresh weight per cutting differed significantly at 30, 60 and 90 DAP. The highest fresh weight of shoot was seen in dragon fruit cuttings treated with (T₈) Azotobacter (2 ml) + PSB (1 ml) (14.97, 24.69 and 51.34 g, respectively). The lowest fresh weight was found in (T₁) control (9.51, 18.46 and 24.28 g, respectively).

3.6 Dry Weight of Shoot

The stem cuttings of dragon fruit showed significant differences among the treatments in terms of shoot dry weight. The untreated cuttings showed the least dry weight of shoot at all their growth stages. This may be because of slow sprout initiation, minimum leaf area and smaller number of leaves.

The differences among the treatments may be due to bio-fertilizers which activate shoot growth, resulting in elongation of stems and leaves through cell division accounting for higher dry weight of shoot. The results are in agreement with the findings of Devi et al. [13] in lemon. Among the treatments, application of (T₈) Azotobacter (2 ml) + PSB (1 ml) showed the higher weight of shoot dry matter. This could be due to earliness in sprouting, increase in number of leaves and leaf area and higher shoot fresh weight. Similar results are in conformity with the findings of Devi et al. [13] in lemon cuttings.

4. CONCLUSION

From the experimental results, it may be concluded that among the different treatments of bio-fertilizers either single or in combination have great potential to accelerate shooting in stem cuttings of dragon fruit. Of the treatments, Azotobacter (2%) + PSB (1%) gave better results with respect to shooting parameters followed by the treatment, Azotobacter (2%) + PSB (2%). It is recommended that vegetative method of

propagation through stem cuttings in dragon fruit treated with bio-fertilizers can be reliable for commercial production of planting materials and it is quick method also.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mizrahi Y, Nerd A. New crops as a possible solution for the troubled Israeli export market. 1996;37-45.
2. Panwar S, Singh KP, Namita N. Assessment of variability, heritable components and grouping of Indian rose. The Indian Journal of Agricultural Sciences. 2012;82(10):875-0.
3. Jayant Raman, Jayant Raman. Response of Azotobacter, Pseudomonas and Trichoderma on growth of apple seedling. 2012;83-90.
4. Abobatta WF, El-Azazy AM. Role of organic and biofertilizers in citrus orchards. Aswan University Journal of Environmental Studies. 2020 Oct 1;1(1):13-27.
5. Pushpakumara DK, Gunasena HP, Karyawasam M. Flowering and fruiting phenology, pollination vectors and breeding system of dragon fruit (*Hylocereus spp.*). Sri Lankan J. Agric. Sci. 2005;42:81-91.
6. Chen JH. The combined use of chemical and organic fertilizers and/or biofertilizer for crop growth and soil fertility. In International workshop on sustained management of the soil-rhizosphere system for efficient crop production and fertilizer use. Land Development Department Bangkok Thailand. 2006 Oct 16;16(20):1-11.
7. Pandey L, Verma RS, Shukla KK, Shukla JK, Pathak S. Effect of IBA and NAA on Rooting of Stem Cutting in Dragon Fruit [*Hylocereus undatus* (Haworth) Britton & Rose]. J. Exp. Agric. Int. 2022;44:1-6.

8. Venkata Siva Prasad P, Sivarama Krishna VNP, Venkata Ramana KT, Thanuja Sivaram G, Padmaja VV, Lalitha Kadiri. Studies on combining effect of media and biofertilizers on survival, rooting and establishment of dragon fruit (*Hylocereus costaricensis* L.) cuttings under protected and open field conditions. Pharma Innovation. 2022;11(9):659-663.
9. Minz V. Effect of growing media and plant growth regulators on root and shoot growth of dragon fruit cuttings. M.Sc. (Horti.) Fruit Science thesis, Indira Gandhi Krishi Vishwavidyalaya, Raipur, Chhattishgarh. 2021;1-78.
10. Abdullahi IN, Chuwang PZ, Isah AD. Effect of biofertilizer application on growth of *Vitellaria paradoxa* seedlings. Journal of Research in Environmental Science and Toxicology. 2012 Dec;1(11):294-7.
11. Kour D, Wali VK, Bakshi P, Bhat DJ, Sharma BC, Sharma V, Sinha BK. Effect of organic and inorganic fertilizers along with azotobacter on growth, yield and quality of Aonla (*Emblica officinalis* Gaertn.) Cv. Na-7. Int. J. Curr. Microbiol. App. Sci. 2019;8(9):1142-51.
12. Verma RS, Lata R, Ram RB, Verma SS, Prakash S. Effect of organic, inorganic and bio-fertilizers on vegetative characters of dragon fruit (*Hylocereus undatus* L.) plant. The Pharma Innovation Journal. 2019;8(6):726-8.
13. Devi BL, Krishna VS, Madhumathi C, Yuvaraj KM, Sireesha Y, Krishna MR. Organics and bio-fertilizers effect on germination process of papaya at nursery level (*Carica papaya* L.). Journal of Pharmacognosy and Phytochemistry. 2019;8(1):2425-7.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/123958>