



Synthesis, Characterization and X-Ray Structure of Mononuclear Cu(II) Complex of the Schiff Base 2,2'-[1,1'-(Ethane-1,2-Diyl) Bis (Nitriloethylidyne)]Diphenol

**Gorgui Awa Seck^a, Mamour Sarr^a, Marc Lenertz^b,
Emilie Delahaye^b, Pierre Rabu^b, Mayoro Diop^{a*},
Farba Bouyagui Tamboura^a and Mohamed Lamine Gaye^a**

^a Department of Chemistry, University Cheikh Anta Diop, Dakar, 10700, Senegal.

^b Institute de Physique et Chimie des Matériaux de Strasbourg, Université de Strasbourg, France.

^c NRS UMR 7504, 67034, Strasbourg Cedex 2, France.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/CSIJ/2024/v33i6935>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/127329>

Received: 22/09/2024

Accepted: 25/11/2024

Published: 03/12/2024

Original Research Article

ABSTRACT

The reactions of the Schiff base 2,2'-[1,1'-(ethane-1,2-diyl)bis(nitriloethylidyne)]diphenol (H₂L), which was synthesised by condensation of 2-hydroxyacetophenone and ethylenediamine in the 2:1 ratio, with acetate copper (II) Salt, afforded a new mononuclear complex formulated as

*Corresponding author: E-mail: mayorodiop2000@yahoo.fr;

Cite as: Seck, Gorgui Awa, Mamour Sarr, Marc Lenertz, Emilie Delahaye, Pierre Rabu, Mayoro Diop, Farba Bouyagui Tamboura, and Mohamed Lamine Gaye. 2024. "Synthesis, Characterization and X-Ray Structure of Mononuclear Cu(II) Complex of the Schiff Base 2,2'-[1,1'-(Ethane-1,2-Diyl) Bis (Nitriloethylidyne)]Diphenol". *Chemical Science International Journal* 33 (6):163-73. <https://doi.org/10.9734/CSIJ/2024/v33i6935>.

[Cu(L)].(H₂O)_{0.5}. The ligand and the complex have been characterized by elemental analysis, FTIR, UV-Vis and ¹H and ¹³C NMR spectroscopies. The characterization of the copper complex was completed by molar conductivity and room temperature magnetic measurements. The structure of the complex has been resolved by X-ray crystallography technic. The mononuclear copper (II) complex crystallises in the monoclinic space group C2/c with the following unit cell parameters $a = 26.131(6) \text{ \AA}$, $b = 7.293(5) \text{ \AA}$, $c = 17.256(4) \text{ \AA}$, $\beta = 106.10(5)^\circ$, $V = 3160(2) \text{ \AA}^3$, $Z = 4$, $R_1 = 0.077$ and $wR_2 = 0.145$. In the structure of the complex the ligand acts in tetradentate fashion and the coordination environment of the copper atom is best described as distorted square planar geometry.

Keywords: 2-hydroxyacetophenone; schiff base; copper; mononuclear; FTIR; XRD.

1. INTRODUCTION

“Schiff bases are organic ligands widely studied in coordination chemistry. They are derived from the condensation of amine derivatives and ketone or aldehyde derivatives and constitute a wide range of organic ligands” [1-3]. These ligands, which can have biological or physical properties, are often used in the preparation of transition metal complexes that can enhance the properties of the free ligands” [4-8]. Schiff bases can generate mononuclear [9,10], dinuclear [11,12] or polynuclear complexes [13-15]. “The interest of chemists in these compounds has increased considerably in recent years because of the exceptional properties observed for these molecules. Some coordination compounds derived from Schiff bases have been tested for their biological activities and have been shown to be antibacterial” [16,17], antifungal [18,19], anticancer [20,21], or antituberculosis [22,23]. In these compounds, the presence of several metal ions can induce cooperation between the metal centers and improve physical properties such as magnetism [24], fluorescence [25] or optics Turan et al., [26] and catalytic properties [27,28]. “Schiff bases and their coordination compounds have also been used in the field of biomimetic chemistry to understand the behavior of some metalloenzymes” [29-31]. “In particular, hydroxyacetophenone is a keto-precursor that enables the synthesis of Schiff bases used in the preparation of biomimetic complexes” [32-35]. “In our previous studies, we reported the preparation and use of 2-hydroxyacetophenone-derived Schiff bases to prepare transition metal complexes with novel structures and interesting properties such as antioxidation” [36-39]. This study aims to isolate and characterize a copper(II) Schiff base complex obtained by the condensation reaction between 2-hydroxyacetophenone and ethylenediamine.

2. MATERIALS AND METHODS

2.1 Experimental Analysis

“All chemicals and solvents were of analytical reagent grade and were used directly without further purification. Elemental analyses of C, H and N were recorded on a VxRio EL Instrument. Infrared spectra were obtained on a FTIR Spectrum Two of Perkin Elmer spectrometer in the 4000-400 cm⁻¹ region. The UV-Visible spectra were recorded on a Perkin Elmer Lambda UV-Vis spectrophotometer. The ¹H NMR spectra were recorded at 300 MHz and ¹³C{¹H} NMR spectra at 75 MHz on a Bruker AC-300 instrument. The molar conductance of 1×10⁻³ M in DMF solutions of the metal complex was measured at 25 °C using a WTW LF-330 conductivity meter with a WTW conductivity cell. Room temperature magnetic susceptibility of the powdered sample was measured using a Johnson Matthey scientific magnetic susceptibility balance (Calibrant: Hg[Co(SCN)₄])” [40].

2.2 Synthesis of 2,2'-[1,1'-(ethane-1,2-diyl)bis(nitriloethylidyne)] diphenol (H₂L)

The Schiff base H₂L was synthesized by mixing 2-hydroxyacetophenone (6.02 mL, 0.050 mmol) and ethylenediamine (1.70 mL, 0.025 mmol) in 50 mL of absolute ethanol. The reaction mixture was heated under reflux for four hours. After cooling the mixture at room temperature, yellowish powder appears. The compound was filtered and recrystallized from ethanol [40]. Yield, 83 %, M. P. 199 °C.

2.3 Synthesis of the Copper(II) Complex

In a 100 mL round bottomed flask containing 20 mL of methanol, H₂L (0.2961 g; 1 mmol) was introduced with stirring. Then 10 mL of

Table 1. Crystallographic data and refinement parameter for the complex.

Chemical formula	[2(C ₁₈ H ₁₈ CuN ₂ O ₂)]·H ₂ O
<i>M_r</i>	733.80
Crystal system, space group	Monoclinic, C2/c
Temperature (K)	293
<i>a</i> (Å)	26.131 (6)
<i>b</i> (Å)	7.293 (5)
<i>c</i> (Å)	17.256 (4)
β (°)	106.10 (5)
<i>V</i> (Å ³)	3160 (2)
<i>Z</i>	4
Radiation type	Mo K α
μ (mm ⁻¹)	1.40
Crystal size (mm)	0.19 × 0.08 × 0.05
No. of measured reflections	16310
No. of independent reflections	4504
No. of observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	2135
<i>R</i> _{int}	0.077
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)]	0.077
<i>wR</i> (<i>F</i> ²)	0.145
GOF	1.11
No. of reflections	4504
No. of parameters	215
$\Delta\rho_{\max}, \Delta\rho_{\min}$ (e Å ⁻³)	1.03, -0.72

methanolic solution containing Cu(OAC)₂·4(H₂O) (0.2 g; 1 mmol) was added and the mixture was heated to reflux for two hours. After cooling, the brown solution was filtered, and the filtrate was left to evaporate slowly for two weeks. Brown single crystals suitable for XRD are collected by filtration [40]. Yield, 67%.

2.4 Crystal Structure Determination

“Details of the X-rays crystal structure solution and refinement are given in Table 1. Diffraction data were collected using an ENRAF NONIUS Kappa CCD diffractometer with graphite monochromatized MoK α radiation ($\lambda = 0.71073$ Å). All data were corrected for Lorentz and polarization effects. No absorption correction was applied. Complex scattering factors were taken from the program package SHELXTL [41]. The structures were solved by direct methods which revealed the position of all non-hydrogen atoms. All the structures were refined on *F*² by a full-matrix least-squares procedure using anisotropic displacement parameters for all non-hydrogen atoms” [42]. “The hydrogen atoms of water molecules and NH groups were located in the Fourier difference maps and refined. Others H atoms (CH and CH₃ groups) were geometrically optimized and refined as riding models by AFIX instructions. Molecular graphics were generated using ORTEP-3” [43].

3. RESULTS AND DISCUSSION

3.1 General Study


The symmetrical Schiff base ligand 2,2'-[1,1'-(ethane-1,2-diyl)bis (nitriloethylidene)]diphenol (H₂L) was prepared by refluxing acetophenone and ethylene diamine in 2:1 ratio as shown in Scheme 1. The spectroscopic data are summarized in Table 2. The ¹H NMR spectrum of the ligand exhibits a broad signal pointed at 12.93 ppm attributed to the phenolic protons. The signals of the aromatic protons appear in the range 6.80-7.55 ppm as a complex multiplet. The signals due to the methylene and the methyle protons appear, respectively, at 3.99 ppm and 2.52 ppm. The ¹³C NMR spectrum reveal two signals at 172.39 and 163.63 ppm attributed to the carbon atoms of the azomethine group and the C_{ipso}-OH of the aromatic ring. The signals due to the other aromatic carbon atoms appear in the range 117.49-132.79 ppm. The signals due to the methyle and the methylene carbon atoms are pointed, respectively, at 14.80 and 50.12 ppm. The FTIR of the free ligand shows a broad band at 3423 cm⁻¹ attributed to the $\nu_{\text{O-H}}$ stretching of the phenolic group. The $\nu_{\text{C=N}}$ stretching band is pointed at 1612 cm⁻¹. The $\nu_{\text{C-O}}$ stretching appears at 1285 cm⁻¹. The bands which appear in the range 1550-1450 cm⁻¹ are

attributed to the C=C of the aromatic ring. The electronic spectrum of the free ligand shows two bands pointed at 243 nm and 324 nm which are attributed to the $\pi \rightarrow \pi^*$ and $n \rightarrow \pi^*$ of the phenyl ring and -C=N- bond [44], respectively. The copper complex was synthesized by mixing H₂L ligand in 15 mL of methanol with copper acetate in a molar ratio 1:1. Elemental analyses gives result in agreement with the chemical formula obtained from X-ray diffraction study. The FTIR spectrum of the complex reveals the absence of the band due to ν_{O-H} which appeared in the spectrum of the free ligand at 3423 cm⁻¹. This observation is indicative of the deprotonation of the phenolic groups. The band due to $\nu_{C=N}$ stretching shifts to low frequencies and appears at 1588 cm⁻¹. This fact is indicative of the involvement of the nitrogen atom of the azomethine moiety in the coordination. The band due to ν_{C-O} , which appears at 1285 cm⁻¹ in the

spectrum of the free ligand is pointed at 1220 cm⁻¹, suggesting the coordination of the phenolate oxygen to the copper(II). The electronic spectrum of the complex shows four bands. The bands at 230 and 279 nm are attributed to the $\pi \rightarrow \pi^*$ and $n \rightarrow \pi^*$ of the aromatic ring and the -C=N- bond. The band at 372 nm is due to the MLCT [45]. The electronic spectrum displays also a d→d band at 554 nm which is attributed to ${}^2B_{1g} \rightarrow {}^2A_{1g}$ transition [46]. The magnetic moment value of 1.76 μ_B of the copper(II) complex is in good agreement with paramagnetic copper(II) complexes [47] having one unpaired electron with d⁹ system. The conductimetry measurement (10 S·cm²·mol⁻¹) showed that the copper (II) complex is neutral electrolyte type [48]. On standing for two weeks the conductance remains quite constant, indicating good stability of the complex in DMF solution.

Table 2. Spectroscopic data for the ligand and the complex

H ₂ L	[Cu(L)]·(H ₂ O) _{0.5}		
FTIR (ν (cm⁻¹))			
ν_{O-H}	3423	3458	
$\nu_{C=N}$	1612	1588	
$\nu_{C=C}$	1550-1450	1455-1550	
$\delta_{C=O}$	1285	1220	
UV-Visible (λ (nm))			
$\pi \rightarrow \pi^*$	243	230	
$n \rightarrow \pi^*$	324	279	
MLCT	-	372	
${}^2B_{1g} \rightarrow {}^2A_{1g}$	-	554	
¹H NMR (δ (ppm))		Λ (S·cm²·mol⁻¹)	
δ_{CH_3}	2.52	Fresh solution	10
δ_{CH_2}	3.99	Old solution	12
$\delta_{H_{Ar}}$	6.80-7.55	Magnetic Moment	
δ_{OH}	12.93	μ_{eff} (μ_B)	1.76
¹³C NMR (δ (ppm))			
δ_{CH_3}	14.80	-	
δ_{CH_2}	50.12	-	
$\delta_{C_{Ar}}$	117.49-132.79	-	
$\delta_{C_{Ar-OH}}$	163.63	-	
$\delta_{C=N}$	172.39	-	
Elemental analyses			
% C : Calc. (Found)	72.95 (72.91)	58.92 (58.88)	
% H : Calc. (Found)	6.80 (6.77)	5.22 (5.17)	
% N : Calc. (Found)	9.45 (9.40)	7.64 (7.59)	



Scheme 1. Chemical diagrams for the ligand and the complex

3.2 Description of the Structure of the Complex $[\text{Cu}(\text{C}_{18}\text{H}_{18}\text{O}_2\text{N}_2)] \cdot (\text{H}_2\text{O})_{0.5}$

The molecular structure of the copper(II) complex presented in this work was solved by X-ray diffraction and is illustrated Fig. 1. The main bond lengths and angles are shown in Table 3. The asymmetric unit is composed of one copper(II) ion, one dinegative ligand molecule and a half uncoordinated water molecule. The ligand is coordinated through two nitrogen atoms of azomethine moieties and two phenolate oxygen atoms, yielding a tetracoordinated copper(II) ion which is situated in an N_2O_2 environment. For elucidating the geometry around the metal ion in tetracoordinated complexes, the trigonality parameter τ_4 ($\tau_4 = [360^\circ - (\alpha + \beta)]/141$ (α and β are the largest angles around the metal center) is used: $\tau_4 = 1$ define a perfect tetrahedral geometry and $\tau_4 = 0$ describe a perfect square planar geometry [49]. The τ_4 value of 0.068 is indicative of a distorted square planar geometry around the copper(II) for the title complex. The *cisoid* angles which are in the range $[86.83 (14)^\circ - 92.99 (15)^\circ]$ deviate from the ideal value of 90° . The *transoid* angles values of $174.91(16)^\circ$ $[\text{N}1\text{-Cu}1\text{-O}2]$ and $175.55 (17)^\circ$ $[\text{N}2\text{-Cu}1\text{-O}1]$ deviate from the ideal value of 180° . The sum of the angles subtended by the atoms defining the square planar geometry is 360.39° . The coordinating atoms $\text{N}1/\text{N}2/\text{O}1/\text{O}2$, from the ligand are, approximately, in the same plane (planar rms deviation of 0.0785 \AA) and the

copper atom is located 0.004 \AA out of this plane. Upon coordination of the ligand to the copper(II), one five-membered ring and two six-membered rings were formed. The atoms defining the five membered-ring $\text{Cu}1/\text{N}1/\text{C}9/\text{C}10/\text{N}2$ are coplanar with rms deviation of 0.0835 \AA . The atoms defining the six membered-ring $\text{Cu}1/\text{O}1/\text{C}1/\text{C}6/\text{C}7/\text{N}1$ and $\text{Cu}1/\text{O}2/\text{C}18/\text{C}13/\text{C}11/\text{N}2$ are coplanar with rms deviation of 0.0676 \AA and 0.0462 \AA , respectively. The mean plane of the five-membered ring forms dihedral angle of $5.852(2)^\circ$ and $4.591(2)^\circ$ with the means planes of the two six-membered-rings, respectively. The means planes of the two six-membered-rings form a dihedral angle of $4.472(1)^\circ$. The Cu-N bond lengths $[\text{Cu}1\text{-N}1 = 1.931 (4) \text{ \AA}$ and $\text{Cu}1\text{-N}2 = 1.934 (4) \text{ \AA}]$ are longer than the Cu-O $[\text{Cu}1\text{-O}1 = 1.878 (3) \text{ \AA}$ and $\text{Cu}1\text{-O}2 = 1.876 (3) \text{ \AA}]$ distances. These values are close proximity to the values reported for the complex Chlorido(2-{1-[(2-morpholinoethyl)-imino]ethyl}phenolato- $\kappa^3\text{N,N,O}$)-copper(II) [50]. The bond length of $1.299(5) \text{ \AA}$ for C1-O1 and $1.303(5)$ for C18-O2 are compatible with data reported for C-O_{phenolate} distances [51,52]. Additionally, C7-N1 $[1.286 (5) \text{ \AA}]$ and C11-N2 $[1.292 (6) \text{ \AA}]$ have double bond character and the values of $1.471(5) \text{ \AA}$ and $1.461(5) \text{ \AA}$ for C9-N1 and C10-N2 are indicative of single bond character [53]. No hydrogen-bonds are found in the structure. The packing diagram of the structure is illustrated in Fig. 2.




Fig. 1. Crystal structure of the mononuclear copper (II) complex




Fig. 2. The packing of the complex in the crystal structure

Table 3. Selected geometric parameters (Å, °)

Cu1—O2	1.876 (3)	O1—C1	1.299 (5)
Cu1—O1	1.878 (3)	N2—C11	1.292 (6)
Cu1—N1	1.931 (4)	N2—C10	1.461 (5)
Cu1—N2	1.934 (4)	N1—C7	1.286 (5)
O2—C18	1.303 (5)	N1—C9	1.471 (5)
O2—Cu1—O1	86.83 (14)	O2—Cu1—N2	92.99 (15)
O2—Cu1—N1	174.91 (16)	O1—Cu1—N2	175.55 (17)
O1—Cu1—N1	92.54 (15)	N1—Cu1—N2	88.03 (15)

4. CONCLUSION

The prepared organic ligand H₂L coordinates to copper(II) ion yielding a mononuclear complex. The compounds are characterized by elemental analysis, FTIR, UV-Vis and NMR spectroscopies, room temperature magnetic moment measurement and molar conductivity. The structure of the mononuclear copper (II) complex is established by single-crystal X-ray diffraction. The complex is non-ionic electrolyte in DMF solution. The organic molecule H₂L acts in its dinegative form as tetradentate ligand through two nitrogen azomethine atoms and two phenolate oxygen atoms. Considering the magnetic moment, infrared spectra and the electronic spectrum data for Cu(II) complex, square planar geometry is proposed for the Cu(II) complex as observed in X-ray diffraction study.

SUPPORTING INFORMATION

CCDC-1844618 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via Available: <https://www.ccdc.cam.ac.uk/structures/>, or by, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Haj NQ, Mohammed MO, Mohammad LE. Synthesis and Biological Evaluation of Three New Chitosan Schiff Base Derivatives. ACS Omega. 2020;5(23):13948–13954. Available: <https://doi.org/10.1021/acsomeg.a.0c01342>
2. Mighani H. Schiff Base polymers: Synthesis and characterization. Journal of Polymer Research. 2020;27(6):168. Available: <https://doi.org/10.1007/s10965-020-02080-x>
3. Segura JL, Mancheño MJ, Zamora F. Covalent organic frameworks based on Schiff-base chemistry: synthesis, properties and potential applications. Chem. Soc. Rev. 2016;45(20):5635–5671. Available: <https://doi.org/10.1039/C5CS00878F>
4. Al Zoubi W, Al-Hamdani AAS, Ahmed SD, Ko YG. Synthesis, characterization, and biological activity of Schiff bases metal complexes. Journal of Physical Organic Chemistry. 2018;31(2):e3752. Available: <https://doi.org/10.1002/poc.3752>
5. El-Sonbati AZ, Mahmoud WH, Mohamed GG, Diab MA, Morgan Sh. M, Abbas SY. Synthesis, characterization of Schiff base metal complexes and their biological investigation. Applied Organometallic Chemistry. 2019;33(9):e5048. Available: <https://doi.org/10.1002/aoc.5048>
6. Gul S, Jan F, Alam A, Shakoor A, Khan A, AlAsmari AF, Alasmari F, Khan M, Bo L. Synthesis, molecular docking and DFT analysis of novel bis-Schiff base derivatives with thiobarbituric acid for α -glucosidase inhibition assessment. Scientific Reports. 2024;14(1):3419. Available: <https://doi.org/10.1038/s41598-024-54021-z>
7. Hosny S, Abdelfatah A, Gaber GA. Synthesis, characterization, synergistic inhibition, and biological evaluation of novel Schiff base on 304 stainless steel in acid solution. Scientific Reports. 2024;14(1):470. Available: <https://doi.org/10.1038/s41598-023-51044-w>
8. Venkatesh G, Vennila P, Kaya S, Ahmed SB, Sumathi P, Siva V, Rajendran P, Kamal C. Synthesis and Spectroscopic Characterization of Schiff Base Metal Complexes, Biological Activity, and Molecular Docking Studies. ACS Omega. 2024;9(7):8123–8138. Available: <https://doi.org/10.1021/acsomeg.a.3c08526>
9. Middy P, Sarkar SD, Chattopadhyay S. Use of tridentate NNO donor salicylidine Schiff base ligands for the synthesis of mononuclear cobalt complexes having the general formula [Co(ABC) (DE)X]: Synthetic strategy, structure and application. Journal of Molecular Structure. 2025; 1322:140242.

- Available:<https://doi.org/10.1016/j.molstruc.2024.140242>
10. Paul S, Sanjurani T, Talukdar J, Barman P, Das A, Pandey P. Unraveling the Potential of Mononuclear Zn(II) and Cu(II) Schiff Base Metal Complexes: Microwave-Aided Synthesis, Theoretical Analysis and Application in Biomimetic Catalysis, CT-DNA Interaction, and In Vitro Biological Assays. *Applied Organometallic Chemistry*. 2024;38(12):e7749.
Available:<https://doi.org/10.1002/aoc.7749>
 11. Uyada KAH, Mahesha, Pampa KJ, Harohally NV, Krishnamurthy C, Jathi K, Ahmad A, Alshammari MB, Lokanath NK. Structural Investigation of Schiff Base Ligand and Dinuclear Copper Complex: Synthesis, Crystal Structure, Computational, and Latent Fingerprint Analysis. *ACS Omega*. 2024;9(28):30109–30119.
Available:<https://doi.org/10.1021/acsomega.3c07536>
 12. Huang M, Jiao L, Yang J, Ning D, Xu J, Wu Q, Weng Z. Regulating effect of the halogen atoms on the cofacial dinuclear Schiff base complexes: Synthesis, spectroscopy, electrochemistry, and DFT calculations. *Applied Organometallic Chemistry*. 2024;38(7):e7536.
Available:<https://doi.org/10.1002/aoc.7536>
 13. Chakraborty P., Chatterjee, A., Benmansour, S., Gómez-García, C. J., & Dey, S. K. (2024). Synthesis, characterization and magnetic properties of a hexanuclear copper (II) complex with hydroxido, alkoxido and phenoxido bridges. *Inorganica Chimica Acta*. 2024; 122442.
Available:<https://doi.org/10.1016/j.ica.2024.122442>
 14. Mahapatra TS, Roy BC, Dutta B, Lengyel J, Shatrak M, Ray D. Structures and magnetic properties of a trinuclear angular [Ni₃] and a heptanuclear wheel-like [Ni₇] complexes with a Schiff base ligand. *Polyhedron*. 2024;249:116782.
Available:<https://doi.org/10.1016/j.poly.2023.116782>
 15. Nowicka D, Kubicki M, Patroniak V, Łuczak T, Gorczyński A. Self-assembly of simple Schiff base ligand into unique saddle-type [4x4] tetranuclear architecture and its application as selective voltammetric dopamine sensor in aqueous conditions. *Electrochimica Acta*. 2024;476: 143754.
Available:<https://doi.org/10.1016/j.electacta.2023.143754>
 16. Anacona JR, Santaella J, Al-shemary RKR, Amenta J, Otero A, Ramos C, Celis F. Ceftriaxone-based Schiff base transition metal(II) complexes. Synthesis, characterization, bacterial toxicity, and DFT calculations. Enhanced antibacterial activity of a novel Zn(II) complex against *S. aureus* and *E. coli*. *Journal of Inorganic Biochemistry*. 2021;223:111519.
Available:<https://doi.org/10.1016/j.jinorgbio.2021.111519>
 17. Kargar H, Ashfaq M, Fallah-Mehrjardi M, Behjatmanesh-Ardakani R, Munawar KS, Tahir MN. Unsymmetrical Ni(II) Schiff base complex: Synthesis, spectral characterization, crystal structure analysis, Hirshfeld surface investigation, theoretical studies, and antibacterial activity. *Journal of Molecular Structure*. 2022;1265:133381.
Available:<https://doi.org/10.1016/j.molstruc.2022.133381>
 18. Amirthaganesan K, Vadivel T, Dhamodaran M, Chandraboss VL. *In vitro* antifungal studies of Ruthenium (III) complex derived from chitosan Schiff bases. *Materials Today: Proceedings*. 2022;60:1716–1720.
Available:<https://doi.org/10.1016/j.matpr.2021.12.265>
 19. Frei A, King AP, Lowe GJ, Cain AK, Short FL, Dinh H, Elliott AG, Zuegg J, Wilson JJ, Blaskovich MAT. Nontoxic cobalt(III) Schiff base complexes with broad-spectrum antifungal activity. *Chemistry – A European Journal*. 2021;27(6):2021–2029.
Available:<https://doi.org/10.1002/chem.202003545>
 20. Alorini TA, Al-Hakimi AN, Saeed SES, Alhamzi EHL, Albadri AEAE. Synthesis, characterization, and anticancer activity of some metal complexes with a new Schiff base ligand. *Arabian Journal of Chemistry*. 2022;15(2):103559.
Available:<https://doi.org/10.1016/j.arabj.2021.103559>
 21. Jiang M, Su X, Zhong X, Lan Y, Yang F, Qin Y, Jiang C. Recent development of Schiff-base metal complexes as therapeutic agents for lung cancer. *Journal of Molecular Structure*. 2024;1318:139403.
Available:<https://doi.org/10.1016/j.molstruc.2024.139403>
 22. Bufarwa SM, El-Sefait RM, Thbayh DK, Belaidi M, Al-Shemary RK, Abdusamea RM, El-Ajaily M. M, Fiser B, Bader HA, Saleh AA, Bufarwa MM. Antituberculosis,

- antimicrobial, antioxidant, cytotoxicity and anti-inflammatory activity of Schiff base derived from 2,3-diaminophenazine moiety and its metal(II) complexes: Structural elucidation, computational aspects, and biological evaluation. *Reviews in Inorganic Chemistry*; 2024.
Available:<https://doi.org/doi:10.1515/revic-2024-0007>
23. Malav R, Ray S. Recent developments on the synthesis of copper and cobalt-Schiff base complexes and their assessment as anti-tuberculosis drugs. *Chemical Papers*. 2024;78(8): 4623–4646.
Available:<https://doi.org/10.1007/s11696-024-03425-2>
 24. Chen JT, Yan H, Wang T, Sun WB. Heteronuclear complexes [MDyM] (M=Cu; Zn; Ni) constructed by Schiff base ligands with different amine backbone exhibiting significant single-molecule magnets. *Journal of Molecular Structure*. 20224; 1311:138473.
Available:<https://doi.org/10.1016/j.molstruc.2024.138473>
 25. Dong QW, Yang LZ, Yue TC, Wang DZ, Wang LL. New complexes based on reduced Schiff base carboxylic acid ligand: Electric properties, fluorescence properties and detection of Fe³⁺. *Journal of Molecular Structure*. 20224;1295:136747.
Available:<https://doi.org/10.1016/j.molstruc.2023.136747>
 26. Turan N, Seymen H, Gündüz B, Buldurun K, Çolak N. Synthesis, characterization of Schiff base and its metal complexes and investigation of their electronic and photonic properties. *Optical Materials*. 2024;148:114802.
Available:<https://doi.org/10.1016/j.optmat.2023.114802>
 27. Parada J, Atria AM, Avilés C, Mendoza J, Basualto C, Parra-Melipan S, Zúñiga C, Aguirre P. Recyclable Ni(II), Cu(II), and Co(II) complexes containing Schiff base immobilized on nanosilica as catalysts in oxidation of hydrocarbons. *Inorganica Chimica Acta*. 2024;572:122303.
Available:<https://doi.org/10.1016/j.ica.2024.122303>
 28. Rezaie MA, Khojastehnezhad A, Shiri A. Post-synthetic modification of Zr-based metal organic framework by schiff base zinc complex for catalytic applications in a click reaction. *Scientific Reports*. 2024;14(1):24644.
Available:<https://doi.org/10.1038/s41598-024-76199-y>
 29. Mandal A, Sarkar A, Adhikary A, Samanta D, Das D. Structure and synthesis of copper-based Schiff base and reduced Schiff base complexes: A combined experimental and theoretical investigation of biomimetic catalytic activity. *Dalton Transaction*. 2020;49(43):15461–15472.
Available:<https://doi.org/10.1039/D0DT02784G>
 30. Mureseanu M, Bleotu I, Spînu CI, Cioatera N. Anchoring of Copper(II)–Schiff base complex in SBA-15 matrix as efficient oxidation and biomimetic catalyst. *International Journal of Molecular Sciences*. 2024;25(2).
Available:<https://doi.org/10.3390/ijms25021094>
 31. Shahraki S. Schiff base compounds as artificial metalloenzymes. *Colloids and Surfaces B: Biointerfaces*. 2022;218: 112727.
Available:<https://doi.org/10.1016/j.colsurfb.2022.112727>
 32. Banerjee S, Paine TK. Bioinspired iron(II)-β-diketonate and iron(II)-α-hydroxy ketone complexes of a carbanionic N₃C ligand: Oxidation of metal center vs C-C bond cleavage of co-ligand with dioxygen. *Inorganica Chimica Acta*. 2020;501: 119200.
Available:<https://doi.org/10.1016/j.ica.2019.119200>
 33. Chandra S. Copper (II) complexes with some semicarbazones; Possible biomimetic models. *Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry*. 1984;14(7):957–967.
Available:<https://doi.org/10.1080/00945718408058277>
 34. Mir JM, Roy S, Vishwakarma PK, Maurya RC. *cis*-Dioxomolybdenum(VI) complex of N-*o*-hydroxyacetophenone-isonicotinic acid hydrazide as nosocomial anti-infectious agent: Experimental and theoretical study. *Journal of the Chinese Advanced Materials Society*. 2018;6(3): 282–300.
Available:<https://doi.org/10.1080/22243682.2018.1466727>
 35. Nasr-Esfahani M, Montazerzohori M, Akhlaghi P. Efficient Biomimetic Oxidative Decarboxylation of Some Carboxylic Acids Catalyzed by a Manganese (III) Schiff Base Complex. *Bulletin Korean Chemical Society*. 2009;30(7):1583–1587.

- Available:https://doi.org/chrome-extension://efaidnbmnnnibpcajpcglclefindmkaj/Available:https://web.archive.org/web/20170819211938id_/http://ocean.kisti.re.kr/download/volume/chemical/JCGMCS/2009/v30n7/JCGMCS_2009_v30n7_1583.pdf
36. Fall N, Faye FD, Gaye AA, Diouf O, Gaye M. Synthesis of mono and bis-substituted asymmetrical compounds, 1-(2'-hydroxy-3'-methoxybenzylidene)carbonohydrazide and 1-(2'-hydroxy-3'-methoxybenzylidene)-5-(1'pyridylmethylene)carbonohydrazide: Structural characterization and antioxidant activity study. *Journal of Applied Chemistry (IOSR-JAC)*. 2020;13(12):22–30. Available:<https://doi.org/10.9790/5736-1312022230>
 37. Faye F, Guèye A, Camara PS, Gaye AA, Tamboura FB, Gruber N, Gaye M. Synthesis of Schiff bases compounds from oxamic hydrazide: spectroscopic characterization, X-ray diffraction structure and antioxidant activity study. *American Journal of Applied Chemistry*. 2021;9(1):6–12. Available:<https://doi.org/10.11648/j.ajac.20210901.12>
 38. Kébé M, Thiam IE, Sow MM, Diouf O, Barry AH, Sall AS, Retailleau P, Gaye M. (Hexanuclear copper(II) complex of 2-hydroxy-*N,N*-bis[1-(2-hydroxyphenyl)ethylidene]propane-1,3-diamine incorporating an open-cubane core. *Acta Crystallographica Section E*. 2021;77(7):708–713. Available:<https://doi.org/10.1107/S2056989021005570>
 39. Mamour S, Mayoro D, Elhadj Ibrahima T, Mohamed G, Aliou Hamady B, Ellena J. (.{1-[1-(2-Hydroxyphenyl)ethylidene]-2-(pyridin-2-yl- κ N)hydrazine- κ^2 N,O}]-[1-(2-oxidophenyl)ethylidene]-2-(pyridin-2-yl- κ N)hydrazine- κ^2 N,O}nickelate(II) nitrate hemihydrate. *Acta Crystallographica Section E*. 2018;74(5):642–645. Available:<https://doi.org/10.1107/S2056989018005261>
 40. Kumar C. Synthesis, characterization and biological evaluations of heterobinuclear complexes of Cu^{II} and Ni^{II} Schiff base of ethylenediamine and o-hydroxyacetophenone with alkali metal salts of o-nitrophenol. *Journal of Indian Chemical Society*. 2017;94:857-861. Available:<https://doi.org/10.5281/zenodo.5595126>
 41. Sheldrick GM. Crystal structure refinement with SHELXL. *Acta Crystallographica Section C*. 2015a;71(1):3–8. Available:<https://doi.org/10.1107/S2053229614024218>
 42. Sheldrick GM. SHELXT – Integrated space-group and crystal-structure determination. *Acta Crystallographica Section A*. 2015b;71(1):3–8. Available:<https://doi.org/10.1107/S2053273314026370>
 43. Farrugia LJ. WinGX and ORTEP for Windows: An update. *Journal of Applied Crystallography*. 2012;45(4):849–854. Available:<https://doi.org/10.1107/S0021889812029111>
 44. Kendel A, Miljanić S, Kontrec D, Soldin Ž, Galić N. Copper(II) complexes of aroylhydrazones: Preparation and structural characterization. *Journal of Molecular Structure*. 2020;1207:127783. Available:<https://doi.org/10.1016/j.molstruc.2020.127783>
 45. Saha S, Sasmal A, Choudhury CR, Gómez-García CJ, Garribba E, Mitra S. A new linear double phenoxide-bridged trinuclear Cu(II) Schiff base complex: Synthesis, crystallographic elucidation, magneto-structural correlation and DFT Study. *Polyhedron*. 2014;69:262–269. Available:<https://doi.org/10.1016/j.poly.2013.12.006>
 46. Dey L, Rabi S, Hazari SKS, Roy TG, Buchholz A, Plass W. Copper(II) and nickel(II) complexes of an N-pendent bis-(cyanoethyl) derivative of an isomeric hexamethyl tetraazamacrocyclic ligand: Synthesis, characterization, electrolytic behavior and antimicrobial studies. *Inorganica Chimica Acta*. 2021;517:120172. Available:<https://doi.org/10.1016/j.ica.2020.120172>
 47. Srivastava AK, Yadav P, Srivastava K, Prasad J. Synthesis, characterization, biological and electrochemical investigation of copper (II) complexes containing 4-chloro-2-[2, 6-diisopropylphenylimino] methyl] phenol Schiff base ligand and aromatic diinines. *Chemical Data Collections*. 2021;32:100659. Available:<https://doi.org/10.1016/j.cdc.2021.100659>
 48. Geary WJ. The use of conductivity measurements in organic solvents for the characterisation of coordination

- compounds. Coordination Chemistry Reviews. 1971;7(1):81–122.
Available:[https://doi.org/10.1016/S0010-8545\(00\)80009-0](https://doi.org/10.1016/S0010-8545(00)80009-0)
49. Singh YP, Patel RN, Singh Y, Choquesillo-Lazarte D, Butcher RJ. Classical hydrogen bonding and stacking chelate rings in new copper(ii) complexes. Dalton Trans. 2017;46(9):2803–2820.
Available:<https://doi.org/10.1039/C6DT04661D>
50. Ikmal Hisham NA, Suleiman Gwaram N, Khaledi H, Mohd Ali H. Chlorido(2-{1-[(2-morpholinoethyl)imino]ethyl}phenolato- κ^3N,N',O)copper(II). Acta Crystallographica Section E. 2011;67(1):m57.
Available:<https://doi.org/10.1107/S1600536810051160>
51. Zhang J, Chen XD, Zhang HH, Sun BW. Bis{ μ -2-[1-(2-Pyridylmethylimino)ethyl]phenolato}bis-[azidocopper(II)]. Acta Crystallographica Section E. 2009;65(9):m1136.
Available:<https://doi.org/10.1107/S1600536809030475>
52. Zhao GQ, Xue LW, Hao CJ, Chen LH, Wu HT. [Bis(3,5-dimethylpyrazol-1-yl)methane]{N-[1-(2-oxidophenyl)ethylidene]-DL-alaninato}copper(II) monohydrate. Acta Crystallographica Section E. 2008;64(12):m1553.
Available:<https://doi.org/10.1107/S1600536808037264>
53. Seck TM, Faye FD, Gaye AA, Thiam IE, Diouf O, Gaye M, Retailleau P. Synthesis of mono and bis-substituted asymmetrical compounds, (1-(pyridin-2-yl)ethylidene) carbonohydrazide and 1-(2'-hydroxybenzyl idene) -5-(1'-pyridylethylidene) carbonohydrazone : Structural characterization and antioxidant activity study. European Journal of Chemistry. 2020;11(4): Article 4.
Available:<https://doi.org/10.5155/eurjchem.11.4.285-290.2023>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/127329>