



Impact of Selected Soil Physical and Chemical Properties Regulating Phosphate Availability in Hadejia-Jama'are Wetland Soils

Shehu, Yusuf ^{a*}, James, Patricia Aliyu ^b, Solomon, Rejoice Ibrahimn ^c, Musa, Sharhabil Yahaya ^a and Joseph, Jerry ^d

^a Department of Soil Science, Ahmadu Bello University, Zaria, Nigeria.

^b Department of Soil Science and Land Resources Management, Federal University, Wukari, Taraba State, Nigeria.

^c Soil Science Department, Modibbo Adama University, Yola, Adamawa State, Nigeria.

^d Chemistry Department, Kaduna State College of Education Gidan Waya, Kafanchan, Kaduna State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.56557/ajopss/2024/v9i1108>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://prh.ikpress.org/review-history/12249>

Original Research Article

Received: 14/08/2024

Accepted: 17/10/2024

Published: 06/12/2024

ABSTRACT

The objectives of this study were to examine the major soil routine parameters regulating P availability and transformation in Hadejia-Jama'are wetlands. Three wetland locations; Sinamu, Tandanu and Masama and their corresponding adjacent dryland soils as control were strategically selected. Six profile pits were dug (one each on the six research wetlands). Triplicate soil samples

*Corresponding author: E-mail: yusufshehu2@gmail.com;

Cite as: Yusuf, Shehu, James, Patricia Aliyu, Solomon, Rejoice Ibrahimn, Musa, Sharhabil Yahaya, and Joseph, Jerry. 2024. "Impact of Selected Soil Physical and Chemical Properties Regulating Phosphate Availability in Hadejia-Jama'are Wetland Soils". *Asian Journal of Plant and Soil Sciences* 9 (1):74-85. <https://doi.org/10.56557/ajopss/2024/v9i1108>.

were collected strategically at 0-35, 35-70, 70-105 cm soil depths. The following standard laboratory procedures were used; Particle size distribution (soil organic carbon, Electrical conductivity, effective cation exchange capacity (ECEC) by summation, pH, available phosphorus. The clay content of the soils manifest decrease from top to bottom except in Sinamu and Tandanu upland soils. The pH of the wetland soils varied between strongly acidic (pH 4.1) to moderately acidic (pH 5.5) while that of the drylands varied between moderately acidic (pH 5.3) to slightly acidic (pH 6.8). There is significant difference in soil solution pH and exchangeable acidity between the two land forms $P = .05$. The wetland have higher EA values 0.9 to 1.3 $\text{cmol}(+)\text{kg}^{-1}$ compared to dryland soils 0.4 to 0.6 $\text{cmol}(+)\text{kg}^{-1}$. There was a significant difference Mg^{2+} content between the wetlands and the drylands ($P = .05$). The values of Potassium ions recorded were generally very low both in the wetlands 0.010 to 0.026 $\text{cmol}(+)\text{kg}^{-1}$ and the drylands 0.02 to 0.036 $\text{cmol}(+)\text{kg}^{-1}$. There was significantly difference $P = .05$ between the wetland and dryland soils with the dryland soils having higher Ca^{2+} ions. The difference in effective cation exchange capacity (ECEC) between the wetland and the dryland soils was statistically significant $P = .05$ with the dryland soils having higher ECEC values. there was significant difference in ECEC between the surface soils 0 to 35cm and the sub- surface soils 35cm up to 105 cm $P = .05$. The available P content of the two land forms were significantly different $P = .05$. Base on the findings above, the soil parameters regulating P availability and transformation in both Hadejia-Jama'are wetland and dryland soils in order of increasing magnitude are silicate clays, pH, O.C, ECEC, EA, Ca and Mg.

Keywords: Wetlands; phosphate; clay content; organic matter; pH.

1. INTRODUCTION

The reaction types that fix P ions in soils varies from soil to soil and have strong connection to soil solution pH. High P fixation occur at a very low or very high soil solution pH [1]. The concentrations of H_2PO_4^- and HPO_4^{2-} in soil largely depends on the soil solution pH [2]. Soil clay content affect P sorption by attracting P ions onto smaller clay particles such as free Fe and Al oxides, thus soils with high clay content tend to have high P fixing capacity [3]. Soil organic matter (O.M) greatly influences P sorption-desorption through its binding effect [4]. The chelating ability of the naturally synthesized oxyl, hydroxyl and carboxyl ligands synthesized by microbes from soil organic matter can affect phosphate availability by blocking some of the reactive sites on the soil matrix surfaces thereby reducing phosphate adsorption [5]. Humic acids on their own have the ability to inhibit crystallization and polymerization of Fe and Al oxides [6]. High clay content and high O.M in soil have the potential to increase the soil cation exchange capacity CEC [3]. The greatest problem of available phosphorus (P) in soil is that P can be sufficiently present yet un-available for plants uptake due to strong covalent bond formed with aluminum/iron oxides and/or undergo precipitation reaction with calcium and magnesium ions in the soils readily. Lack of knowledge about this problem by farmers, makes lots of farmers to apply excess amount of P

source in an attempt to replenish the soil P in order to achieve bumper harvest which in turn leads to eutrophication of the wetland water bodies and rapid growth of invasive species of grasses that claims many of the suitable portions of the wetlands used for crops cultivation. In situations where this problem exist, application of sufficient organic matter can help solve or at least reduce the problem to a minimum level, otherwise liming can be recommended if the problem is severe. As at when this research was carried out, no research was conducted to specifically examine the parameters controlling P availability in Hadejia-Jama'are wetland and its dryland soils. Most of the established researches [7]. limit their findings to pedogeochemical assessment of the Hadejia- Jama'are wetlands which is non-specific to P and/or parameters controlling P availability in soils, [8] limit their findings to exchangeable bases of the Hadejia-Jamaare wetlands, [9]. limit their findings to land use and Mohammed, et al. [10]. limit their findings to statistical assessment of pH, exchangeable bases and base saturation of Hadejia-Nguru wetland soils.

2. MATERIALS AND METHODS

2.1 Site Description

Three wetland locations, namely; northern wetlands, western wetlands, southern wetlands of Hadejia-Jama'are river basin and their corresponding adjacent drylands as control were

strategically selected. The sampling locations and coordinates of the sampling point were; Sinamu wetland at 12.450°N and 10.044°E, Sinamu dryland at 12.455°N and 10.054°E, Tandanu wetland at 12.405°N and 10.100°E, Tandanu dryland at 12.452°N and 10.110°E, Masama wetland at 12.447°N and 10.130°E, Masama dryland at 12.501°N and 10.111°E.

2.2 Sampling

A total of six profile pits were sank, one on each of the six research wetlands.

Triplicate undisturbed and disturbed soil samples were collected at each of the selected soil depths 0-35, 35-70, 70-105 cm. The soil Samples collected were immediately transferred into air-sealed plastic bags, labeled appropriately and transported to the laboratory.

2.3 Treatment of the Samples

The undisturbed soil samples collected were used for bulk density analysis while the disturbed soil samples were air dried, crushed, sieved through 2mm mesh sieve, labeled and stored in polythene lathers for particle-size-distribution and further laboratory chemical analyses. Grade analytical reagents and distilled water were used all through the laboratory experiments. All the glass wares and plastic containers used were washed with detergent, swirl with nitric acid, then rinsed very well with distilled water.

2.4 Analytical Procedures/ Instrumentations

The following standard laboratory procedures were employed; Bulk density analysis [11]. Particle size distribution was determined by the Bouyoucus standard hydrometer method described by Uyovbisere, et al. [12]. Soil organic carbon content was determined using Walkley-Black method described by SSSA, [13] and calculated using the formula:

$$\% \text{ OC} = \frac{\text{Blank} - \text{Titter} \times \text{NF} \times 0.003 \times \text{CF}}{\text{Weight of the soil sample used} \times 100} \text{ and finally converted to } \text{gkg}^{-1} \text{ as } \% \text{ OC} \times 10.$$

Where NF stands for normality of ferrous sulphate and CF stands for correction factor.

The soil pH was determined using Advance (m/v) digital pH meter in 1:2.5 soil/deionized water described by Brevick, and Burgess, [14]. Effective cation exchange capacity (ECEC) was

determined using ammonium acetate method described by Soil Survey Staff, [13]. Electrical conductivity was determined in 1:2.5 soil/water ratio using Rhoades, and Manteghi, [15] procedure. Available P was determine using Bray 1 procedure described by Sparks, [16]. Data obtanine was subjected to ANOVA using Statistical Analysis System SAS [17]. Statistical Software package version 9.3.

3. RESULTS AND DISCUSSION

The clay content and the textural class (Tables 1 and 2) recorded in the wetland soils ranged between 10 to 28 % clay while in the dryland soils a range of 9 to 12 % clay was recorded. The textural class of the wetland soils varied from silt loam, loam, loamy sand, sandy loam and clay loam while that of the dryland soils were dominantly sandy loam and loam. Out of the eighteen soil samples, eight were classified as loamy soils, seven sandy loam, one silt loam, one loamy sand and one clay loam. The clay content recorded in the wetlands reveal higher values compared to the drylands, having dominant clay values between 14 and 28 % (Table 2). The dryland soils manifest very low clay content, having dominant clay values between 9 to 12 % clay. Consistent increase in clay content down the soil profile was observed in all the wetlands except the northern wetland. On the other hand, the dryland soils manifest consistent decreases from surface to sub- surface except in southern dryland. A significant difference $P = .05$ in clay content (Table 2) between the wetland and the dryland soils was observed and the highest mean value was recorded in the wetland soils. The bulk density values across the three wetlands were found slightly higher 1.40 to 1.56 gcm^{-3} compared to those recorded in the dryland 1.40 to 1.54 gcm^{-3} revealing the true nature of a wetland's geology. The BD from 70 cm downward manifest some evidence of compaction. However, values reported were in conformity with those recorded by Aliyu et al. [7] 1.53 gcm^{-3} in Hadejia-Jama'are wetland soils. The Ph of the wetland soils (Table 2) varied between strongly acidic (Ph 4.1) to moderately acidic (Ph 5.5) while the dryland Ph varied between moderately acidic (Ph 5.3) to slightly acidic (Ph 6.8). Masama upland appeared to be more acidic compared to Sinamu and Tandanu uplands. The wetland soils also manifest similar trend, suggesting that, soil properties down-slope are usually a reflection of the soil properties up- slope. The wetland soils are generally acidic compared to the dryland soils, indicating abundance of soluble aluminum

and hydrogen ions and probably manganese in that pedo-environment. This might be as a result of anaerobic condition of the wetlands and the lithologic origin of the parent material from which weatherable materials were washed, transported and deposited down-slope. All the drylands show decrease in pH down the soil profile except in Tandanu dryland. Aliyu, et al. [7] reported a pH of 4.88 in Hadejia-Jama'are wetland soils which corroborate with the values recorded in this work. Shehu et al. [18] obtained a pH range of 5.0 to 5.5 in Sudan savanna soils which also corroborate with the values observed in the upland soils in this work. There is however, significant difference in soil solution pH (Table 3) between the two land forms $P = .05$, suggesting dominance of soluble aluminum in the wetland soil solution. There was significant difference in soil solution pH between the surface (0-35cm) and subsurface (70-105cm) but the two selected soil depths mentioned above were statistically the same with the sub-surface (35 to 70 cm) soil depth at $P = .05$. This may be attributed to leaching of soluble products within 0 to 70 cm soil depth. There is significant interaction in pH between the land forms and the selected soil depths (Table 5).

3.1 Exchangeable Acidity (EA)

The wetland soils appeared to have higher EA values 0.9 to 1.3 $\text{cmol}(+)\text{kg}^{-1}$ compared to dryland soils 0.4 to 0.6 $\text{cmol}(+)\text{kg}^{-1}$. Similarly, there is a significant difference ($P = .05$) (Table 3) in exchangeable acidity between the two land forms probably due to protonation by hydrogen ions from water molecule and soluble aluminum ions released from the parent rock, trapped and incorporated into the soil and became part of the clay mineralogy, there by changing the composition of the soil-mineral. All the wetlands showed increasing E.A down the soil profile except southern wetland that manifest an inconsistent pattern. Shehu et al. [18] reported values $< 1.0 \text{ cmol}(+)\text{kg}^{-1}$ in Nigeria Sudan savanna which agree with the values recorded in this work.

3.2 The Exchangeable Sodium (Na^+)

Exchangeable sodium concentration of the wetland soils (Table 2) ranged from 0.87 to 1.24 $\text{cmol}(+)\text{kg}^{-1}$ while values obtained in the dryland soils ranged from 0.16 to 1.08 $\text{cmol}(+)\text{kg}^{-1}$. The soils have no evidence of sodicity, since for a soil

to be considered sodic it must have exchangeable sodium percentage (ESP) $>15\%$, $\text{EC} < 4 \text{ (dSm}^{-1}\text{)}$, $\text{pH} \geq 8.2$ and/or manifest visible effects on the soil structure FAO, [19]. Shehu et al. [18] reported very low values (0.1 to 0.3 $\text{cmol}(+)\text{kg}^{-1}$) in Nigerian Sudan Savanna soils. Yakubu, [20] reported 0.01 to 0.12 $\text{cmol}(+)\text{kg}^{-1}$ in Nigerian Guinea Savanna. All the values cited here were low compared to the values reported in this work.

3.3 The Exchangeable Magnesium (Mg^{2+})

The exchangeable magnesium concentration recorded in the dryland soils is higher 0.22 to 0.51 $\text{cmol}(+)\text{kg}^{-1}$ compared to 0.38 to 2.65 $\text{cmol}(+)\text{kg}^{-1}$ observed in the wetland soils and there is a significant difference ($P = .05$) Mg^{2+} content between the wetland and the dryland soils (Table 3). Values obtained in both locations especially in the dryland soils were between low to moderate, since values < 0.3 even up to 0.4 $\text{cmol}(+)\text{kg}^{-1}$ can still be considered low (Leo et al. n.d.; Daniel et al. [21]. Shehu et al. [18] reported a range of 0.3 to $> 1.0 \text{ cmol}+\text{kg}^{-1}$. some of the values he reported agree with the values recorded in this work.

3.4 The Exchangeable Potassium (K^+)

Values obtained were generally very low both in the wetlands 0.010 to 0.026 $\text{cmol}(+)\text{kg}^{-1}$ and 0.02 to 0.036 $\text{cmol}(+)\text{kg}^{-1}$ in the drylands, attributed to non-exchangeable behavior of K^+ , often trapped in the inter-layer of clay minerals, such that only very little amount is freely available. Shehu et al. [18] reported very low values (< 0.15) which were still higher compared to the values obtained in this work. Values reported in Hadejia-Jamaare wetlands 0.41 to 16.07 mg kg^{-1} by Goni et al. [8] were extremely low compared to the values recorded in the same wetland soils of Hadejia-Jama'are river basin, probably associated to the difference in seasons within the year when the soil sampling was established. The K^+ content between soil depth 35 to 70 cm and 70 to 105 cm were statistically the same ($P = .05$) but significantly different with the K^+ content at soil depth 0 to 35 cm. The K fixation in the interlayer of the clay minerals is probably high at soil depth 35 to 105 cm than at soil depth 0 to 35 cm.

3.5 The Exchangeable Calcium (Ca^{2+})

The Ca^{2+} concentration in the dryland soils appeared to be much higher 0.35 to 0.77

cmol(+)kg⁻¹ compared to 1.11 to 8.20 cmol(+)kg⁻¹ in the wetland soils despite high clay content was observed in the wetland soils. Probably due to low EA and high pH observed in the dryland soils. Abdu and Udofot and Abdu, [22] reported that, Ca is the dominant cation in the exchange site of savanna drylands. Shehu et al. [18] reported 2.0 to 5.0 cmol(+)kg⁻¹. There is significantly difference $P = .05$ between the wetland and dryland soils with the dryland soils having higher Ca²⁺. The Ca²⁺ content between the soils of Sinamu and Masama locations were statistically at par $P = .05$ but significantly different from the Tandanu research location. The distribution pattern down the soil profile showed no significant difference $P = .05$ between the two soil depths 0-35 cm and 70-105 cm but was significantly different from 35 to 70 cm soil depth $P = .05$ indicating presence of a bleached zone somewhere within 35 to 70 cm soil depth, attributed to co-migration of Ca with clays especially in the wetland soils during the period of heavy downpour usually in the month of August and due to continuous push by irrigation water during the dry season irrigation activities by farmers. There was significant interaction between landforms and research locations (Table 4) The highest ranked Ca²⁺ interaction between land form and research locations (Table 4) was recorded at Tandanu upland but it wasn't significantly different from values recorded in Sinamu and Masama dryland soils. However, it was significantly different from Ca²⁺ values obtained due to the effect of interaction between landform and research locations in all the wetland soils. Similarly, effect of interaction recorded at Sinamu and Masama drylands was statistically the same with the Ca²⁺ value recorded in all the wetland soils, attributed to sharp variation in calcium content between the two land forms, across the research fields and down the soil sampling depths.

3.6 The Effective Cation Exchange Capacity (ECEC)

The ECEC (Table 2) ranged from 2.68 to 3.40 cmol(+)kg⁻¹ in the wetland soils and 2.52 to 11.10 cmol(+)kg⁻¹ in the dryland soils. Values observed in the dryland soils appeared to be higher compared to values observed in the wetland soils. High concentration of exchangeable calcium recorded in the dryland soils may explain this difference. The soil ECEC recorded, show increase from surface to sub-surface in the drylands except in Tandanu dryland while in the

wetland soil, the distribution is almost uniform with little variation from 0 up to 105 cm soil depth. Values up to 7.22 cmol(+)kg⁻¹ were reported by Aliyu et al. [7]; Umar et al. [9] in wetland soils. The difference in effective cation exchange capacity (ECEC) between the wetland and the dryland soils (Table 3) was statistically significant $P = .05$, with the dryland soils having higher ECEC values attributed to abundance of Ca ions as earlier observed in the soil physicochemical properties, which contribute greatly to the ECEC values compared to contribution by other exchangeable cations, indicating high tendency for P precipitation in the drylands by Ca ions under high pH than there will be in the wetlands. However, across the three research locations, the ECEC in Sinamu and Tandanu soils were statistically the same but significant difference exist between.

Masama and the two other locations above all at $P = .05$, having Sinamu and Tandanu soils with higher ECEC values compared to Masama soil, suggesting that, Masama soils may have probably been formed from a different parent material, different from Sinamu and Tandanu soils. It may probably be attributed to calcium loading from calcium phosphate fertilizer added by farmers or from organic manure added by farmers prior and or during farming activities. There was no significant difference in ECEC from 70 cm up to 105 cm soil depth, but there was significant difference in ECEC between the surface soil 0 to 35cm and the sub-surface soil 35cm up to 105cm $P = .05$, this can be linked to difference in O.M content which is usually high in the surface soils of most soils than there is in the sub-surface soils. There was significant interaction between land forms and research locations (Table 3) attributed to profound variations revealed by clay content, organic carbon content, exchangeable acidity and exchangeable basis (Table 2). Highest ranked ECEC value due to interaction between land forms and research locations (Table 4) was recorded at Tandanu wetland but it was not significantly different from the value recorded at Sinamu wetland. However, it was significantly different from ECEC values obtained between land forms and locations at all the drylands and at Masama wetland. This may in addition be attributed to high E.A recorded in Tandanu and Sinamu wetlands which leads to serious increase in the ECEC of these two locations over Masama.

Table 1. Particle-Size Distribution of Hadejia-Jama'are Wetlands and Dryland Soils

Sampling Locations	Depth (cm)	% Clay	% Silt	% Sand	Textural class
SINAMU	0-35	14.00	WETLANDS 51.00	35.00	Silt loam
	35-70	14.00	35.00	52.00	Loam
	70-105	12.00	38.00	50.00	Loam
TANDANU	0-35	26.00	31.00	43.00	Loam
	35-70	16.00	32.00	52.00	Loam
	70-105	13.00	25.00	62.00	Loamy sand
MASAMA	0-35	28.00	28.00	44.00	Clay loam
	35-70	20.00	22.00	58.00	Sandy loam
	70-105	10.00	39.00	51.00	Loam
SINAMU	0-35	11.00	DRYLANDS 37.00	52.00	Loam
	35-70	10.00	21.00	69.00	Sandy loam
	70-105	9.00	32.00	59.00	Sandy loam
TANDANU	0-35	11.00	29.00	60.00	Sandy loam
	35-70	11.00	30.00	59.00	Sandy loam
	70-105	10.00	38.00	52.00	Loam
MASAMA	0-35	12.00	38.00	50.00	Loam
	35-70	9.00	24.00	67.00	Sandy loam
	70-105	10.00	15.00	75.00	Sandy loam

Table 2. Selected Soil Physical and Chemical Properties of Hadejia-Jama'are Wetlands

Loc.	Depth(cm)	BD (gcm ⁻³)	Clay %	T/class	pH CaCl ₂	EA	Na ⁺	Mg ²⁺ (cmol(+)k g ⁻¹)	K ⁺	Ca ²⁺	ECEC	EC(dSm ⁻¹)	Av.P (mgkg ⁻¹)	OC (gkg ⁻¹)
SN	0-35	1.45	14	SiL	5.0	0.9	1.02	0.34	0.017	0.35	2.68	1.02	19.6	5.2
	35-70	1.46	14	L	5.1	0.9	1.03	0.40	0.017	0.77	3.17	1.21	14.2	4.1
	70-105	1.49	12	L	4.8	1.0	0.97	0.31	0.010	0.51	2.79		12.4	2.9
TA	0-35	1.54	26	L	5.1	1.3	1.24	0.35	0.011	0.35	3.45	1.02	16.2	7.7
	35-70	1.54	16	L	5.1	1.1	0.90	0.43	0.026	0.71	3.26	0.92	14.6	4.6
	70-105	1.44	13	LS	4.4	1.3	1.14	0.22	0.019	0.55	3.00	0.89	10.5	2.9
MA	0-35	1.56	28	CL	4.1	1.3	1.10	0.32	0.012	0.35	3.09	0.99	13.4	2.5
	35-70	1.40	20	SL	5.3	0.9	0.90	0.41	0.017	0.66	3.04	0.86	9.4	1.9
	70-105	1.54	10	L	4.1	1.1	0.87	0.33	0.018	0.60	2.08	0.87	8.0	1.5

Loc.= Location, SIN=Sinamu, TA =Tandanu, MA = Masama, SiL = Silt Loam, SL = Sandy Loam, LS = Loamy Sand, L = Loamy Soil, CL = Clay Loam

Table 3. Contened Selected Soil Physical and Chemical Properties of Hadejia-Jama'are Drylands

Loc	Depth (cm)	Bd (gcm ⁻³)	Clay %	T.class	pH CaCl ₂	EA	Na ⁺	Mg ²⁺ (cmol(+)kg ⁻¹)	K ⁺	Ca ²⁺	ECEC	EC (dSm ⁻¹)	A.P (mgkg ⁻¹)	O.C gkg ⁻¹
SN	0-35	1.45	11	L	6.8	0.4	0.93	0.76	0.036	3.18	5.33	1.81	16.6	5.6
	35-70	1.40	10	SL	5.6	0.5	1.07	0.73	0.028	3.14	5.52	1.91	10.6	4.6
	70-105	1.46	9	SL	5.5	0.5	0.16	0.57	0.002	1.29	2.52	1.24	12.1	3.9
TA	0-35	1.49	11	SL	6.0	0.4	0.96	0.38	0.035	5.50	8.29	2.11	13.5	5.8
	35-70	1.45	11	SL	6.1	0.5	1.04	0.75	0.031	8.80	11.10	2.01	11.5	5.6
	70-105	1.54	10	L	6.2	0.4	1.08	1.08	0.002	6.20	8.76	1.18	10.0	5.0
MA	0-35	1.46	12	L	5.6	0.5	1.05	1.18	0.036	1.11	4.00	1.95	16.0	5.0
	35-70	1.54	9	SL	5.5	0.6	0.93	0.60	0.028	2.44	4.65	1.10	10.9	4.6
	70-105	1.42	10	SL	5.5	0.4	0.99	2.65	0.002	1.44	5.48	1.00	11.1	2.9

Loc. = Location, SN = Sinamu, TA = Tandanu, MA = Masama, LS = Loamy Sand, L = Loamy Soil,

Table 4. Effect of Treatments on Selected Soil Physical and Chemical Properties in Hadejia-Jama'are River Basin

Trt.	Bd (gcm ⁻³)	Clay gkg ⁻¹	pH CaCl ₂	EC (dSm ⁻¹)	OC (gkg ⁻¹)	TN	ECEC	E.A (cmol(+)kg ⁻¹)	Na+	K+	Ca ²⁺	Mg ²⁺	Av.P (mgkg ⁻¹)
Land forms													
Wetlands	1.492	16.67 ^a	4.68 ^b	1.23	4.8	2.83	4.29 ^b	2.081 ^a	1.01	0.019	0.94 ^b	0.49 ^b	13.19 ^a
Drylands	1.469	10.33 ^b	6.03 ^a	2.20	3.7	3.14	8.82 ^a	0.993 ^b	0.91	0.022	2.68 ^a	1.37 ^a	11.70 ^b
LSD	0.0595	5.171	1.03	1.44	1.4	0.81	1.197	0.79	0.23	0.0095	0.82	0.35	1.205
Location													
Sinamu	1.453	14.88	5.38	2.03	4.4 ^{ba}	2.85	6.02 ^a	1.230	0.86	0.018	1.04 ^b	0.54	14.25 ^a
Tandanu	1.500	14.00	5.73	1.21	5.3 ^a	2.85	7.52 ^a	1.138	1.04	0.024	3.19 ^a	0.67	12.88 ^{ab}
Masama	1.488	11.07	5.38	1.20	3.1 ^b	3.27	4.63 ^b	1.023	0.97	0.018	0.90 ^b	0.54	11.70 ^b
LSD	0.073	0.33	0.59	2.04	1.7	0.99	2.66	1.335	0.28	0.012	0.99	1.65	1.333
Depth (cm)													
0-35	1.492	15.83	5.92 ^a	2.18	3.8	3.87 ^a	5.48 ^a	1.535	0.87	0.009 ^b	1.60 ^b	1.38	15.55 ^a
35-70	1.468	13.33	5.70 ^{ab}	2.20	4.3	2.63 ^b	3.06 ^{ab}	0.822	0.98	0.025 ^a	2.92 ^a	0.60	11.87 ^b
70-105	1.482	11.33	5.20 ^b	1.22	4.7	2.47 ^b	4.62 ^a	1.035	1.03	0.028 ^a	1.31 ^b	0.56	11.42 ^b
LSD	0.073	6.33	0.59	2.04	1.7	0.99	2.26	1.335	0.28	0.012	0.99	1.65	2.33
Interactions													
Lf x Loc	NS	NS	NS	NS	NS	NS	*	NS	NS	NS	*	NS	*
Lf x Dth	NS	NS	*	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Loc x Dth	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
LfxLocxDth	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Lf = Landform, Loc = Location, Dth = Depth, Trt = Treatments

Table 5. Interaction of Land Forms by Locations on ECEC, Ca²⁺ and Av.P

Locations	ECEC (cmol(+)kg ⁻¹)			Ca ²⁺ (cmol(+)kg ⁻¹)			Av.P (mgkg ⁻¹)		
	Sinamu	Tandanu	Masama	Sinamu	Tandanu	Masama	Sinamu	Tandanu	Masama
Wetlands	8.82 ^{ab}	12.97 ^a	8.04 ^b	3.08 ^b	3.07 ^b	3.07 ^b	14.41 ^{ab}	23.06 ^{ab}	27.28 ^a
Uplands	6.34 ^b	7.08 ^b	6.96 ^b	5.76 ^{ab}	8.78 ^a	4.24 ^{ab}	9.11 ^b	24.01 ^{ab}	22.99 ^{ab}
SE±	1.99	1.99	1.99	2.24	2.24	2.24	7.82	7.82	7.82

Table 6. Interaction of Landforms and Selected Soil Depths on pH CaCl₂

Depth (cm)	Sinamu	Tandanu	Masama
Wetlands	5.467 ^b	5.433 ^b	5.967 ^a
Uplands	5.267 ^b	6.10 ^a	5.467 ^b
SE±	0.197	0.197	0.197

3.7 Electrical Conductivity (EC)

The electrical conductivity (EC) ranged from 0.57 to 1.21 dSm⁻¹ in the wetland soils of the study area while 1.24 to 2.41 dSm⁻¹ was recorded in the drylands (Table 2). It was obvious that the dryland soils manifest higher EC values compared to the wetland soils. This may suggest more soluble salts probably contributed by calcium and magnesium salts. The EC was more at the surface than the sub-surface soils, probably due to upward capillary movement that pushes soluble salts and eventually got deposited or adsorbed to the soil particle surface. Umar, et al. [9] reported a range of 0.01 to 1.1 dSm⁻¹ in savanna soils which was in harmony with the range of values obtained in this study. Mean values of the electrical conductivity (EC) were statistically the same $P = .05$ between the two land forms, across the three research locations and also down the three selected soil depths. All the interactions were non-significant $P = .05$.

3.8 Soil Organic Carbon (OC gkg⁻¹)

The soil organic carbon in the wetland soils (Table 2) ranged from 1.5 to 7.7 gkg⁻¹ while values observed in the dryland soils ranged from 2.9 to 5.8 gkg⁻¹. Values obtained are very low, since values less than 20 gkg⁻¹ were generally considered low. Umar, et al. [9] reported a range of values from 1.0 to 7.8 gkg⁻¹ in savanna soils. Values reported 6.3 to 127 gkg⁻¹ by Aliyu, et al. [7] does not agree with the values reported in this work. The distribution pattern down the soil profile indicates consistent decrease from 0 up to 105 cm selected soil depths. Soil organic carbon content (OC gkg⁻¹) in the wetland soils and the dryland soils (Table 3) are statistically the same $P = .05$. The OC gkg⁻¹ content between the three selected soil depths were also statistically the same $P = .05$. There was a significant difference in O.C gkg⁻¹ between Tandanu and Masama

research locations but the two locations were statistically the same with Sinamu research location all at $P = .05$.

3.9 Available Phosphorus (Av. P)

The available P in the wetland soils (Table 2) ranged from 8.0 to 19.6 mgkg⁻¹ and 10.0 to 16.6 mgkg⁻¹ was recorded in the dryland soils. Higher values observed in the wetlands may suggest deposition of organic materials and inorganic phosphate ions mobilized and transported from upslope into the wetlands via run-off water. However, values recorded were classified within low to very low, since values less than < 20 ppm can still be considered low [23].

The distribution pattern down the soil profile revealed sharp decrease from surface to sub-surface soils across all the fields, suggesting that most of the P was concentrated at the surface soils in other words, available P decrease with increase in soil depth [24]. Habib, et al. [25] reported available P of 2.17 mgkg⁻¹ in Ungogo, Kano, Sudan Savanna, Nigeria. Mohammeda, et al. [10] reported a range of 18.6 to 24.0 mgkg⁻¹. Orji, et al. [26] reported a range of 3.30 to 40.00 mgkg⁻¹. All the values reported above were in agreement with the values obtained from wetland and dryland soils in Hadeja-Jama'are river basin in this study. The available P content of the two land forms (wetlands and dryland soils) (Table 3) were significantly different $P = .05$, suggesting mobilization and import of several soluble products such as herbicides, pesticides, fungicides, soluble inorganic P from phosphate fertilizers and organic materials brought in by run-off from the drylands into the wetland areas. Supporting the general opinion that wetland soils are often characterized by accumulation of organic matter [27], in addition, much higher temperature in the agro-ecological zone of the research area, especially in the dryland areas

may have contributed to this effect observed. The available P content of Sinamu and Masama research locations are significantly different ($P = .05$), but statistically the same with Tandanu research location ($P = .05$). The vertical spatial distribution of available P content down the soil profile indicates that, the selected surface soil (0 to 35 cm) was significantly different $P = .05$ from the sub-surface soils (35 to 70 cm and 70 to 105 cm). However, the two sub-surface soil depths (35 to 70 cm and 70 to 105 cm) were statistically the same $P = .05$ in available P content, giving more evidence the widely accepted opinion that P content in most soils is concentrated at the surface, usually 0 to 5 cm or 0 to 1 cm except in rare cases [28,29]. There was significant interaction between the land forms and research locations (Table 3) probably due to variations in Fe and Al oxides associated to clay content, or variation in calcium and magnesium content or variation in organic carbon content or combine. The highest ranked interaction (Table 4) was observed in Masama wetland and it was not significantly different from Sinamu, Tandanu wetlands and Tanadanu, Masama drylands, however there is significant difference in the interaction between Masama wetland and Sinamu drylands $P = .05$.

4. CONCLUSION

Base on the findings above, the soils parameters regulating P availability and transformation in both wetlands and dryland soils in the order of increasing magnitude include silicate clays, pH, O.C, ECEC, EA, Ca and Mg.

Base on the Statistical findings obtained, phosphate availability is higher in the wetlands of Hadejia-Jama'are than it is in the drylands suggesting deposition of organic materials and inorganic phosphate ions mobilized, transported from up-slope by run-off and deposited in to the wetland soils.

Phosphate ions were more available in Sinamu wetlands, moderately available in Tandanu wetlands and less available in Masama wetlands which cannot be unconnected to the low nature of organic carbon observed in Masama wetland soils compared to Sinamu and Tandanu wetland soils.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image

generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Hamoud YA, Shaghaleh H, Zhang K, Okla KM, Alaraidh IA, and SM, Elgawad HA. Increasing Soil Clay Content Increases Soil Phosphorus Availability and Improves the Growth, Physiology, and Phosphorus Uptake of Rice Under Alternative Wetting and Mild Drying Irrigation. *Environmental Technology & Innovation*. 2024;35.
2. Ducouso-Detrez A, Fontain J, Lounes-Hadj Sahraoui A, Hijri M. Diversity of Phosphate Chemical Forms in Soils and their Contributions on Soil microbial community structure changes. *Microorganisms*. 2022;10(3):609.
3. Goyal SK, Singh AK. Phosphate sorption and desorption in alkaline soils. *Agricultural Research Journal*. 2020;57(2):598-604.
4. Leinweber P, Morshedizad M, Debicka M. The effects of Dissolved Organic Matter Derived from Agricultural Waste Materials on Phosphate Sorption in Sandy Soils. *Agriculture*. 2023;13 (11):2164.
5. Ahmed AA, Leinweber P, Kuhn O. Advances in Understanding the Phosphate Binding to Soil Constituents: A Computation Chemistry Perspective. *Science of the Total Environment*. 2023; 887:163692
6. Khundzhua DA, Yuzhakov VI, Korvatovskiy BN, Paschenko VZ, Kulyabko LS, Kydraliev KA, Patsaeva SV. Spectroscopic manifestation of interaction of humic acids with trivalent iron ions in aqueous solutions. *Mosc. Univ. Phys. Bull*. 2020;73:632–637.
7. Aliyu M, Nuhu OI, Saminu AI, Azare IM, Abdurkadir M, Abdu N. Pedogeochemical assessment of wetland soils in Hadejia-Jama'are river basin in the Nigerian Sahel savanna. *International Journal of Environmental Quality*. 2022;51.
8. Goni I, Vassolo S, Sherif M, Aji M, Bura BB, Ibrahim Y. Geochemical and isotopic studies in parts of the Hadejia-Jamaare-Komadugu-Yobe Basin, northeast Nigeria. *Hydrogeology Journal*. 2023;32:113–129.

9. Umar G.1*, Magaji MJ, Buji IB. Comparative Analysis of Some Soil Properties in Different Land Use of Auyo Local Government Area, Jigawa, North West, Nigeria. *Journal of Arid Zone Economy*. 2023;2(1):47–57.
10. Mohammeda SE, Aderibigbeb DO, Saeed MD, Nelanad SM, Musaa A, Ayandae OS. Seasonal quality and statistical assessment of physicochemical properties and base saturation of Hadejia-Nguru wetland soils, Northwest/Eastern Nigeria. *The geographical journal: Water Practice & Technology*. 2024;19;3:761. DOI: 10.2166/wpt.2024.035
11. Lal R. (Ed). *Encyclopedia of Soil Science (4th Ed)* CRC Press; 2021.
12. Uyovbisere EO, Ogunwole JO, Odigie VO, Abdu N. *Laboratory Manual of Routine Soil, Water, Plant and Fertilizer Analyses*. A compilation of the Department of Soil Science, Faculty of Agriculture, Institute of Agricultural Research, Ahmadu Bello University Zaria; 2013.
13. Soil Survey Staff. *Soil Survey Manual (Agriculture Handbook No. 18)*. United State Department of Agriculture, Natural Resources Conservation Service; 2017.
14. Brevick EC, Burgess LC. (Eds). *Soil Health Across Landscape: Opportunities and Challenges (2nd ed.)*, Academic Press; 2021.
15. Rhoades JD, Manteghi NA. Measurement O Soil Salinity: Electrical Conductivity and Total Dissolve Solids. In J.D. Rhoades and D.L. Corwin (Eds.), *Soil Salinity Assessment: Methods and Interpretation of Electrical Conductivity Measurements*. 2018;5-11.
16. Sparks DL. (Ed.) *Advances in Agronomy*. 2021;173.
17. *Statistical Analysis System SAS*. Statistical package Software version 9.3.SAS Institute, Cary, North Carolina; 2011.
18. Shehu BM, Jibrin JM, Samndi AM. Fertility Status of Selected Soils in the Sudan Savanna Biome of Northern Nigeria *International Journal of Soil Science*. 2015;10(2):74-84.
19. Food and Agriculture Organization F.A.O *Sodic Soils and their Management*; 2019. Available:www.fao.org/3/x5871e/5871e05.. Page visited on 7/6/2019
20. Yakubu A. Interaction of Microbial, Organic and Inorganic Sources of Phosphorus on the Performance of Maize (*Zea mays* L.) in Two Soils of Different Fertility. (Project for Bachelor's Degree in Agriculture, Ahmadu Bello University); 2016.
21. Daniel CJ, Frid L, Sleeter BM, Fortim MJ. State-and-transition Simulation Model: A Framework for Forecasting Landscape Change. *Methods in Ecology and Evaluation*. 2016;(7):11.
22. Abdu N, Udofot EA. Fifteen-year Fallow Altered the Dynamics of Soil Phosphorus and Cationic Balance in a Savanna Alfisol. *Archives of Agronomy and Soil Science*. 2015;61:645-656.
23. Chimdi A, Negasa D, Chala G. Effects of Rhizobium Inoculation and P Fertilizer Leves o Selected Soil Properties, Yield, and Yield Components of Foba Bean (*Vicia foba* L.): The Case of Abuna Gindeberat, West Shewa Zone, Ethiopia; 2022. Available:https://onlinelibrary.wiley.com/doi/full/10.1155/2022/3635989
24. Johnbosco C, Egbueri JC, Ighalo JO, Pande CAB. Editors. *Climate Change Impacts on Nigeria Environment and Sustainable Development*. 2023; 339.
25. Habib DW, Umar AM, Usman YA, Abdurrahman BL. Effect of Municipal Solid Waste Treatment Techniques on Soil Chemical Properties in Sudan Savannah, Kano State, Northern Nigeria. *Journal of Wastes and Biomass Management*. 2023; 5(2):82-87. DOI:http://doi.org/10.26480/jwbm.02.2023.8 2.87
26. Orji JE, Igwe CA, Ezeaku PI. Effect of Climate Change on Soil Organic Carbon Storage in Four Land Use Types in Abakaliki, South Eastern Nigeria. Johnbosco C, Egbueri JC, Ighalo JO, Pande CAB. (Ed). *Climate Change Impacts on Nigeria Environment and Sustainable Development*. 2023; 135.
27. Fu C, Li Y, Zeng L, Tu C, Wang X, Ma H, Xiao L, Christie P, Luo Y. Climate and Mineral Accretion as Drivers of Mineral-Associated and Particulate Organic Matter Accumulation in Tidal Wetland Soils. *Global Change Biology*. 2023;30(1).

28. Leo E, Nathan S, Morteza M, (n.d) Understanding Numbers on your Soil Test Report. University of Arkansas, USA. Available: <https://www.uaex.edu>>PDF Site visited 29/6/2019
29. Soil Science Society of America (SSSA). Soil Science Society of America. Methods Book (3rd Ed). Soil Science Society of America; 2016.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://prh.ikpress.org/review-history/12249>