



Impact of Organic Foliar Formulations on Growth and Yield of Okra (*Abelmoschus esculentus*)

Ghodke P. B. ^a, Jakku Prasanna ^{b*}, Ubale S. P. ^c,
Warpe S. T. ^d and Mirza I. A. B. ^e

^a Department of Agronomy, Punjab Agricultural University, Ludhiana, India.

^b Central Research Institute for Dryland Agriculture, Hyderabad, India.

^c Department of Genetics and Plant Breeding, College of Agriculture, Pune, MPKV, Rahuri, India.

^d Department of Agronomy, College of Agriculture, Pune, MPKV, Rahuri, India.

^e Department of Agronomy, VNMKV, Parbhani, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i121811>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/127697>

Original Research Article

Received: 11/10/2024

Accepted: 17/12/2024

Published: 26/12/2024

ABSTRACT

Aims: To evaluate the effect of foliar application of different organic formulations on growth and yield of okra.

Study Design: Completely Randomized Block Design

Place and Duration of Study: Agronomy organic farm, College of Agriculture, Pune, during summer season of 2019-20.

Methodology: The experiment was laid out in Completely Randomized Block Design with eight treatments and three replications. The gross and net plot sizes were 3.30 m x 3.00 m and 2.70 m x

*Corresponding author: E-mail: Prasanna.j10@gmail.com;

Cite as: P. B., Ghodke, Jakku Prasanna, Ubale S. P., Warpe S. T., and Mirza I. A. B. 2024. "Impact of Organic Foliar Formulations on Growth and Yield of Okra (*Abelmoschus Esculentus*)". *Journal of Advances in Biology & Biotechnology* 27 (12):637-42. <https://doi.org/10.9734/jabb/2024/v27i121811>.

2.40 m, respectively. The treatments consisted seven organic formulations with one control i.e. no foliar spray viz. control (no foliar spray), T₁: control (no foliar spray), T₂: foliar application of cow urine at 5%, T₃: foliar application of jeevamrut at 5%, T₄: foliar application of EM solution at 1%, T₅: foliar application of humic acid at 2%, T₆: foliar application of fulvic acid at 2%, T₇: foliar application of vermiwash at 5% and T₈: foliar application of seaweed extract at 0.25%.

Results: The results revealed that all the growth contributing characters viz., plant height, number of branches plant-1, plant spread, number of nodes plant-1, internodal length and leaf area plant-1 were found significantly higher with the foliar application of seaweed extract at 0.25%. This was followed by foliar application of vermiwash at 5%. However, substantial reduction in growth characters was observed with control (no foliar spray).

Conclusion: The expression of growth characters in okra like Plant height, number of branches per plant, Plant spread, Number of nodes, internodal length and Leaf area were recorded higher with the foliar application of sea weed extract at 0.25% and were found statistically difference among all growth characters except the Internodal length which was found to be non-significant.

Keywords: Foliar spray; sea weed extract; plant height; okra; malvaceae family.

1. INTRODUCTION

Okra (*Abelmoschus esculentus* L.) is an annual, erect, broadleaved, hairy plant belonging to the Malvaceae family. It is originated in Ethiopia (Sathish and Eshwar, 2013) and was then propagated in North Africa, in the Mediterranean, in Arabia and India by the 12th century B.C. (Nzikou et al., 2006). Among the various vegetable crops, okra is one of the fruit vegetables. Okra is widely cultivated vegetable and can be found in almost every market all over India (Acharya et al., 2021).

The exploitative agriculture for a long time in our country has brought down the fertility status of the soil to a level that even provision of high rate of fertilizers is unable to sustain the productivity of soil. So as to attain the productivity of soil and promote the health of the soil, use of organic sources is imperative (Bhattacharyya and Mathur, 2017).

Recent developments in intensive agriculture, though contributed immensely towards surplus food, caused degradation of fertile land and left residues in food products. Thus, there is need of increasing awareness throughout the world about the organic, sustainable agricultural practices.

2. MATERIALS AND METHODS

Field experiment was conducted to study the effect of foliar sprays of organic formulations on growth and yield of okra at the Agronomy organic farm, College of Agriculture, Pune, during summer season of 2019-20. The experiment was laid out in Completely

Randomized Block Design with 08 treatments which include T₁: control (no foliar spray), T₂: foliar application of cow urine at 5%, T₃: foliar application of jeevamrut at 5%, T₄: foliar application of EM solution at 1%, T₅: foliar application of humic acid at 2%, T₆: foliar application of fulvic acid at 2%, T₇: foliar application of vermiwash at 5% and T₈: foliar application of seaweed extract at 0.25% imposed to okra variety Phule Vimukta. The foliar sprays were taken at 30 and 45 days after sowing. The gross and net plot sizes were 3.30 m x 3.00 m and 2.70 m x 2.40 m, respectively. The okra seed was dibbled at a spacing of 30 x 15 cm².

A full dose of Nitrogen, Phosphorus and Potassium was applied through vermicompost as a basal application at the time of sowing. *Azotobacter*, PSB and *Trichoderma* seed treatment was given to all the treatments.

Plant protection measures were undertaken as per the necessity to protect the crop from pests and disease incidence by using organic measures. The spraying of NSKE at 10 ml /lit, Metarrhizium at 5 g/lit and Dashparni Ark at 25 ml/lit was used for control of sucking pests.

3. RESULTS AND DISCUSSION

Plant height, Number of branches, Plant spread, Number of nodes, Internodal length and Leaf area: Effect of foliar application of organic formulations were significantly influenced plant height, number of branches, plant spread and number of nodes per plant at 42, 56, 70 and 84 days after sowing whereas leaf area was influenced significantly at all the

growth stages and number of nodes per plant was found to be influenced non-significantly at all the growth stages.

Plant height: The significantly taller plants were recorded with the foliar spray of sea weed extract at 0.25% (T₈). However, it was on par with foliar application of vermiwash at 5% (T₇) and with foliar application of cow urine at 5% (T₂), foliar application of jeevamrut at 5% (T₃), foliar application of humic acid at 2% (T₅) and foliar application of vermiwash at 5% (T₇).

The taller plants observed with the application of seaweed extract might be due to the plant growth regulators that are present in the seaweed and seaweed extracts, including cytokinins, auxins and GA's (Sadeghipour et al., 2022). It is hypothesized that seaweed synthesizes relatively high concentration of plant growth regulator's because this confers an adaptive advantage that allows it to survive the extreme temperature range (Crouch et al. 1990). Gollan and Wright (2006) concluded that the increase in shoot characteristics due to the auxins content in the seaweed extracts which have an effective role in cell division and enlargement which leads to increase the shoot growth (Kim et al., 2023).

Number of branches: The maximum number of branches plant⁻¹ was obtained with the foliar application of sea weed extract at 0.25% which was significantly superior over all other treatments. It was on par with foliar application of jeevamrut at 5% (T₃) and foliar application of vermiwash at 5% (T₇) and it was on par with foliar application of cow urine at 5% (T₂). The higher number of branches recorded with the application of sea weed extract was 1.67.

This increase in number of branches plant⁻¹ was owing to more availability of essential nutrients in a balanced form to plants which supported the plant growth and development. Similar results were recorded by Rathore et al. (2009) and Sharma et al. (2022).

Plant spread: The differences in plant spread due to different organic formulations were influenced significantly. The foliar application of seaweed extract at 0.25% recorded significantly higher plant spread as compared to rest of the treatments. However, it was on par with the foliar application of vermiwash at 5% (T₇).

The increase in plant spread may be due to availability of optimum amount of nutrients from the sea weed extract.

Number of nodes: The foliar application of seaweed extract at 0.25% produced significantly more number of nodes plant⁻¹ than all other treatments. The next best treatment was found to be vermiwash at 5%, whereas the lower number of nodes were recorded from control treatment in which no foliar spray was undertaken. The results were found to be on par with the foliar application of cow urine at 5%, foliar application of jeevamrut at 5%, foliar application of humic acid at 2% and foliar application of vermiwash at 5%.

Internodal length: The data revealed that effect of different foliar application treatments on internodal length in okra was found to be non-significant. However, numerically internodal length was recorded higher with foliar application of seaweed extract at 0.25% (T₈). Whereas, no foliar spray (T₁) recorded significantly the lowest internodal length in okra.

Leaf area: The difference in leaf area plant⁻¹ due to different organic formulations was statistically significant at all the crop growth stages. The foliar application of sea weed extract at 0.25% resulted in significantly higher leaf area plant⁻¹ as compared to rest of the treatments at all the crop growth stages. However, it was on par with the foliar application of cow urine at 5%, Jeevamrut at 5%, Humic acid at 2% and Vermiwash at 5%. This increase in growth and biomass accumulation in the said nutrient combination could be due to various metabolic processes which could have led to the enhanced leaf area plant⁻¹. The minimum leaf area plant⁻¹ was reported under control i.e. no foliar spray (T₁).

Kavitha et al. (2008) reported that seaweed extract foliar spray significantly influenced the growth attributes at all the stages of observation and also observed that the maximum LAI (6.38) of rice was recorded with spraying of seaweed extract at 0.3% twice at 50% flowering and milk stages than other treatments and was followed by spraying of sea weed extract at 0.3% at milk stage (Selvakumar et al., 2022).

Table 1. Treatment variability in plant height, number of branches, plant spread, number of nodes, internodal length and leaf area

Treatment	Plant height	Number of branches	Plant spread	Number of nodes	Internodal length	Leaf area
T ₁ : Control (No folia spray)	90.26	1.20	38.95	22.53	4.01	17.00
T ₂ : Foliar application of cow urine at 5%	107.71	1.47	41.54	26.27	4.10	24.60
T ₃ : Foliar application of Jeevamrut at 5%	109.14	1.53	41.98	26.49	4.12	27.23
T ₄ : Foliar application of EM-Solution at 1%	93.07	1.33	39.18	22.78	4.03	17.17
T ₅ : Foliar application of Humic acid at 2%	102.33	1.40	40.35	25.08	4.08	23.47
T ₆ : Foliar application of Fulvic acid at 2%	97.74	1.40	40.16	24.07	4.06	19.40
T ₇ : Foliar application of Vermiwash at 5%	115.82	1.53	43.99	28.78	4.13	27.78
T ₈ : Foliar application of Sea weed extract at 0.25%	119.79	1.67	46.48	28.86	4.15	27.83
S.E.m±	5.79	0.09	1.42	1.40	0.27	1.94
C.D. at 5 %	17.57	0.26	4.32	4.24	NS	5.95
General mean	104.48	1.44	41.58	25.61	4.08	23.06

4. CONCLUSION

The expression of growth characters in okra like plant height, number of branches per plant, plant spread, number of nodes, internodal length and leaf area were recorded higher with the foliar application of sea weed extract at 0.25% and were found statistically significant among all growth characters except the internodal length which was found to be non-significant.

Recent developments in intensive agriculture, though contributed immensely towards surplus food, caused degradation of fertile land and left residues in food products. Thus, there is need of increasing awareness throughout the world about the organic, sustainable agricultural practices.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENTS

A Sincerely thankful to Proff. A. A. Sheikh (Retd.) Division of Agronomy, College of Agriculture, Mahatma Phule Rahuri Kishi Vidhyapeeth for providing the required facilities for the current research experiment.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Acharya, R., Skalicky, M., Maitra, S., Umair, M., Bejandi, T. K., Brestic, M., & Hossain, A. (2021). Insight into the morphological, physiological, biochemical, and molecular responses of okra (*Abelmoschus esculentus* L.) under drought stress. *Plants*, 10(10), 2087. <https://doi.org/10.3390/plants10102087>
- Bhattacharyya, P., & Mathur, R. K. (2017). Organic farming in India: Opportunities and challenges. In *Organic farming in India: Opportunities and challenges*

- (pp. 1-15). Springer Singapore. <https://doi.org/10.1007/978-981-10-5217-3>
- Crouch, L. J., Beckett, R. P., & Van Staden, J. (1990). Effect of seaweed concentrate on the growth and mineral nutrition of nutrient-stressed lettuce. *Journal of Applied Phycology*, 2(3), 269-272.
- Gollan, J. R., & Wright, J. T. (2006). Limited grazing pressure by native herbivores on the invasive seaweed *Caulerpa taxifolia* in a temperate Australia estuary. *Marine and Freshwater Research*, 57(7), 685-694.
- Kavitha, M. P., Ganesaraja, V., & Paulpandi, V. K. (2008). Effect of foliar spraying of seaweed extract on growth and yield of rice (*Oryza sativa* L.). *Agricultural Science Digest*, 28(2), 127-129.
- Kim, D. H., Lee, S. Y., Jeong, M. J., Kwon, O. W., Kim, S. H., & Oh, S. E. (2023). Effects of seaweed extract on growth, physiological characteristics, and antioxidant activity of lettuce (*Lactuca sativa* L.) under different light conditions. *Horticulturae*, 9(4), 456.
- Nzikou, J., Mvoula-Tsieri, M., & Matouba, E. (2006). A study on *Gumbo* seed grown in Congo Brazzaville for its food and industrial applications. *African Journal of Biotechnology*, 5(24), 2469-2475.
- Rathore, S. S., Chaudhary, D. R., Boricha, G. N., Ghosh, A., Bhatt, B. P., Zodape, S. T., & Patolia, J. S. (2009). Effect of seaweed extract on the growth, yield, and nutrient uptake of soybean (*Glycine max*) under rainfed conditions. *South African Journal of Botany*, 75(2), 351-355.
- Sadeghipour, O., Aghili, H., Dehghani, F., & Khalili, M. (2022). Foliar application of seaweed extract improves growth, yield, and biochemical characteristics of green bean (*Phaseolus vulgaris* L.) under different irrigation regimes. *Plants*, 11(16), 2168.
- Satish, D., & Eshwar, A. (2013). A review on: *Abelmoschus esculentus* (okra). *International Research Journal of Pharmaceutical and Applied Sciences*, 3(4), 129-132.
- Selvakumar, P., Ramya, B., Thiruppathi, M., & Ananthi, S. (2022). Impact of seaweed liquid fertilizer on growth and yield of rice (*Oryza sativa* L.) under different nutrient management practices. *Journal of Soil Science and Plant Nutrition*, 22(3), 3456-3467.

Sharma, A., Kumar, V., Chaudhary, A., & Singh, B. (2022). Nutrient management and its impact on branching pattern and

yield attributes of vegetable crops. *Journal of Plant Nutrition*, 45(8), 1145-1156.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/127697>