



Annual Research & Review in Biology

Volume 40, Issue 1, Page 44-49, 2025; Article no.ARRB.129048

ISSN: 2347-565X, NLM ID: 101632869

(Past name: Annual Review & Research in Biology, Past ISSN: 2231-4776)

Impact of Organic Inputs (Jeevamruth and Beejamruth) on Quality Parameters of Fenugreek

Sugumaran M.P^{a*}, Goveanthan A.S^b, Kiruba M^{c*},
Bharathi Kumar K^{a*}, Akila S^d, Somasundaram E^b,
Natarajan K^a, Gayathry G^a and Sinduja M^d

^a Krishi Vigyan Kendra, TNAU, Vridhachalam, India.

^b Tamil Nadu Agricultural University, Coimbatore, India.

^c Krishi Vigyan Kendra, TNAU, Sandhiyur, India.

^d National Agro Foundation, Chennai, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/arrb/2025/v40i12188>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129048>

Original Research Article

Received: 05/11/2024

Accepted: 07/01/2025

Published: 11/01/2025

ABSTRACT

The present study on Biochemical evaluation of organic inputs (Jeevamruth and Beejamruth) and their efficacy on Greens was carried out at the Department of sustainable organic agriculture, Tamil Nadu Agricultural University, Coimbatore and the laboratory experiments were carried out at the Department of Environmental Sciences. A field experiment was conducted in a randomized block design with three replications. Biometric observations were taken during 10th, 20th and 30th days after sowing. Soil samples were collected during initial and at the stage of harvest and analysed for

*Corresponding author: E-mail: sugumaran.mp@tnau.ac.in; kiruba.m@tnau.ac.in, bharathisolunam@gmail.com;

Cite as: M.P, Sugumaran, Goveanthan A.S, Kiruba M, Bharathi Kumar K, Akila S, Somasundaram E, Natarajan K, Gayathry G, and Sinduja M. 2025. "Impact of Organic Inputs (Jeevamruth and Beejamruth) on Quality Parameters of Fenugreek". Annual Research & Review in Biology 40 (1):44-49. <https://doi.org/10.9734/arrb/2025/v40i12188>.

chemical and biological properties. Plant samples were analysed for protein and chlorophyll contents. Quality parameters like protein content was found to be high in both treatments Jeevamruth @ 5 % Spray (using Green gram flour) and Jeevamruth @ 5 % Spray (using Green gram +Black gram flour); and the chlorophyll content was high in Jeevamruth @ 5 % Spray (Green gram flour). From the results, the application of Jeevamruth as a 5 % spray was observed as a viable organic approach to improve soil and eco-friendly fenugreek production.

Keywords: Jeevamruth; beejamruth; biochemical evaluation; organic inputs; protein content; chlorophyll content.

1. INTRODUCTION

“With the growing awareness for safe and healthy food, the demand for organic food is increasing. Concern for deteriorating soil health and increasing contamination in ground water and surface water bodies have also added to the need for alternative technologies which not only can ensure safe and healthy food but are also environment friendly, contribute to the long-term fertility and sustainability of the soils. During the last 10 years, many farmers due to various reasons, have switched over to the organic methods of cultivation and are now successfully growing comparable productivities with much less costs. In the absence of any scientific support, majority of the farmers relied on the traditional wisdom” (Reddy et al., 2022). “On-farm experimentation has developed innovative inputs which has to be prepared on-farm with local resources. Among these innovative inputs most common and successful inputs are Panchagavya, Jeevamruth and Vermiwash. Among the liquid formulations, panchagavya is one of the most important liquid formulations to be considered for the study of shelf life as it not only proves to be most effective for crop growth but also widely used by the farmers” (Sugumaran et al., 2018).

“Nowadays organic farming practices are gaining importance as farmers have realized the benefits of organic farming in terms of soil fertility, soil health and sustainable productivity. Farmers are well aware with the use of organic liquid manure such as Panchagavya, Beejamruth and Jeevamruth as well as bio digester in organic farming. These organic liquid manures play a key role in promoting growth and providing immunity to plant system. The spray of Panchgavya on crop imparts dark green colour leaves within 10 days. Its role as plant growth promoter has already” been reported by Kumar, (2022) and Sreenivasa et al., (2010).

2. MATERIALS AND METHODS

2.1 Field Experimental Details

The experiment was conducted in a randomised block design with three replications. The experimental layout was kept undisturbed throughout the period of investigation and fenugreek seeds were soaked for 1 hr in the Beejamruth solution before sowing and were sown in the field.

Design : RBD

Number of treatments : 7

Number of replications : 3

The treatment details are given below:

Treatments:

T₁–Control

T₂-Jeevamruth @ 3% Spray (Green gram flour)

T₃-Jeevamruth @ 5 % Spray (Green gram flour)

T₄-Jeevamruth @ 3 % Spray (Black gram flour)

T₅-Jeevamruth @ 5 % Spray (Black gram flour)

T₆-Jeevamruth @ 3 % Spray (Green gram + Black gram flour)

T₇-Jeevamruth @ 5 % Spray (Green gram +Black gram flour)

2.2 Field Preparation and Sowing

The field was thoroughly prepared to a fine tilth. The soil type of experimental plot was clay loam with a pH of 7.42. Seeds of fenugreek variety Co1 selection were sown in lines adopting a spacing of 30 × 15 cm at the rate of 40 grams per bed (size 2.0×2.0). Plants were thinned at 3 days after sowing.

2.3 Leaf Protein

The leaf protein was estimated as per the method described by Lowrey et al., (1951).

2.4 Leaf Chlorophyll Content

The amount of chlorophyll was estimated according to Yoshida et al. (1971).

3. RESULTS AND DISCUSSION

3.1 Protein Content

The treatments Jeevamruth @ 5 % Spray (using Green gram flour) and Jeevamruth @ 5 % Spray (using Green gram flour+Black gram flour) are on par with each other and the protein contents were 1.493 mg/100 g and 1.492 mg/100g of greens respectively. Lowest protein content was recorded in control (0.795 mg/100 g) (Fig. 2). The nutritional quality of fenugreek plant is decided by parameters like protein content and chlorophyll content. Prabhakaran & Pitchai (2002) reported that the application of recommended dose of nitrogen through poultry manure recorded higher protein in tomato fruit compared with inorganic fertilizer. The present investigation results go in support of Beulah (2001). Montagu & Goh (1990) also found that the rate of nitrogen through fertilizer significantly decreased vitamin C concentration and fruit visual quality in tomato. Boomiraj (2003) had also reported the increased protein content of Bendi fruit due to panchagavya application and Jeevamruth application. The microorganisms

present in the Jeevamruth would have helped in the fixation of atmospheric nitrogen and its adsorption and utilization for protein synthesis. Increased nitrogen content would increase protein content. These results are in line with earlier findings of Roy & Seth (1971) in radish, Venter (1979) in carrot, Bome et al. (1987) in cabbage, Kohil et al. (1992) in pea, Beulah (2001) in moringa and Somasundaram (2003) in maize, sunflower and green gram. "The plant height, root length and single plant weight in fenugreek are high in the treatment as Jeevamruth 5% spray was observed as a viable organic approach to improve soil and eco-friendly fenugreek production" (Goveanthan et al., 2020). "In another study, seeds treated with Panchagavya and Jeevamruth separately, the maximum shoot length and root length were recorded in Panchagavya treatment and minimum shoot length and root length of was recorded in Jeevamruth treated seeds and also the panchagavya treated seeds registered the maximum vigour index" (Akila et al., 2020). "Compounds Erioflorin and nagilactone A present in Beejamruth are plant growth regulators and hence Beejamruth is a valid and effective alternative fertilizer for the production of safe and good quality food produces which could cater the needs of the modern Indian requirements" (Goveanthan et al., 2019).



Fig. 1. Experimental field view




Fig. 2. Effect of organic nutrient source (jeevamrutham) on (µg/100 g) of Fenugreek plant leaf protein content

Table 1. Effect of organic nutrient source (Jeevamruth) on Fenugreek plant leaf Chlorophyll content

Treatment	Chlorophyll A	Chlorophyll B	Total
T ₁ - Control	0.5441	0.3619	1.3423
T ₂ - Jeevamruth @ 3% Spray (Green gram)	1.5854	0.3228	1.474
T ₃ - Jeevamruth @ 5 % Spray (Green gram)	2.3765	0.1045	2.854
T ₄ - Jeevamruth @ 3 % Spray (Black gram)	1.4663	0.6776	1.795
T ₅ - Jeevamruth @ 5 % Spray (Black gram)	1.7760	0.3338	2.352
T ₆ - Jeevamruth @ 3 % Spray (Green gram + Black gram)	1.3977	0.3179	1.911
T ₇ - Jeevamruth @ 5 % Spray (Green gram + Black gram)	1.5778	0.4441	2.453
SEd	0.0913	0.021	0.1182
CD (P = 0.05)	0.199	0.0458	0.2575

In another study, the soluble protein content (0.87 mg/g) and total sugar content (11.20 µg/g) were found to be improved in Panchagavya (Groundnut cake instead of ghee) @ 3% spray treatment and Recommended dose of NPK fertilizer applied plants (Sugumaran et al., 2019).

3.2 Chlorophyll Content

The chlorophyll A content was high (2.377) in the treatment Jeevamruth @ 5% Spray (Black gram flour) (1.776) followed by Jeevamruth @ 5% Spray (using Green gram flour+Black gram flour) (1.578). The lowest chlorophyll A content was

found in control (0.544). The chlorophyll B content was high (0.678) in the Jeevamruth @ 5% Spray (using Green gram flour). The lowest chlorophyll B content was found in control (0.362). The total chlorophyll content was high (2.854) in the treatment Jeevamruth @ 5% Spray (using Green gram flour). The total chlorophyll content was found to be on par with Jeevamruth @ 5% Spray (using Green gram flour+Black gram flour) (2.453) and Jeevamruth @ 5% Spray (Black gram flour) (2.352). The lowest total chlorophyll content was found in control (1.342) (Table 1). Magnesium is essential for normal plant growth and development (Bose

et al., 2011). Mg is an integral part of chlorophyll, essential for photosynthesis. Thus the presence of magnesium in the Jeevamruth will supplement the magnesium requirement of the crop. The maize plants in treatment TNAU Panchagavya @ 3% spray and Panchagavya (Groundnut cake instead of ghee) @ 3% spray increased the quantity of photosynthetic pigments and root oxidation activity than in the control plants (Sugumaran et al., 2019). "Compounds Isoanthranic acid, Columbianetin and Lomatolin, 1, 6-Hexanediol, Mevastatin and Gitoxigenin, Dibutoxy anthracene, Erioflorin and nagilactone, Trimegestonea, Rofe Coxib, Clupanodonic acid are present in Jeevamruth which helps in the plant metabolism and also improves its growth" (Goveanthan et al., 2021).

4. CONCLUSION

From the field experiment conducted at Research Farm, Department of Sustainable Organic Agriculture, Tamil Nadu Agricultural University, Coimbatore to test the effect of Jeevamruth and Beejamruth on Fenugreek, it is inferred that, the quality parameters like protein content was found to be high in both Jeevamruth @ 5 % Spray (using Green gram flour) and Jeevamruth @ 5 % Spray (using Green gram flour+Black gram flour); and the chlorophyll content was high in the treatment Jeevamruth @ 5 % Spray (using Green gram flour).

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

Akila, S., Sugumaran, M. P., Suganya, K., & Somasundaram, E. (2020). Studies on testing the efficacy of liquid organic inputs (Panchagavya and Jeevamruth) on maize (*Zea mays* L.) germination. *Current Journal of Applied Science and Technology*, 39(23), 134-137.

Beaulah, A. (2001). *Growth and development of moringa (Moringa oleifera Lem.) under organic and inorganic system of culture* (Ph.D. thesis). Tamil Nadu Agricultural University, Coimbatore.

Bome, N. A., & Petunina, T. L. (1987). Evaluation of the combining ability of lucerne breeding material by the polycross method.

Boomiraj, K. (2003). *Evaluation of organic sources of nutrients, Panchagavya and botanicals spray on Bhendi (Abelmoschus esculentus Moench)* (M.Sc. thesis). Tamil Nadu Agricultural University, Coimbatore.

Bose, J., Babourina, O., & Rengel, J. (2011). Role of magnesium in alleviation of aluminium toxicity in plants. *Journal of Experimental Botany*, 62, 2251-2264.

Goveanthan, A. S., Sugumaran, M. P., & Somasundaram, E. (2019). Biochemical analysis of Beejamruth and its plant promoting factors. *International Journal of Current Research and Academic Review*, 7(5), 1-4.

Goveanthan, A. S., Sugumaran, M. P., & Somasundaram, E. (2021). Scientific validation of organic liquid formulation—Jeevamruth by studying its characteristics. *International Journal of Plant Sciences*, 16(1), 15-18.

Goveanthan, A. S., Sugumaran, M. P., Gudimetha, G. K., Akila, S., Suganya, K., & Somasundaram, E. (2020). Studies on organic inputs (Jeevamruth and Beejamruth) and their efficacy on fenugreek. *The Pharma Innovation Journal*, 9(11), 92-94.

Kohil, V. K., Thakur, I. K., & Shukla, Y. R. (1992). A note on response of pea (*Pisum sativum* L.) to P and K application. *Horticultural Journal*, 5(1), 59-61.

Kumar, K. (2022). Significance of Liquid Organic Manures in Indian Agriculture—A Review. *Agriculture*, 5(1), 17-24.

Lowry, O. H., Rosebrough, N. J., Farr, A. L., & Randall, R. J. (1951). Protein measurement with the Folin phenol reagent. *J Biol Chem*, 193(1), 265-275.

Montagu, K. D., & Goh, K. M. (1990). Effects of forms and rates of inorganic nitrogen fertilizers on the yield and some quality indices of tomato. *Journal of Crop and Horticultural Science*, 18, 31-37.

Prabhakaran, C., & Pitchai, G. J. (2002). Effect of different organic nitrogen sources on pH, total soluble solids, titratable acidity, reducing and non-reducing sugars, crude protein, and ascorbic acid content of

- tomato fruits. *Journal of Soils and Crops*, 12(2), 160-166.
- Reddy, G. S., Rai, P. K., & Nagar, S. (2022). Pre-sowing seed treatment with Panchagavya, Beejamruth, and leaf extract of Moringa and neem on growth, yield, and yield-attributing traits of cowpea (*Vigna unguiculata* L.) cv-Bali265. *International Journal of Plant & Soil Science*, 34(22), 906-916.
<https://doi.org/10.9734/ijpss/2022/v34i2231451>.
- Roy, R. N., & Seth, J. (1971). Nutrient Uptake and Quality of Radish (*Raphanus Satjvus* L.) As Influenced by Levels of Nitrogen, Phosphorus and Potassium and Methods of Their Application. *Indian Journal of Horticulture*, 28(2), 144-149.
- Somasundaram, E. (2003). *Evaluation of organic sources of nutrients and Panchagavya spray on the growth and productivity of maize-sunflower-green gram system* (Ph.D. thesis). Tamil Nadu Agricultural University, Coimbatore.
- Sreenivasa, M.N., Nagaraj M. Naik and Bhat, S.N., 2010, Beejamruth: A source for beneficial bacteria. *Karnataka J. Agric. Sci.*, 17(3): pp.72-77.
- Sugumaran, M. P., Akila, S., & Somasundaram, E. (2018). Studies on analyzing the shelf life of Panchagavya with different alternatives for ghee. *International Journal of Agriculture Sciences*, 10(24), 7655-7656.
- Sugumaran, M. P., Akila, S., & Somasundaram, E. (2019). Studies on analysis of biochemical characters of leaf over liquid organic inputs (Panchagavya and Jeevamruth) on maize (*Zea mays* L.). *Journal of Pharmacognosy and Phytochemistry*, 8(5), 1794-1797.
- Venter, F. (1979, June). Nitrate contents in carrots (*Daucus carota* L.) as influenced by fertilization. In *Symposium on Quality of Vegetables* 93 (pp. 163-172).
- Yoshida, S., Forno, D. A., & Cock, J. H. (1971). *Laboratory manual for physiological studies of rice*.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/129048>