



The Morbidities Experienced during and after the 2017 Flood in Port Harcourt

A. S. Amaechi^{1*} and B. Ordinioha¹

¹Department of Preventive and Social Medicine, University of Port Harcourt, P.M.B. 5323 Choba, Port Harcourt, Nigeria.

Authors' contributions

This work was carried out in collaboration between both authors. Author ASA designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors ASA and BO managed the analyses of the study. Author ASA managed the literature searches. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRID/2019/v2i130093

Editor(s):

(1) Dr. Jamal Hussaini, Assistant Professor, Department of Medical Microbiology, Universiti Teknologi MARA, Malaysia.

Reviewers:

(1) Fasanmi Olubunmi Gabriel, University of Pretoria, South Africa.

(2) Soon Ae Kim, International Vaccine Institute, SNU Research Park, Nakseongdae-Dong, Kwanak-Gu, Seoul, South Korea.

(3) Alicia Garcia Falgueras, Spanish University, Spain.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/45450>

Original Research Article

Received 04 January 2019

Accepted 19 March 2019

Published 02 April 2019

ABSTRACT

Background: Floods are a type of natural phenomena, identified as the world most common natural disaster with a major ravaging impact claiming lives, causing property damage, destruction of environment and infrastructure, and increasing health impacts. Port Harcourt Metropolis, the capital of Rivers State was greatly affected by the 2017 flood. Hence this study aims to determine the morbidities experienced during and after the 2017 flood.

Materials and Methods: The study was a combination of quantitative and qualitative research for effective triangulation. A cross-sectional study was carried out. Data were collected through self-administered semi-structured questionnaires and key informant interviews. Data was analysed using Microsoft excel for editing, SPSS version 20 for quantitative analysis and Nvivo version 12.0 was used for qualitative analysis.

Results: A total of 210 respondents were administered questionnaires but had response rate of 96.67% and a total of 3 key informants were interviewed with a 100% response rate. Amongst these respondents were 44.8% male and 55.2% female, with respondents' mean age of

*Corresponding author: E-mail: sandycute.aa@gmail.com;

35.96±11.15. The frequencies of occurrence of morbidities of the residents and the 2017 flood were analyzed using percentage and chi-square test and the result showed a statistical significance ($p < 0.05$) between both variables.

Conclusion: The findings showed that the morbidities experienced during the flood had higher prevalence than post flood morbidities, but an exception was dark urine. These flood risks and morbidity outcomes can however be controlled through *adequate preventive measures and recommendable interventions*.

Keywords: Floods; morbidities; experiences and health.

1. INTRODUCTION

Floods stand as the most common and critical catastrophic event occurring in most parts of the world [1], which have resulted in loss of human lives and sources of livelihood, prolonged health impact, damage and deterioration of the environment, as well as retardation to development and economic losses [2,3]. Flooding is the most frequent global natural hazard, in terms of occurrence, with incidence and impact on the increase worldwide with a trend that is set to continue to increase in frequency and intensity due to climate change accompanied by rising sea levels and more frequent and extreme precipitation [4,5].

The Centre for Research on the Epidemiology of Disasters (CRED) defined flooding as “a significant rise of water level in a stream, lake, reservoir or coastal region” [6]. The occurrence of floods are influenced by natural phenomena and human involvement as the events and factors that leads to flood are diverse, multifaceted, and interrelated. Some of the factors that are attributed to the weather conditions include; heavy or sustained precipitation, snowmelts, or storm surges from cyclones while some of the important human factors include structural failures of dams, alteration of absorptive land cover with impervious surfaces and inadequate drainage systems [7].

In tropical regions, such as the Asia and the Pacific regions, flooding of high magnitude that has resulted in serious consequences has been caused by heavy rainstorms, hurricanes, snow melt and dam failures [8]. The United Kingdom National Risk Register of Civil Emergencies [7] puts the winter season of 2015/2016 as the second wettest winter on record as a series of storms (including ‘Desmond’ and ‘Eva’) resulted in heavy and sustained rainfall which resulted in flooding of about 17,600 UK properties and an

estimated £1.6 billion Economic damage. In Nigeria, the incident of floods is becoming a reoccurring decimal in most rural and urban areas leading to colossal loss of properties and lives [8]. This could be said to be evident in the two days of heavy down pour of rainfall in August 2008 in Benue state threw the residents of Makurdi out of their residences and their farmlands [7]. Also, the 2012 rainy season between August and September, has been the worst than previous years as it led to serious floods which inundated most part of the country [3]. By September 29 of the same year, the floods became so drastic that it affected over 134,371 people, displaced 64,473, injured 202 and killed 148; hence it was declared a national disaster. By the end of October, more than 7.7 million people had been affected by the floods, more than 2.1 registered as Internally Displaced People (IDP), and about 363 people were reported dead; almost 600,000 houses had been damaged, submerged or destroyed. Out of Nigeria's 36 states, 30 were affected by the floods [3]. The states affected with this bitter experience were those located within the Mangrove and Fresh water belts among which are Rivers, Lagos, Delta, and Bayelsa [3,7] In Port-Harcourt, which is the capital of Rivers State, the depth of flood water in affected areas has escalated significantly in the previous years due to the combined effects of uncoordinated, uncontrolled rapid urbanization, development of swamps, flood plains and poor drainage channels [8].

Port Harcourt Metropolis, which like most urban areas of the third world, has in most times experienced accelerated population growth that has led to changes in the land use activities. The city is faced with a number of environmental challenges, among which is flooding. Floods are major natural events that may not only lead to immediate loss of life and property but may have caused physical disability and severe psychological trauma among survivors. As a result of fears and actions taken to protect family

or belongings, experience of flooding and long-term uncertainties around insurance [9-11], often result in reduction in quality of life [12,13]. The IPCC in 2001 stated that the consequence of persistent rise in sea level and altered patterns of precipitation as a result of climate change are expected to increase the frequency and intensity of floods in many regions of the world. The incidence of flood disasters has been noted to globally increase as a result of various factors such as; population growth in areas at risk of flooding, climate change (which increases the variability and severity of weather, such as record-breaking rainfall and possibly more severe tropical cyclones) as well as changes to catchments (such as deforestation or urbanisation) that lead to increased run-off [14,15]. This thereby increases the impact of flood on health of the populace.

Taking a look at the 2017 flood in Port Harcourt during, one would likely want to ascertain if the health of the populace in the community was affected by the flood, thereby leading to morbidity (as the focus of this study), owing to the fact that morbidity in total is commonly defined as "departure from an overall state of health," but more specifically often referred to as the effect of illness, disease or injury in a population [16]. This paper hence aims to determine the morbidities experienced during and after the 2017 flood, so as to establish timely and adequate preventive measures and recommendable interventions to reduce the risks of flood and flood-related morbidity outcomes. Regarding the aim of this paper, the following questions and the answers that will be provided, forms the fundamentals and focus of this paper: What was the prevalence of morbidities experienced during and after the 2017 flood in Port Harcourt? Is there an association between the 2017 flood and the morbidities experienced in Port Harcourt?

2. METHODOLOGY

2.1 Study Location

This study was conducted within the metropolis of Port Harcourt, Rivers State, and South-South region of Nigeria. It is situated along the Bonny River and is located in the Niger Delta. As of 2016, the Port Harcourt urban area has an estimated population of 1,865,000 inhabitants, up from 1,382,592 as of 2006 [17].

2.2 Study Design

A descriptive, cross-sectional study design was employed in this study. In carrying out this study, the study populations were heads of households aged ≥18 years residing in the flood affected areas/quarters of the community.

2.3 Sample Size

The minimum sample size was derived using the Fisher's formula:

$$n = \frac{(Z^2) \times pq}{(d^2)} \quad [18]$$

Where: p = proportion of group p = 14.0% which was assumed because there is no similar study done so far. $p = 14 \div 100 = 0.14$; d = error margin = 5% = $(1 \times 5) \div 100 = 0.05$; z = corresponding value to C.I (z = 1.96); q = non-proportion of group = $1 - p = 1 - 0.14 = 0.86$. Thus, $n = (1.96)^2 (0.14) (0.86) / (0.05)^2 = 185$. Considering a 15% non-response rate = $15\% \times 185 = 27.75 = 185 + 27.75 = 212.75 = 210$ (2 s. f.); a final sample size of two hundred and ten (210) sample size was selected. A multistage sampling was conducted. The first stage was a clustered sampling of a centralized flood affected area. This made homogeneity and recruitment of sampling unit (houses) achievable. The second stage of sampling in this paper involved a systematic sampling of the sampling units which was achieved by deriving the sampling interval given as:
$$= \frac{\text{estimated number of houses in clustered streets}}{\text{allocated number of respondents}}$$

In the case where eligible participants were unavailable during the data collection, the next individual in charge of the house aged ≥18 years was administered the questionnaire and whereby there were non-available, the next immediate household was taken.

2.4 Data Collection

The research data was primarily generated through the use of both quantitative and qualitative research methods. For the quantitative method, the study instrument used was the semi-structure self-administered questionnaire. This was designed to ensure ease of answers, taking into cognizance, the differences in assimilation of various respondents. The questionnaire was made up of both close and unrestricted questions which is

grouped into sections, namely; socio-demographic characteristics, the 2017 flood experience, and the morbidity experiences (during and after the flood). In the qualitative method, the key informant interview was employed. The survey was supported with direct observation. During data collection, an official permission from the appropriate community leaders such as the community development committee chairman was first sorted for after presentation of ethical clearance, and then the selected respondents were enlightened with the study objectives before finally administering the questionnaires to the respondents and interviewing the community key informants. The quality of the data was assured by giving emphasis in designing the data collection tool, pre-testing the data and training the data collectors.

2.5 Statistical Analysis

After data collection, the obtained field data was entered into the computer and then edited using Microsoft Excel to ensure order of the information. After entering and editing, the Statistical Package for Social Sciences (SPSS) version 20 was used to analyse the quantitative data while Nvivo version 12.0 was used for qualitative analysis to get statistical results displayed using tables. Analysis of the output/data involved descriptive and inferential statistics. The descriptive statistics was done by

deriving mean, frequency, percentage and standard deviation of the data and the inferential statistics include chi-square test which was used to check if the differences that exist between variables are statistically significant. Meaningful conclusions from the study were drawn from these tests.

3. RESULTS

A total of 210 questionnaires were administered and 203 were sufficiently completed remaining 7 questionnaires, this gave a response rate of 96.67%.

The socio-demographic characteristics of the respondents that were interpreted (as seen in Table 1) include; sex, age, marital status and education. Under the sex composition of the respondents, a total of 55.2% female and 44.8% male were involved in the survey. The age distribution of the respondents in Table 1. showed that, majority of the respondents fell between the age of 26-35 years while the age group with the least participants' number were 66 and above, and the respondents mean age was 35.96±11.15. It was indicated that 116_(57%) were married, while 87_(43%) were unmarried. And based on the level of education of the respondents, majority of the respondents (55.2%) had secondary education while minority (3.4%) of the respondents had no education.

Table 1. Distribution of the socio-demographic characteristics of the respondents

Characteristics	Frequency (N = 203)	Percentage (%)
Sex		
Male	91	44.8
Female	112	55.2
Age (years)		
18 - 25	25	12.3
26 - 35	90	44.3
36 - 45	48	23.6
46 - 55	27	13.3
56 - 65	10	4.9
≥66	3	1.5
Mean ± SD	35.96 ± 11.15	
Marital Status		
Married	116	57.1
Unmarried	87	42.9
Educational Status		
None	7	3.4
Primary	17	8.4
Secondary	112	55.2
Tertiary	67	33.0

3.1 Quantitative Findings

3.1.1 Frequencies of occurrence of morbidities experienced

Table 2, shows the prevalence of morbidities experienced during and after the 2017 flood in Port Harcourt. 188(92.6%) indicated participants had morbidity experiences throughout the flood while 73(36.0%) indicated participants had morbidity experiences after the flood. Amongst those who had morbidity experiences during and after/post the 2017 flood, the prevalence of the observed symptoms during the flood which include fever (85.8%), shaking chills (64.9%), body pains (54.0%), limb weakness (36.4%), diarrhea (53.2), Rice-water stool (9.0%), blood in

stool (12.8%), fatigue (35.1%), nausea (32.4%), vomiting (34.6%), loss of appetite (56.6%), headache (63.3%), catarrh (68.9%), dry cough (38.6%), breathing difficulty (70.9%), sore throat (70.7%), rashes (59.8), yellow skin (19.7%) and yellow eyes (15.5%), were higher compared to the observed symptoms after the flood. An exception of a higher prevalence of observed symptoms during flood as compared to after the flood is dark urine. The prevalence of dark urine is higher (56.2%) compared to that of during the flood (31.9%). For experienced injuries such as bruise, fracture and cut, the prevalence during the flood were higher (26.1%, 18.6, 20.2% respectively) compared to the experienced injuries after the flood. The prevalence of the psychological morbidity during the 2017 flood

Table 2. Prevalence of the morbidity experiences of respondents during and post the 2017 floods in Port Harcourt

Question(s)	During		After	
	Yes (%)	No (%)	Yes (%)	No (%)
Did you experience any Morbidity	188 (92.6)	15 (7.4)	73 (36.0)	130 (64.0)
Morbidity Experienced				
Illnesses				
Fever	162 (85.8)	26 (14.2)	40 (54.8)	33 (45.2)
Shaking chills	122 (64.9)	66 (35.1)	35 (47.9)	38 (52.1)
Body pains	101 (54.0)	87 (46.0)	38 (52.1)	35 (47.9)
Limb weakness	69 (36.4)	119 (63.6)	23 (31.5)	50 (68.5)
Dark urine	60 (31.9)	128 (59.1)	41 (56.2)	32 (43.8)
Diarrhea	100 (53.2)	88 (46.8)	29 (39.7)	44 (60.3)
Rice-water stool	17 (9.0)	171 (91.0)	6 (8.2)	67 (91.8)
Blood in stool	24 (12.8)	164 (87.2)	9 (12.3)	64 (87.7)
Fatigue	66 (35.1)	122 (64.9)	20 (27.4)	53 (72.6)
Nausea	61 (32.4)	127 (67.6)	18 (25.0)	55 (75.0)
Vomiting	65 (34.6)	123 (65.4)	24 (32.9)	49 (67.1)
Loss of appetite	106 (56.6)	82 (43.4)	32 (43.8)	41 (56.2)
Headache	119 (63.3)	69 (36.7)	44 (60.3)	29 (39.7)
Catarrh	131 (68.9)	57 (31.1)	15 (20.5)	58 (79.5)
Dry cough	73 (38.6)	115 (61.4)	12 (16.4)	61 (83.6)
Breathing difficulty	133 (70.9)	55 (29.1)	18 (24.7)	55 (75.3)
Sore throat	133 (70.7)	55 (29.3)	19 (26.0)	54 (74.0)
Rashes	113 (59.8)	75 (40.2)	15 (20.5)	58 (79.5)
Yellow skin	37 (19.7)	151 (80.3)	08 (10.9)	65 (89.1)
Yellow eyes	29 (15.5)	159 (84.5)	08 (10.9)	65 (89.1)
Injuries				
Bruise	49 (26.1)	139 (73.9)	14 (19.1)	59 (80.9)
Fracture	35 (18.6)	153 (81.4)	11 (15.8)	62 (84.2)
Cut	38 (20.2)	150 (79.8)	09 (12.3)	64 (87.7)
Psychological effects				
Anxiety	154 (81.5)	34 (18.5)	40 (54.8)	33 (45.2)
Stress	162 (85.3)	26 (14.7)	62 (84.9)	11 (15.1)
Depression	151 (79.9)	37 (20.1)	45 (61.6)	28 (38.4)
Others (such as; chilblains, shock, sprain, bites, etc)	103 (54.5)	85 (45.5)	35 (47.9)	38 (52.1)

which include; anxiety (81.5%) and stress (85.3%), were higher compared to psychological morbidity prevalence after the flood. Unlike the prevalence of anxiety and stress, worried of loss (79.9%) during the flood has a lower prevalence as compared to after the 2017 flood. Other morbidity experiences such as shock, sprain/strain, foot sores, object pierce, chilblains and bites indicated by the respondents were higher (54.5%) in prevalence during the flood, compared with after the flood (47.9%).

Table 3, displays the chi-square test of association between the 2017 flood and the morbidities experienced in Port Harcourt. The result showed that the 2017 flood was significantly associated with the morbidity experiences of the respondents during and after the flood occurrence at p -value = 0.00 ($X^2 = 141.88$; 95% C.I: 12.26, 40.63). Under the observed symptoms, flooding was significantly associated with an increased number of fever cases during the flood which was 5.14 times higher than the fever case after the flood ($X^2 = 29.59$, p -value <0.05). For shaking chills, the odds amongst respondents with morbidity experiences during the flood were 2.01 times significantly higher than the cases after the flood, with a 95% C.I ranging from 1.16 to 3.47 ($X^2 = 6.30$, p -value <0.05). For cases of dark urine, the number after the flood had 0.37 times significant higher odds than the cases during the flood, with a 95% C.I ranging from 0.21 to 0.64 ($X^2 = 13.03$, p -value <0.05). For diarrhoea cases, the odds amongst respondents during the flood were 1.72 times significantly higher than the cases after the flood, with a 95% C.I ranging from 1.00 to 2.99 ($X^2 = 3.81$, p -value <0.05). Also respondents with cases of catarrh during the flood had 8.89 times significantly higher odds than the catarrh cases after the flood, with a displayed 95% C.I ranging from 4.65 to 16.98 ($X^2 = 51.50$, p -value <0.05). The odds for the cough cases during the floods were identified to be 3.23 times significantly higher than the cough cases after the flood, and the 95% C.I ranged from 1.63 to 6.40 ($X^2 = 12.00$, p -value <0.05). Based on the cases of breathing difficulty, a significantly higher odd of 3.79 were identified during the flood as compared to the cases following the flood ($X^2 = 45.80$, p -value <0.05).

Similar to the breathing difficulty cases, the odds of sore throat cases during the flood were 6.87 times significantly higher than the cases of sore throat after the flood and this showed a 95% C.I ranging between 3.73 and 12.65 ($X^2 = 43.23$, p -

value <0.05). Finally under the observed morbidity symptoms, the odd cases of rashes were 5.83 times significantly higher during the flood than after the flood with a 95% C.I ranging from 3.08 to 11.03 ($X^2 = 32.93$, p -value <0.05). Under the morbidity experienced injuries, Table 3 indicated that there was no statistical significance in the chi-square association between the 2017 flood and the morbidity experiences. Based on the psychological morbidity experiences of the respondents, Table 3 indicated a statistical significance of association between the 2017 flood and morbidity experiences (at p -value <0.05); where the odd cases of anxiety during the flood were 0.26 times significantly higher than the anxiety case after the flood ($X^2 = 6.72$, p -value <0.05); and the odd cases of respondents worried of loss during the flood were 0.17 times significantly higher than the cases after the flood with a 95% C.I ranging from 3.08 to 11.03 and $X^2 = 32.93$ (p -value <0.05). Other morbidity experiences (such as sprain/strain, chilblains, foot sores, bites and shock) identified by the respondents, were not statistically significant (as shown in Table 3), but the odd cases were 1.32 times higher during the flood than after the flood. Of these illnesses experienced, the odds ratio of some of them (including fever, shaking chills, dark urine, diarrhea, catarrh, cough, difficulty in breathing, sore throat and rashes) was significantly high (at p <0.05), indicating an association between floods and morbidity illnesses (Table 3).

3.1.2 Qualitative findings

The qualitative finding of this study is presented on Table 4. This was obtained after an interview (guided by the study objectives) with three (3) key informants in Port Harcourt.

The key informants that were interviewed identified several morbidities experienced during and after the 2017 flood. The morbidities experienced during the flood were water borne and vector borne diseases (11), physical injuries (1) while trying to evacuate, and social and psychological conditions (5). Few morbidity experiences were identified after the flood (3) such as high blood pressure.

The responsibilities assumed by the local authorities so as to manage the flood effects were recognized as intervention and advocacy (2), flood management strategies (2) such as encouraging affected residents evacuation and provision of relief materials, and flood prevention

Table 3. Chi-square test of association between the 2017 flood and the morbidity experiences

Response	Morbidity Experiences		Chi-square		Odds Ratio	95% C.I	
	During flood	After flood	X ²	P-value		Lower	Upper
Yes	188 (92.6)	73 (36.0)	141.88	0.00	22.32	12.26	40.63
No	15 (7.4)	130 (64.0)					
Morbidity Experienced	During flood	After flood	X ²	P-value	Odds Ratio	Lower	Upper
Illnesses							
Fever							
Yes	162 (85.8)	40 (54.8)	29.59	0.00	5.14	2.77	9.55
No	26 (14.2)	33 (45.2)					
Shaking chills							
Yes	122 (64.9)	35 (47.9)	6.30	0.01	2.01	1.16	3.47
No	66 (35.1)	38 (52.1)					
Body pains							
Yes	101 (54.0)	38 (52.1)	0.06	0.81	1.07	0.62	1.84
No	87 (46.0)	35 (47.9)					
Limb weakness							
Yes	69 (36.4)	23 (31.5)	0.62	0.43	1.26	0.71	2.24
No	119 (63.6)	50 (68.5)					
Dark urine							
Yes	60 (31.9)	41 (56.2)	13.03	0.00	0.37	0.21	0.64
No	128 (59.1)	32 (43.8)					
Diarrhea							
Yes	100 (53.2)	29 (39.7)	3.81	0.05	1.72	1.00	2.99
No	88 (46.8)	44 (60.3)					
Rice-water stool							
Yes	17 (9.0)	6 (8.2)	0.04	0.83	1.11	0.42	2.94
No	171 (91.0)	67 (91.8)					
Blood in stool							
Yes	24 (12.8)	9 (12.3)	0.01	0.92	1.04	0.46	2.36
No	164 (87.2)	64 (87.7)					
Fatigue							
Yes	66 (35.1)	20 (27.4)	1.41	0.23	1.43	0.79	2.60
No	122 (64.9)	53 (72.6)					
Nausea							
Yes	61 (32.4)	18 (25.0)	1.51	0.22	1.47	0.79	2.71
No	127 (67.6)	55 (75.0)					
Vomiting							
Yes	65 (34.6)	24 (32.9)	0.07	0.80	1.08	0.61	1.91
No	123 (65.4)	49 (67.1)					
Loss of appetite							
Yes	106 (56.6)	32 (43.8)	3.32	0.07	1.66	0.96	2.86
No	82 (43.4)	41 (56.2)					
Headache							
Yes	119 (63.3)	44 (60.3)	0.21	0.65	1.14	0.65	1.98
No	69 (36.7)	29 (39.7)					
Catarrh							
Yes	131 (68.9)	15 (20.5)	51.50	0.00	8.89	4.65	16.98
No	57 (31.1)	58 (79.5)					
Dry cough							
Yes	73 (38.6)	12 (16.4)	12.00	0.00	3.23	1.63	6.40
No	115 (61.4)	61 (83.6)					
Breathing difficulty							
Yes	133 (70.9)	18 (24.7)	45.80	0.00	7.39	3.98	13.71
No	55 (29.1)	55 (75.3)					

Sore throat							
Yes	133 (70.7)	19 (26.0)	43.23	0.00	6.87	3.73	12.65
No	55 (29.3)	54 (74.0)					
Rashes							
Yes	113 (59.8)	15 (20.5)	32.93	0.00	5.83	3.08	11.03
No	75 (40.2)	58 (79.5)					
Yellow skin							
Yes	37 (19.7)	08 (10.9)	2.80	0.09	1.99	0.88	4.51
No	151 (80.3)	65 (89.1)					
Yellow eyes							
Yes	29 (15.5)	08 (10.9)	0.86	0.35	1.48	0.64	3.41
No	159 (84.5)	65 (89.1)					
Injuries							
Bruise							
Yes	49 (26.1)	14 (19.1)	1.36	0.24	1.49	0.76	2.90
No	139 (73.9)	59 (80.9)					
Fracture							
Yes	35 (18.6)	11 (15.8)	0.46	0.50	1.29	0.62	2.70
No	153 (81.4)	62 (84.2)					
Cut							
Yes	38 (20.2)	09 (12.3)	2.21	0.14	1.80	0.82	3.94
No	150 (79.8)	64 (87.7)					
Psychological effects							
Anxiety							
Yes	154 (81.5)	40 (54.8)	6.72	0.01	0.26	0.09	0.77
No	34 (18.5)	33 (45.2)					
Stress							
Yes	162 (85.3)	62 (84.9)	0.07	0.80	1.11	0.52	2.37
No	26 (14.7)	11 (15.1)					
Depression							
Yes	151 (79.9)	45 (61.6)	9.82	0.00	0.17	0.05	0.59
No	37 (20.1)	28 (38.4)					
Others (shock, sprain, bites, etc)							
Yes	103 (54.5)	35 (47.9)	0.99	0.32	1.32	0.77	2.26
No	85 (45.5)	38 (52.1)					

strategies (2); such as encouraging settlements outside flood prone areas and education on waste discarding appropriateness.

The responsibilities that the Government, NGOs and other relevant bodies assumed in an attempt to curb the flood damages caused were noted as provision of relief materials (3); like food and mattresses, and public shelter (1).

The opinions/recommendations given by the key informants on how to control flood menace include; proper environmental management (3) to control environmental abuse (such as disposal of waste in drainages and building of structures at flood prone areas), improved attitudes and implementation of effective policies relating to flood (3), provision of effective and prompt flood warning systems against flood events (1), effective intervention or management of flood

proceedings (2) by government and other agencies/ institutions.

4. DISCUSSION

4.1 Morbidities Experienced

This study provides detailed morbidities experienced during and after the 2017 flood in Port Harcourt, which were majorly categorized into; illness, injuries and psychological effects (Tables 2 and 3), similar to several studies from Germany [18], England and Wales [19]. This study is in conformity with WHO, which reported that the health effects observed during and after floods include injuries, infections, and poisoning and greater mental-health problems [1]. Generally as revealed by this study, the prevalence of morbidities experienced amongst flood-affected respondents were significantly

higher (92.6%), during the flood as compared to after the flood, with increased odds of 22.32 and $X^2= 141.88$ (at $p<0.05$). Hence, this study rejects the null hypothesis (H_0) and retains the alternate hypothesis (H_1) which states that: there is an association between the 2017 flood and the morbidities experienced in Port Harcourt, at $p<0.05$. The findings of this study corroborate to the submissions of Landoh et al. and Carroll et al. [20,21].

4.1.1 Illness

A number of illnesses such as; fever, shaking chills, body pains, dark urine, diarrhea, limb weakness, Rice-water stool, blood in stool, fatigue, nausea, vomiting, headache, catarrh, cough; was identified in this study to have been experienced during and after the 2017 flood in Port Harcourt. The prevalence of these illnesses were noted to be higher during the flood

Table 4. Frequency, meaning and explanation of themes derived from key informant interview

Theme	Meaning	Categories	Frequency	Total	Evidence
Morbidity experiences	Diseases and other health related conditions suffered by the people of Eneka Community during the 2017 flood	Water and vector borne diseases/ conditions	11	20	The health of the people was greatly affected during the flood. So many had foot sores, rashes, and diarrhea. The children, especially those in families who remained, were very sick (Key informant 3).
		Physical injuries	1		
		Social and Psychological conditions	5		
		Post flood conditions	3		
		Evacuation/ Relocation	2		
Local Authorities	The role played by Local authorities in order to manage the effects of the flood	Intervention/ advocacy	2	6	They called on governments' attention for adequate construction of drainages and provision of relief supplies to the affected residents (Key informant 2).
		Flood management strategies	2		
		Flood prevention strategies	2		
Institutional assistance	The role played by Government, NGOs and other relevant bodies in an effort to control the flood caused damages	Relief materials	3	4	Some relief materials were sent by the government and NGOs (Key informant 2).
		Shelter	1		
Public opinion	Suggestions by the Community members and leaders on how to present and or manage flooding should it occur	Proper environmental management	3	9	Early warnings should be given about floods. Residents should avoid settling in flood prone localities (Key informant 1).
		Attitude/ Policies	3		
		Early warning	1		
		Intervention/ management	2		

(<http://www.sciencedomain.org/page/publication-charge>)

than after the 2017 flood and the illness with the highest frequency during the flood was fever (85.8%); but an exception of the prevalence of these illnesses is dark urine with a higher prevalence after flood event. These illnesses may be caused by different agents (such as viral, bacteria, fungi and protozoa), due to unhygienic water exposure which can occur through various routes of infections such as inhalation, ingestion and insect transmission. These illnesses may also have resulted from the poor sewage disposal and flood water depth. This study is similar to other studies like in Germany [18,22] and to the multicentre research of Obanga [23] in Ahoada East and Ahoada West Local government area. Also, fever which could be seen as a major symptom of malaria and other vector borne diseases was seen to be the highest occurring decimal according to the survey. This corroborates with the study carried out by Ahern and colleagues in 2005 which revealed that there is a potential for increased vector-borne illnesses and endemic levels of diarrheal disease, especially in areas with poor sanitation. Also, the 2012 study of Orijin on the flooding that occurred that same year in Rivers state also outlined fever and gastrointestinal disorders (cholera, dysentery and diarrhoea) as the most occurring morbidities, and attributed them to results from contaminated drinking-water and exposure to waste water facilities [24]. The findings this study were not too far from that of Obanga [23] when he studied the effects of flooding menace on health and housing in two communities of Ahoada east and west local government areas of Rivers state. Although his result showed that the morbidity with the highest prevalence was Cough (45%), it was closely followed by malaria/fever (44%), in supporting the outcome of the present study where fever (85.8%) was shown to have the highest frequency. The illnesses reported in this study are similar to certain general illnesses reported by US [19] and Germany [22], which were detailed as: respiratory illness, gastrointestinal illness, skin and eye irritation and infection.

Also, the respiratory illnesses namely: catarrh, cough, sore throat and difficulty in breathing were all reported by the findings of this study to be significantly associated to the 2017 flood (at $p < 0.05$). According to a study carried out by the Flood Hazard Research Centre (FHRC), in conformity to this study, reported chest infections, asthma, flu, coughs and colds to be due to the flooding in the North East of England

[19]. The gastrointestinal illnesses such as diarrhea, rice-water stool, blood in stool, nausea, vomiting, and loss of appetite are flood related illnesses commonly implicated by vector and water borne infections [25]. Amongst these gastrointestinal illnesses, the study reveals that the peak in diarrhea morbidity is associated with flooding. This is in conformity to several researches made by; Acuinjet et al., Wade et al. and Cann et al. [18,20,21]. The skin and eye illnesses include; yellow eyes, yellow skin and rashes which were all identified to have higher prevalence during the 2017 flood than post the flood. Amongst these illnesses, rashes was revealed to be significantly associated with the 2017 flood, at $p = 0.00$. This conforms to the study of Tunstall et al. and WHO [19,26].

4.1.2 Injuries

The relatively minor flood injuries that occurred during and post the 2017 flood include bruises and cuts; while some others indicated include sprain/strain, bites, foot sores, chilblains and object pierce. The more serious flood injury revealed was fracture, which was less experienced by a frequency of 18.6% during the flood and 15.8% after the flood. The injuries which according to Bich et al. [27] could be attributable to falls and clattering into some unobserved items beneath the water flooded areas during the flood and may have been sustained in the process of evacuation (while trying to remove themselves, family and valuables), while the post-flood injuries may have been sustained during the cleanup process, when the evacuated residents begin to return to their homes [28]. The prevalence of the injuries experienced during the flood were higher than the post-flood injuries; where the odds of bruise, cuts and fractures respectively were 1.49, 1.80 and 1.29 times respectively higher. Irrespective of the prevalence of the flood injuries, this study result revealed that the injuries experienced during and post the 2017 flood in Eneka community was not flood significant. This is similar to the CCASHH project in Europe that revealed no survey information on significant flood injuries [28]. In agreement with this study is the research conducted by the Health Protection Agency in London, which revealed that the significance of an injury will depend on the local hazards and type of flood [29].

4.1.3 Psychological effects

Living throughout a flood event according to Jermacane et al. [5], can be distressing and the

consequence on people's mental health can be profound. This forms the bases of several studies carried out on the common effects of flood on psychological disorders, amongst which are; the Psychosocial impact of the summer 2007 floods in England by Paranjothy et al. [30], and English National Cohort Study of Flooding and Health by Waite et al. [11]; which were no different from the findings from this paper, which also pointed towards stress, anxiety and depression as the common psychological disorders accompanying the flooding event. The findings of this study showed that the prevalence (Table 2) of the three most common psychological disorder (anxiety, stress and depression respectively), were higher during the flood (81.5%, 85.3% and 79.9% respectively), as compared to the psychological morbidities experienced after the flood (54.8%, 84.9% and 61.6% respectively), and the odds were significantly high (Table 3) showing an association between the 2017 flood and the psychological morbidities (at $p < 0.05$). These reported flood-common mental health outcome according to Tunstall et al. [19] could be attributed to certain flood vulnerability factors like the depth of flood, worried for loss, the strenuous evacuation process, contamination of water, less or warning system and recovery process. The prevalence of mental health disorder was higher amongst flood affected homes [29,30]. This present study also corroborates to the study of Carroll et al., where they conducted interviews with people who were affected by flood during the Carlisle floods and noted that many respondents spoke of psychological stress [21]. On the contrary, Udoimuk et al., researched flood-hazards influence on health in the State of Cross River [16]. Their study adopted a descriptive survey method and the result revealed that flood has no relative effect or wellbeing implication of those residing in such areas. This means that health implications and flood had no significant relationship.

5. CONCLUSION

The morbidities experienced during and after the 2017 floods in Port Harcourt were substantial and significant on the households and community, causing them to be physically injured, psychologically unstable and highly exposed to certain illnesses.

6. RECOMMENDATION

The primary, secondary and tertiary flood preventive measures are addressed with a range of recommendable interventions.

- **Primary Preventive Measures:** Organization of public enlightenment programmes, creating and implementing policies and legal status, Providing flooding insurance and Land use management.
- **Secondary Preventive Measure:** Effective flood forecasting and warning systems and Practical emergency planning measures.
- **Tertiary Preventive Measures:** Providing relief resources to flood victims, Adopting measures to track and ensure total wellbeing of affected populace and extension of emergency preparedness plan.

CONSENT

It is not applicable.

ETHICAL APPROVAL

During data collection, an official permission from the appropriate community leaders such as the community development committee chairman was first sorted for after presentation of ethical clearance.

ACKNOWLEDGEMENT

My profound appreciation goes to God Almighty for His infinite love, providence and guidance to achieve this final piece of academic exertion.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization. Flooding and Communicable Diseases Fact Sheet: The World Health Organization. Available: <http://www.who.int/hac/techguidance/ems/flood> (Accessed 2013)
2. Alderman K, Turner L, Tong S. Floods and human health: A systematic review. *Environment International*. 2012;47:37-47.
3. Wizer C, Week D. Geospatial mapping and analysis of the 2012 Nigeria flood disaster extent in Yenagoa City, Bayelsa State, Nigeria. *Journal of Environment and Earth Science*. 2014;4(10).
4. Ramin B, McMichael A. A climate change and health in Sub-Saharan Africa: A case-

- based perspective. *Eco Health*. 2014;6:52-57.
5. Jermacane D, Waite D, Beck R, Bone A, Amlôt R, Reacher M, Kovats S, Armstrong B, Leonardi G, Rubin J, Oliver I. The English national cohort study of flooding and health: The change in the prevalence of psychological morbidity at year two. *BMC Public Health*. 2018;18:1-8.
 6. Emergency Events Database. EM-DAT: The International disaster database. Retrieved; 2009. Available:www.emdat.be
 7. Hunt JC. Inland and coastal flooding: Developments in prediction and prevention. *Philosophical Transactions. Series A, Mathematical, Physical, and Engineering Sciences*. 2005;363(1831): 1475-1491.
 8. Jeb DN, Aggerwal SP. Flood inundation hazard modelling of the River Kaduna using remote sensing and geographic information systems. *Journal of Applied Sciences Research*. 2008;4(12):1822-1833.
 9. United Kingdom. National risk register of civil emergencies. Available:https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/644968/UK_National_Risk_Register_2017.pdf (Accessed on 12 January 2018)
 10. Dai W, Kaminga A, Tan H, Wang J, Lai Z, Wu X, Liu, A. Long-term psychological outcomes of flood survivors of hard-hit areas of the 1998 dongting lake flood in China: Prevalence and risk factors. *PLoS One*. 2017;12(2):e0171557.
 11. Waite T, Chaintarli K, Bec C, Bone A, Amlôt R, Kovats S, Armstrong B, Leonardi G, Rubin J, Oliver I. The English national cohort study of flooding and health: Cross-sectional analysis of mental health outcomes at year one. *BMC Public Health*. 2017;17:129.
 12. Haar J, Naderi S, Acerra R, Mathias M, Alagappan K. The livelihoods of Haitian health-care providers after the January 2010 earthquake: A pilot study of the economic and quality-of-life impact of emergency relief. *Int J Emerg Med*. 2012;5:13-8.
 13. Sudaryo M, Besral A, Rivany R, Phalkey R, Marx M, Guha-Sapir D. Injury, disability and quality of life after the 2009 earthquake in Padang, Indonesia: A prospective cohort study of adult survivors. *Glob Health Action*. 2012;5:576-7.
 14. Parker R, Little K, Heuser S. Development actions and the rising incidence of disasters (Evaluation Brief 4). Washington: World Bank; 2007.
 15. Cosgrave J. Responding to flood disasters: Learning from previous relief and recovery operations. ALNAP Lessons Paper. London: ALNAP/OD; 2014.
 16. DesMeules M, Turner L, Cho R. Morbidity experiences and disability among Canadian women. *BMC Women's Health*. 2004;4(1):S10.
 17. Arizona-Ogwu, Chinedu L. Port Harcourt PDP rally stampede: Irregular or deregulated police action? Nigerians in America (16 February 2011). Archived From The Original on 25 June; 2014. (Retrieved 25 June 2014)
 18. Schnitzler J, Benzler J, Altmann D, Mucke I, Krause G. Survey on the population's needs and the public health response during floods in Germany 2002. *J. Public Health Manag. Pract.* 2007;13:461-464.
 19. Tunstall S, Tapsell S, Green C, Floyd P, George C. The health effects of flooding: Social research results from England and Wales. *J. Water Health*. 2006;4:365-380.
 20. Landoh D, Tchamdja P, Saka B, Tint S, Gitta N, Wasswa P, Jager D. Morbidity and mortality due to malaria in Est mono District, Togo, from 2005 to 2010: A times series analysis. *Malar J*. 2012;11:389.
 21. Carroll B, Balogh R, Morbey H, Araoz G. Health and social impacts of a flood disaster: Responding to needs and implications for practice. *Disasters*. 2010;34(4):1045-1063.
 22. Steinfuhrer A, Kuhlicke C. Social vulnerability and the 2002 flood: Country report Germany (Mulde River). Flood Site Report T11-07-08; UFZ: Leipzig, Germany; 2007.
 23. Ogbanga MM. Impacts of flooding disaster on housing and health in two communities of Ahoada east and west local government areas of rivers state. *Nigerian Journal of Agriculture, Food and Environment*. 2015;11(1):44-50.
 24. Oriji CC. What to do about climate change-caused flooding and the associated diseases in rivers state of Nigeria. *Global Journal of Human-Social Science: B Geography, Geo-Sciences, Environmental*

- Science & Disaster Management. 2015;15(4):29–34.
25. Cann F, Thomas R, Salmon L, Wyn-Jones P, Kay D. Extreme water-related weather events and waterborne disease. *Epidemiol. Infect.* 2013;141:671–686.
26. World Health Organization. Risk assessment and preventive measures. Flooding and Communicable Diseases Fact Sheet: The World Health Organization. Available:<http://www.who.int/hac/techguidance/ems/flood> (Accessed 2006)
27. Bich TH, Quang LN, le TH, Hanh TT, Guha-Sapir D. Impacts of flood on health: Epidemiologic evidence from Hanoi, Vietnam. *Glob Health Action.* 2014;4: 6356.
28. World Health Organization. Environmental health in emergencies and disasters: A practical guide. (Accessed 2002)
29. Health Protection Agency. Interim guidance for health protection units. Surveillance Activity during Floods, London. Available:http://www.hpa.org.uk/web/HPAwebFile/HPAweb_C/1214291249686 (Accessed 4 August 2010)
30. Paranjothy S, Gallacher J, Amlot R. Psychosocial impact of the summer 2007 floods in England. *BMC Public Health.* 2011;11:145.

© 2019 Amaechi and Ordinioha; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/45450>*